
Communications

This section of the Human Papillomavirus database contains a list of supplemental references captured from the Molecular Biology subset of MEDLINE as communicated through GENINFO at the National Center for Biotechnology Information, National Library of Medicine, National Institutes of Health, Bethesda, Maryland. References for the years 1992, 1993, and 1994 are included.

The enclosed diskettes comprise the electronic portion of Part V, providing HPV and animal papillomavirus nucleic acid and protein sequences in ASCII characters. As a general rule, American and non-European users receive nucleic acid files in GenBank format, which includes protein translations of coding sequences. European users receive sequence files in EMBL and Swiss-Prot formats. Both PC-DOS and Apple Macintosh formatted diskettes are available; which format you receive depends upon what you have requested. If you wish to change the diskette format or the file formats, simply contact us at 505-665-1356, fax a message to us at 505-665-3493, or contact the database by e-mail at `cxc@t10.lanl.gov`. If you wish to contact us via the postal service, our address is: HPV Sequence Database, T-10, MS K710, LANL, Los Alamos, NM, 87545, U.S.A.

Sequences and alignments are also available electronically from a server here at Los Alamos called "atlas". In the following instructions for accessing atlas, what you actually type at your terminal is shown in **boldface** type [with an explanation shown in square brackets]. What the server responds with is shown indented in *italic* type.

ftp atlas.lanl.gov [this ftp command connects your computer to atlas]

Name (atlas.lanl.gov): [atlas asks for your user name]

anonymous [for offsite users anonymous is the only name that works]

Guest login ok, send ident as password.

Password:

[enter your e-mail address or your name as ident]

Guest login ok. access restrictions apply.

cd pub [change to the directory called pub]

CWD command successful

cd papilloma

ls [this lists the files and directories inside of papilloma]

SWISS-PROT-files, RoadMap, Alignments, ReadMe, EMBL-files, GenBank-files

pwd [to show your location in the file hierarchy]

"/pub/papilloma" is current directory.

cd GenBank-files/Human-papilloma [move down 2 directories]

ls [all the GenBank files are listed]

get HPV47.gb [this copies the file HPV47.gb to your home computer]

bye [this disconnects you from the server]

A map of the HPV directories and files on atlas is reproduced on the following page. Many of the directories contain ReadMe files that describe the contents.

Instructions for Accessing ftp Server "atlas"


In the following instructions, what you actually type at your terminal is shown in **boldface** type [with an explanation shown in square brackets]. What the server responds with is shown indented in *italic* type.

```

ftp atlas.lanl.gov
  Name (atlas.lanl.gov):
anonymous
  Guest login ok, send ident as password.
  Password:
[enter your e-mail address as ident]
  Guest login ok, access restrictions apply.
cd pub [change to the directory called pub]
  CWD command successful
cd papilloma
ls [this lists the files and directories inside of papilloma]
  SWISS-PROT-files, RoadMap, Alignments, ReadMe, EMBL-files, GenBank-files
pwd [to show your location in the file hierarchy]
  "/pub/papilloma" is current directory.
cd GenBank-files/Human-papilloma [move down 2 directories]
ls
  [all the GenBank files are listed]
get HPV47.gb [this copies the file HPV47.gb to your home computer]
bye [this disconnects you from the server]
  
```

JUNE 1994* MAP OF HPV DATABASE ON "ATLAS"

* Data subject to continuous revision



- Agrawal R; Pelkonen J; Mantyjarvi RA
Bovine papillomavirus type 1-transformed primary mouse fibroblasts show no correlation between tumorigenicity and viral gene expression, but c-myc gene expression is elevated in tumorigenic cell lines.
J Gen Virol 73 (Pt 6): 1527-32 (1992)
- Agrawal R; Pelkonen J; Rytkonen M; Mantyjarvi RA
Integration of bovine papillomavirus type 1 DNA and analysis of the amplified virus-cell junctions in transformed primary mouse fibroblasts.
J Gen Virol 73 (Pt 1): 201-6 (1992)
- Alcantara Vazquez A; Hernandez Gonzalez M; Amancio Chasin O; Lira Garces TJ
[Behavior of nucleolar organizers in the normal cervix, human papilloma virus infection and cervico-uterine cancer]
Ginecol Obstet Mex 60: 286-9 (1992)
- Alderborn A; Jareborg N; Burnett S
Evidence that the transcriptional trans-activating function of the bovine papillomavirus type 1 E2 gene is not required for viral DNA amplification in division-arrested cells.
J Gen Virol 73 (Pt 10): 2639-51 (1992)
- Altmann A; Jochmus-Kudielka I; Frank R; Gausepohl H; Moebius U; Gissmann L; Meuer SC
Definition of immunogenic determinants of the human papillomavirus type 16 nucleoprotein E7.
Eur J Cancer 28: 326-33 (1992)
- Angell MG; Christensen ND; Kreider JW
An in vitro system for studying the initial stages of cottontail rabbit papillomavirus infection.
J Virol Methods 39: 207-16 (1992)
- Anwar K; Nakakuki K; Shiraishi T; Naiki H; Yatani R; Inuzuka M
Presence of ras oncogene mutations and human papillomavirus DNA in human prostate carcinomas.
Cancer Res 52: 5991-6 (1992)
- Armstrong DJ; Roman A
Mutagenesis of human papillomavirus types 6 and 16 E7 open reading frames alters the electrophoretic mobility of the expressed proteins.
J Gen Virol 73 (Pt 5): 1275-9 (1992)
- Barsoum J; Prakash SS; Han P; Androphy EJ
Mechanism of action of the papillomavirus E2 repressor: repression in the absence of DNA binding.
J Virol 66: 3941-5 (1992)
- Bartsch D; Boye B; Baust C; zur Hausen H; Schwarz E
Retinoic acid-mediated repression of human papillomavirus 18 transcription and different ligand regulation of the retinoic acid receptor beta gene in non-tumorigenic and tumorigenic HeLa hybrid cells.
EMBO J 11: 2283-91 (1992)
- Batova A; Danielpour D; Pirisi L; Creek KE
Retinoic acid induces secretion of latent transforming growth factor beta 1 and beta 2 in normal and human papillomavirus type 16-immortalized human keratinocytes.
Cell Growth Differ 3: 763-72 (1992)
- Bauknecht T; Angel P; Royer HD; zur Hausen H
Identification of a negative regulatory domain in the human papillomavirus type 18 promoter: interaction with the transcriptional repressor YY1.
EMBO J 11: 4607-17 (1992)
- Belaguli NS; Pater MM; Pater A
Nucleotide 880 splice donor site required for efficient transformation and RNA accumulation by human papillomavirus type 16 E7 gene.
J Virol 66: 2724-30 (1992)
- Binnering I; Clavel C; Polette M; Bouterin MC; Chypre C; Alpha B; Birembaut P
A new nonisotopic detection of human papillomavirus DNA using polymerase chain reaction.
Diagn Mol Pathol 1: 239-45 (1992)
- Blangy A; Carle GF; Pierrefite V; Rassoulzadegan M; Cuzin F
Mammalian and viral DNA sequences which interfere with the maintenance of a centromeric vector in yeast.
Biochem Biophys Res Commun 187: 737-43 (1992)
- Blanton RA; Coltrera MD; Gown AM; Halbert CL; McDougall JK
Expression of the HPV16 E7 gene generates proliferation in stratified squamous cell cultures which is independent of endogenous p53 levels.
Cell Growth Differ 3: 791-802 (1992)
- Bossy-Wetzel E; Bravo R; Hanahan D
Transcription factors junB and c-jun are selectively up-regulated and functionally implicated in fibrosarcoma development.
Genes Dev 6: 2340-51 (1992)
- Bowden PE; Woodworth CD; Doniger J; DiPaolo JA
Down-regulation of keratin 14 gene expression after v-Har-ras transfection of human papillomavirus-immortalized human cervical epithelial cells.
Cancer Res 52: 5865-71 (1992)

References (1992)

- Brandsma JL; Yang ZH; DiMaio D; Barthold SW; Johnson E; Xiao W
The putative E5 open reading frame of cottontail rabbit papillomavirus is dispensable for papilloma formation in domestic rabbits.
J Virol 66: 6204-7 (1992)
- Bratthauer GL; Tavassoli FA; O'Leary TJ
Etiology of breast carcinoma: no apparent role for papillomavirus types 6/11/16/18.
Pathol Res Pract 188: 384-6 (1992)
- Braun L; Durst M; Mikumo R; Crowley A; Robinson M
Regulation of growth and gene expression in human papillomavirus-transformed keratinocytes by transforming growth factor-beta: implications for the control of papillomavirus infection.
Mol Carcinog 6: 100-11 (1992)
- Bream GL; Vaillancourt P; Botchan MR
A constitutive enhancer in the bovine papillomavirus upstream regulatory region shares genetic elements with the viral P1 promoter.
J Virol 66: 7319-27 (1992)
- Burnett AF; Barnes WA; Johnson JC; Grendys E; Willett GD; Barter JF; Doniger J
Prognostic significance of polymerase chain reaction detected human papillomavirus of tumors and lymph nodes in surgically treated stage IB cervical cancer.
Gynecol Oncol 47: 343-7 (1992)
- Burnett S; Jareborg N; DiMaio D
Localization of bovine papillomavirus type 1 E5 protein to transformed basal keratinocytes and permissive differentiated cells in fibropapilloma tissue.
Proc Natl Acad Sci U S A 89: 5665-9 (1992)
- Callahan DE; Karim A; Zheng G; Ts'o PO; Lesko SA
Quantitation and mapping of integrated human papillomavirus on human metaphase chromosomes using a fluorescence microscope imaging system.
Cytometry 13: 453-61 (1992)
- Campo MS
Cell transformation by animal papillomaviruses.
J Gen Virol 73 (Pt 2): 217-22 (1992)
- Cardillo MR
Nucleolar organizing regions in human papillomavirus infection and in cervical intraepithelial neoplasia.
Eur J Gynaecol Oncol 13: 277-80 (1992)
- Cason J; Kambo PK; Best JM; McCance DJ
Detection of antibodies to a linear epitope on the major coat protein (L1) of human papillomavirus type-16 (HPV-16) in sera from patients with cervical intraepithelial neoplasia and children.
Int J Cancer 50: 349-55 (1992)
- Chan SY; Bernard HU; Ong CK; Chan SP; Hofmann B; Delius H
Phylogenetic analysis of 48 papillomavirus types and 28 subtypes and variants: a showcase for the molecular evolution of DNA viruses.
J Virol 66: 5714-25 (1992)
- Chan SY; Ho L; Ong CK; Chow V; Drescher B; Durst M; ter Meulen J; Villa L; Luande J; Mgaya HN; et al
Molecular variants of human papillomavirus type 16 from four continents suggest ancient pandemic spread of the virus and its coevolution with humankind.
J Virol 66: 2057-66 (1992)
- Chang YL; Lin CY; Tseng CJ; Cheng HS; Lin HC; Pao CC
Prevalence of genital human papillomavirus infections in patients at a sexually transmitted diseases clinic.
Eur J Clin Microbiol Infect Dis 11: 454-7 (1992)
- Chang YS; Lin YJ; Tsai CN; Shu CH; Tsai MS; Choo KB; Liu ST
Detection of mutations in the p53 gene in human head and neck carcinomas by single strand conformation polymorphism analysis.
Cancer Lett 67: 167-74 (1992)
- Chapman WB; Lorincz AT; Willett GD; Wright VC; Kurman RJ
Epidermal growth factor receptor expression and the presence of human papillomavirus in cervical squamous intraepithelial lesions.
Int J Gynecol Pathol 11: 221-6 (1992)
- Chellappan S; Kraus VB; Kroger B; Munger K; Howley PM; Phelps WC; Nevins JR
Adenovirus E1A, simian virus 40 tumor antigen, and human papillomavirus E7 protein share the capacity to disrupt the interaction between transcription factor E2F and the retinoblastoma gene product.
Proc Natl Acad Sci U S A 89: 4549-53 (1992)
- Chen L; Ashe S; Brady WA; Hellstrom I; Hellstrom KE; Ledbetter JA; McGowan P; Linsley PS
Costimulation of antitumor immunity by the B7 counter-receptor for the T lymphocyte molecules CD28 and CTLA-4.
Cell 71: 1093-102 (1992)

- Chen L; Mizuno MT; Singhal MC; Hu SL; Galloway DA; Hellstrom I; Hellstrom KE
Induction of cytotoxic T lymphocytes specific for a syngeneic tumor expressing the E6 oncoprotein of human papillomavirus type 16.
J Immunol 148: 2617-21 (1992)
- Chen TM; Defendi V
Functional interaction of p53 with HPV18 E6, c-myc and H-ras in 3T3 cells.
Oncogene 7: 1541-7 (1992)
- Chetsanga C; Malmstrom PU; Gyllensten U; Moreno-Lopez J; Dinter Z; Pettersson U
Low incidence of human papillomavirus type 16 DNA in bladder tumor detected by the polymerase chain reaction.
Cancer 69: 1208-11 (1992)
- Chiang CM; Broker TR; Chow LT
Properties of bovine papillomavirus E1 mutants.
Virology 191: 964-7 (1992)
- Chiang CM; Dong G; Broker TR; Chow LT
Control of human papillomavirus type 11 origin of replication by the E2 family of transcription regulatory proteins.
J Virol 66: 5224-31 (1992)
- Chiang CM; Ustav M; Stenlund A; Ho TF; Broker TR; Chow LT
Viral E1 and E2 proteins support replication of homologous and heterologous papillomaviral origins.
Proc Natl Acad Sci U S A 89: 5799-803 (1992)
- Chiang LC; Chiang W; Chang SF; Chen HY
Spontaneous production of growth factors for human lymphocytes from a human papillomavirus type 18-contained foreskin fibroblast cell line.
Kao Hsiung I Hsueh Ko Hsueh Tsa Chih 8: 121-6 (1992)
- Chiang LC; Chiang W; Chang SF; Chen HY; Yu HS
Characterization of an immortalized human cell line derived from neonatal foreskin diploid fibroblasts.
J Dermatol 19: 1-11 (1992)
- Chinami M; Shingu M
Construction of an expression vector for human papillomavirus type 16 E7-glutathione S transferase fusion protein.
Kurume Med J 39: 9-12 (1992)
- Choo KB; Chong KY; Liew LN; Hsu HC; Cheng WT
Unregulated and basal transcriptional activities of the regulatory sequence of the type 18 human papillomavirus genome in transgenic mice.
Virology 188: 378-83 (1992)
- Choudhury S; Woodworth CD; Inamdar A; el-Beik T; DiPaolo JA; Rosenthal LJ
Differences in retention and expression of transfected human cytomegalovirus Towne XbaI-E transforming fragment in human cervical and NIH 3T3 lines.
Intervirology 33: 187-96 (1992)
- Christensen ND; Kreider JW; Shah KV; Rando RF
Detection of human serum antibodies that neutralize infectious human papillomavirus type 11 virions.
J Gen Virol 73 (Pt 5): 1261-7 (1992)
- Chung GT; Huang DP; Lo KW; Chan MK; Wong FW
Genetic lesion in the carcinogenesis of cervical cancer.
Anticancer Res 12: 1485-90 (1992)
- Ciocca DR; Lo Castro G; Alonio LV; Cobo MF; Lotfi H; Teysie A
Effect of human papillomavirus infection on estrogen receptor and heat shock protein hsp27 phenotype in human cervix and vagina.
Int J Gynecol Pathol 11: 113-21 (1992)
- Claas EC; Melchers WJ; Niesters HG; van Muyden R; Stolz E; Quint WG
Infections of the cervix uteri with human papillomavirus and Chlamydia trachomatis.
J Med Virol 37: 54-7 (1992)
- Claas EC; Quint WG; Pieters WJ; Burger MP; Oosterhuis WJ; Lindeman J
Human papillomavirus and the three group metaphase figure as markers of an increased risk for the development of cervical carcinoma.
Am J Pathol 140: 497-502 (1992)
- Clavel C; Zerat L; Binninger I; Bouterin MC; Polette M; Monsonogo J; Birembaut P
DNA content measurement and in situ hybridization in condylomatous cervical lesions.
Diagn Mol Pathol 1: 180-4 (1992)
- Cone RW; Minson AC; Smith MR; McDougall JK
Conservation of HPV-16 E6/E7 ORF sequences in a cervical carcinoma.
J Med Virol 37: 99-107 (1992)
- Cooper K; Herrington CS; Lo ES; Evans MF; McGee JO
Integration of human papillomavirus types 16 and 18 in cervical adenocarcinoma.
J Clin Pathol 45: 382-4 (1992)
- Coulson J; Malcolm AD
Antisense oligonucleotides as antiviral agents.
Ann N Y Acad Sci 660: 339-41 (1992)

References (1992)

- Crook T; Vousden KH
Properties of p53 mutations detected in primary and secondary cervical cancers suggest mechanisms of metastasis and involvement of environmental carcinogens.
EMBO J 11: 3935-40 (1992)
- Crook T; Wrede D; Tidy JA; Mason WP; Evans DJ; Vousden KH
Clonal p53 mutation in primary cervical cancer: association with human-papillomavirus-negative tumours.
Lancet 339: 1070-3 (1992)
- Czegledy J; Poka R; Veress G; Gergely L
Amplification of human papillomavirus type 16 transforming genes from cervical cancer biopsies and lymph nodes of Hungarian patients.
J Clin Microbiol 30: 233-6 (1992)
- Czegledy J; Rogo KO; Evander M; Wadell G
High-risk human papillomavirus types in cytologically normal cervical scrapes from Kenya.
Med Microbiol Immunol (Berl) 180: 321-6 (1992)
- D'Amato L; Pilotti S; Rotola A; Di Luca D; Cassai E; Rilke F
Comparison of Southern blot analysis with isotopic and nonisotopic in situ hybridization for the detection of human papillomavirus sequences in invasive carcinoma of the uterine cervix.
Mod Pathol 5: 207-11 (1992)
- Daniels-McQueen S; Goessling LS; Thach RE
Inducible expression bovine papillomavirus shuttle vectors containing ferritin translational regulatory elements.
Gene 122: 271-9 (1992)
- Das BC; Gopalkrishna V; Sharma JK; Roy M; Luthra UK
Human papillomavirus DNA in urine of women with pre-neoplastic and neoplastic cervical lesions [letter]
Lancet 340: 1417-8 (1992)
- Das BC; Sharma JK; Gopalakrishna V; Luthra UK
Analysis by polymerase chain reaction of the physical state of human papillomavirus type 16 DNA in cervical preneoplastic and neoplastic lesions.
J Gen Virol 73 (Pt 9): 2327-36 (1992)
- Das BC; Sharma JK; Gopalkrishna V; Das DK; Singh V; Gissmann L; zur Hausen H; Luthra UK
A high frequency of human papillomavirus DNA sequences in cervical carcinomas of Indian women as revealed by Southern blot hybridization and polymerase chain reaction.
J Med Virol 36: 239-45 (1992)
- Dasgupta S; Zabielski J; Simonsson M; Burnett S
Rolling-circle replication of a high-copy BPV-1 plasmid.
J Mol Biol 228: 1-6 (1992)
- Davies RC; Vousden KH
Functional analysis of human papillomavirus type 16 E7 by complementation with adenovirus E1A mutants.
J Gen Virol 73 (Pt 8): 2135-9 (1992)
- De Braekeleer M; Sreekantaiah C; Haas O
Herpes simplex virus and human papillomavirus sites correlate with chromosomal breakpoints in human cervical carcinoma.
Cancer Genet Cytogenet 59: 135-7 (1992)
- Del Vecchio AM; Romanczuk H; Howley PM; Baker CC
Transient replication of human papillomavirus DNAs.
J Virol 66: 5949-58 (1992)
- Della Torre G; Donghi R; Longoni A; Pilotti S; Pasquini G; De Palo G; Pierotti MA; Rilke F; Della Porta G
HPV DNA in intraepithelial neoplasia and carcinoma of the vulva and penis.
Diagn Mol Pathol 1: 25-30 (1992)
- Della Torre G; Donghi R; de Campos Lima PO; Pasquini G; Pilotti S; Koronel R; Pierotti MA; De Palo G; Della Porta G; Rilke F
Human papillomavirus genomes in male urethral cells.
Am J Pathol 141: 1181-6 (1992)
- Demeter LM; Stoler MH; Sobel ME; Broker TR; Chow LT
Expression of high-affinity laminin receptor mRNA correlates with cell proliferation rather than invasion in human papillomavirus-associated cervical neoplasms.
Cancer Res 52: 1561-7 (1992)
- Desaintes C; Hallez S; Van Alphen P; Burny A
Transcriptional activation of several heterologous promoters by the E6 protein of human papillomavirus type 16.
J Virol 66: 325-33 (1992)
- Dexeus S; Rubio R; Bassols G; Jakob D; Ojeda J; Labastida R
Papilloma virus infection. Precancer and epidermoid cancer.
Eur J Gynaecol Oncol 13: 167-76 (1992)
- DiLorenzo TP; Tamsen A; Abramson AL; Steinberg BM
Human papillomavirus type 6a DNA in the lung carcinoma of a patient with recurrent laryngeal papillomatosis is characterized by a partial duplication.
J Gen Virol 73 (Pt 2): 423-8 (1992)

- Dickens P; Srivastava G; Liu YT
Human papillomavirus 16/18 and nasopharyngeal carcinoma.
J Clin Pathol 45: 81-2 (1992)
- Dollard SC; Wilson JL; Demeter LM; Bonnez W; Reichman RC; Broker TR; Chow LT
Production of human papillomavirus and modulation of the infectious program in epithelial raft cultures. OFF
Genes Dev 6: 1131-42 (1992)
- Doorbar J; Ely S; Coleman N; Hibma M; Davies DH; Crawford L
Epitope-mapped monoclonal antibodies against the HPV16 E1-E4 protein.
Virology 187: 353-9 (1992)
- Durst M; Glitz D; Schneider A; zur Hausen H
Human papillomavirus type 16 (HPV 16) gene expression and DNA replication in cervical neoplasia: analysis by in situ hybridization.
Virology 189: 132-40 (1992)
- Dyson N; Guida P; Munger K; Harlow E
Homologous sequences in adenovirus E1A and human papillomavirus E7 proteins mediate interaction with the same set of cellular proteins.
J Virol 66: 6893-902 (1992)
- Edwards GM; Huber HE; DeFeo-Jones D; Vuocolo G; Goodhart PJ; Maigetter RZ; Sanyal G; Oliff A; Heimbrook DC
Purification and characterization of a functionally homogeneous 60-kDa species of the retinoblastoma gene product.
J Biol Chem 267: 7971-4 (1992)
- Effert PJ; Frye RA; Neubauer A; Liu ET; Walther PJ
Human papillomavirus types 16 and 18 are not involved in human prostate carcinogenesis: analysis of archival human prostate cancer specimens by differential polymerase chain reaction.
J Urol 147: 192-6 (1992)
- Eschle D; Durst M; ter Meulen J; Luande J; Eberhardt HC; Pawlita M; Gissmann L
Geographical dependence of sequence variation in the E7 gene of human papillomavirus type 16.
J Gen Virol 73 (Pt 7): 1829-32 (1992)
- Evander M; Edlund K; Boden E; Gustafsson A; Jonsson M; Karlsson R; Rylander E; Wadell G
Comparison of a one-step and a two-step polymerase chain reaction with degenerate general primers in a population-based study of human papillomavirus infection in young Swedish women.
J Clin Microbiol 30: 987-92 (1992)
- Finerty S; Tarlton J; Mackett M; Conway M; Arrand JR; Watkins PE; Morgan AJ
Protective immunization against Epstein-Barr virus-induced disease in cottontop tamarins using the virus envelope glycoprotein gp340 produced from a bovine papillomavirus expression vector.
J Gen Virol 73 (Pt 2): 449-53 (1992)
- Fujita M; Inoue M; Tanizawa O; Iwamoto S; Enomoto T
Alterations of the p53 gene in human primary cervical carcinoma with and without human papillomavirus infection.
Cancer Res 52: 5323-8 (1992)
- Furuta Y; Takasu T; Asai T; Shinohara T; Sawa H; Nagashima K; Inuyama Y
Detection of human papillomavirus DNA in carcinomas of the nasal cavities and paranasal sinuses by polymerase chain reaction.
Cancer 69: 353-7 (1992)
- Fuse A; Simizu B; Van Ranst M; Opdenakker G; Van Damme J
The induction of IL-6 and gelatinase B by IL-1 in mouse cell lines transformed with bovine papillomavirus: decreased production in tumorigenic cells.
Lymphokine Cytokine Res 11: 215-20 (1992)
- Galehouse D; Jenison E; DeLucia A
Differences in the integration pattern and episomal forms of human papillomavirus type 16 DNA found within an invasive cervical neoplasm and its metastasis.
Virology 186: 339-41 (1992)
- Georges E; Pehau-Arnaudet G; Orth G
Molecular and biological characterization of cottontail rabbit papillomavirus variant DNA sequences integrated in the VX7 carcinoma.
Virology 186: 750-9 (1992)
- Ghim SJ; Jenson AB; Schlegel R
HPV-1 L1 protein expressed in cos cells displays conformational epitopes found on intact virions.
Virology 190: 548-52 (1992)
- Glew SS; Duggan-Keen M; Cabrera T; Stern PL
HLA class II antigen expression in human papillomavirus-associated cervical cancer.
Cancer Res 52: 4009-16 (1992)
- Goldsborough MD; McAllister P; Reid R; Temple G; Lorincz AT
A comparison study of human papillomavirus prevalence by the polymerase chain reaction in low risk women and in a gynaecology referral group at elevated risk for cervical cancer.
Mol Cell Probes 6: 451-7 (1992)

References (1992)

- Goldstein DJ; Andresson T; Sparkowski JJ; Schlegel R
The BPV-1 E5 protein, the 16 kDa membrane pore-forming protein and the PDGF receptor exist in a complex that is dependent on hydrophobic transmembrane interactions.
EMBO J 11: 4851-9 (1992)
- Goldstein DJ; Kulke R; Dimaio D; Schlegel R
A glutamine residue in the membrane-associating domain of the bovine papillomavirus type 1 E5 oncoprotein mediates its binding to a transmembrane component of the vacuolar H(+)-ATPase.
J Virol 66: 405-13 (1992)
- Goldstein DJ; Toyama R; Dhar R; Schlegel R
The BPV-1 E5 oncoprotein expressed in *Schizosaccharomyces pombe* exhibits normal biochemical properties and binds to the endogenous 16-kDa component of the vacuolar proton-ATPase.
Virology 190: 889-93 (1992)
- Gopalkrishna V; Francis A; Sharma JK; Das BC
A simple and rapid method of high quantity DNA isolation from cervical scrapes for detection of human papillomavirus infection.
J Virol Methods 36: 63-72 (1992)
- Guido MC; Zamorano R; Garrido-Guerrero E; Gariglio P; Garcia-Carranca A
Early promoters of genital and cutaneous human papillomaviruses are differentially regulated by the bovine papillomavirus type 1 E2 gene product.
J Gen Virol 73 (Pt 6): 1395-400 (1992)
- Habiger C; Stelzer G; Schwarz U; Winnacker EL
Two cellular single-strand-specific DNA-binding proteins interact with two regions of the bovine papillomavirus type 1 genome, including the origin of DNA replication.
J Virol 66: 5988-98 (1992)
- Halbert CL; Demers GW; Galloway DA
The E6 and E7 genes of human papillomavirus type 6 have weak immortalizing activity in human epithelial cells.
J Virol 66: 2125-34 (1992)
- Han R; Breitburd F; Marche PN; Orth G
Linkage of regression and malignant conversion of rabbit viral papillomas to MHC class II genes [see comments]
Nature 356: 66-8 (1992)
- Hartman DA; Kuo SR; Broker TR; Chow LT; Wells RD
Intermolecular triplex formation distorts the DNA duplex in the regulatory region of human papillomavirus type-11.
J Biol Chem 267: 5488-94 (1992)
- Heck DV; Yee CL; Howley PM; Munger K
Efficiency of binding the retinoblastoma protein correlates with the transforming capacity of the E7 oncoproteins of the human papillomaviruses.
Proc Natl Acad Sci U S A 89: 4442-6 (1992)
- Hegde RS; Grossman SR; Laimins LA; Sigler PB
Crystal structure at 1.7 Å of the bovine papillomavirus-1 E2 DNA-binding domain bound to its DNA target [see comments]
Nature 359: 505-12 (1992)
- Hermonat PL
Inhibition of bovine papillomavirus plasmid DNA replication by adeno-associated virus.
Virology 189: 329-33 (1992)
- Hermonat PL; Daniel RW; Shah KV
The spermicide nonoxynol-9 does not inactivate papillomavirus.
Sex Transm Dis 19: 203-5 (1992)
- Hietanen S; Auvinen E; Grenman S; Lakkala T; Sajantila A; Klemi P; Maenpaa J
Isolation of two keratinocyte cell lines derived from HPV-positive dysplastic vaginal lesions.
Int J Cancer 52: 391-8 (1992)
- Higgins GD; Phillips GE; Smith LA; Uzelin DM; Burrell CJ
High prevalence of human papillomavirus transcripts in all grades of cervical intraepithelial glandular neoplasia.
Cancer 70: 136-46 (1992)
- Higgins GD; Uzelin DM; Phillips GE; McEvoy P; Marin R; Burrell CJ
Transcription patterns of human papillomavirus type 16 in genital intraepithelial neoplasia: evidence for promoter usage within the E7 open reading frame during epithelial differentiation.
J Gen Virol 73 (Pt 8): 2047-57 (1992)
- Hoppe-Seyler F; Butz K
Activation of human papillomavirus type 18 E6-E7 oncogene expression by transcription factor Sp1.
Nucleic Acids Res 20: 6701-6 (1992)
- Hording U; Daugaard S; Iversen AK; Bock JE; Philip J
Detection of human papillomavirus type 16 DNA sequences in archival cervical tissues by the polymerase chain reaction.
Arch Pathol Lab Med 116: 632-4 (1992)

- Hording U; Teglbjaerg CS; Visfeldt J; Bock JE
Human papillomavirus types 16 and 18 in adenocarcinoma of the uterine cervix.
Gynecol Oncol 46: 313-6 (1992)
- Howard E; Marcus D; O'Brien J; Albert D; Bernards R
Five DNA tumor viruses undetectable in human retinoblastomas.
Invest Ophthalmol Vis Sci 33: 1564-7 (1992)
- Hsu EM; McNicol PJ
Characterization of HPV-16 E6/E7 transcription in CaSki cells by quantitative PCR.
Mol Cell Probes 6: 459-66 (1992)
- Hubbert NL; Sedman SA; Schiller JT
Human papillomavirus type 16 E6 increases the degradation rate of p53 in human keratinocytes.
J Virol 66: 6237-41 (1992)
- Hummel M; Hudson JB; Laimins LA
Differentiation-induced and constitutive transcription of human papillomavirus type 31b in cell lines containing viral episomes.
J Virol 66: 6070-80 (1992)
- Hwang JY; Lin BY; Tang FM; Yu WC
Tamoxifen stimulates human papillomavirus type 16 gene expression and cell proliferation in a cervical cancer cell line.
Cancer Res 52: 6848-52 (1992)
- Icenogle JP; Laga M; Miller D; Manoka AT; Tucker RA; Reeves WC
Genotypes and sequence variants of human papillomavirus DNAs from human immunodeficiency virus type 1-infected women with cervical intraepithelial neoplasia.
J Infect Dis 166: 1210-6 (1992)
- Iftner T; Oft M; Bohm S; Wilczynski SP; Pfister H
Transcription of the E6 and E7 genes of human papillomavirus type 6 in anogenital condylomata is restricted to undifferentiated cell layers of the epithelium.
J Virol 66: 4639-46 (1992)
- Inoue H; Kondoh G; Sudiro TM; Hakura A
Stability of p53 protein in rat cells transformed by various viral transforming genes.
Virology 187: 343-7 (1992)
- Ip EW; Collins RJ; Srivastava G; Cheung AN; Wong LC
Human papillomavirus and its prognostic significance in invasive carcinoma of the cervix in young patients.
Int J Gynecol Pathol 11: 258-65 (1992)
- Ishii K; Inoue H; Hakura A; Araki W; Masuda T
Immortalization of fetal mouse brain glial cells by human papillomavirus type 16 E7 genes.
Cell Struct Funct 17: 197-202 (1992)
- Ishiji T; Lacey MJ; Parkkinen S; Anderson RD; Haugen TH; Cripe TP; Xiao JH; Davidson I; Chambon P; Turek LP
Transcriptional enhancer factor (TEF)-1 and its cell-specific co-activator activate human papillomavirus-16 E6 and E7 oncogene transcription in keratinocytes and cervical carcinoma cells.
EMBO J 11: 2271-81 (1992)
- Iwasaka T; Hayashi Y; Yokoyama M; Hara K; Matsuo N; Sugimori H
'Hit and run' oncogenesis by human papillomavirus type 18 DNA.
Acta Obstet Gynecol Scand 71: 219-23 (1992)
- Iwasaka T; Yokoyama M; Oh-uchida M; Matsuo N; Hara K; Fukuyama K; Hachisuga T; Fukuda K; Sugimori H
Detection of human papillomavirus genome and analysis of expression of c-myc and Ha-ras oncogenes in invasive cervical carcinomas.
Gynecol Oncol 46: 298-303 (1992)
- Jareborg N; Alderborn A; Burnett S
Identification and genetic definition of a bovine papillomavirus type 1 E7 protein and absence of a low-copy-number phenotype exhibited by E5, E6, or E7 viral mutants.
J Virol 66: 4957-65 (1992)
- Jewers RJ; Hildebrandt P; Ludlow JW; Kell B; McCance DJ
Regions of human papillomavirus type 16 E7 oncoprotein required for immortalization of human keratinocytes.
J Virol 66: 1329-35 (1992)
- Jones RE; Heimbrook DC; Huber HE; Wegrzyn RJ; Rotberg NS; Stauffer KJ; Lumma PK; Garsky VM; Oliff A
Specific N-methylations of HPV-16 E7 peptides alter binding to the retinoblastoma suppressor protein.
J Biol Chem 267: 908-12 (1992)
- Kahn T; Friesl H; Copeland NG; Gilbert DJ; Jenkins NA; Gissmann L; Kramer J; zur Hausen H
Molecular cloning, analysis, and chromosomal localization of a mouse genomic sequence related to the human papillomavirus type 18 E5 region.
Mol Carcinog 6: 88-99 (1992)

References (1992)

- Kanda T; Onda T; Zanma S; Yasugi T; Furuno A; Watanabe S; Kawana T; Sugase M; Ueda K; Sonoda T; et al
Independent association of antibodies against human papillomavirus type 16 E1/E4 and E7 proteins with cervical cancer.
Virology 190: 724-32 (1992)
- Kashima HK; Kessiss T; Hruban RH; Wu TC; Zinreich SJ; Shah KV
Human papillomavirus in sinonasal papillomas and squamous cell carcinoma.
Laryngoscope 102: 973-6 (1992)
- Kashiwabara K; Nakajima T
Detection of human papillomavirus DNA in invasive cervical cancers by the polymerase chain reaction and its clinical significance.
Acta Pathol Jpn 42: 876-83 (1992)
- Kato H; Nishida J; Honda T; Miyamoto S; Fujinaga K; Wake N
Chromosome alterations contribute to neoplastic progression of transformed rat embryonal fibroblasts.
Cancer Genet Cytogenet 58: 39-47 (1992)
- Kaur P; Carter WG
Integrin expression and differentiation in transformed human epidermal cells is regulated by fibroblasts.
J Cell Sci 103 (Pt 3): 755-63 (1992)
- Keating MR
Antiviral agents.
Mayo Clin Proc 67: 160-78 (1992)
- Kellokoski JK; Syrjanen SM; Chang F; Yliskoski M; Syrjanen KJ
Southern blot hybridization and PCR in detection of oral human papillomavirus (HPV) infections in women with genital HPV infections.
J Oral Pathol Med 21: 459-64 (1992)
- Khan IM; Malcolm AD; Kamalati T; Buluwela L
Protein-DNA interactions and a putative papillomavirus binding factor.
Biochem Soc Trans 20: 212S (1992)
- Kimbauer R; Booy F; Cheng N; Lowy DR; Schiller JT
Papillomavirus L1 major capsid protein self-assembles into virus-like particles that are highly immunogenic.
Proc Natl Acad Sci U S A 89: 12180-4 (1992)
- Konno R; Sato S; Yajima A
Detection of human papillomavirus DNA in normal epithelium and in squamous metaplasia of the uterine cervix by the polymerase chain reaction.
Tohoku J Exp Med 168: 491-7 (1992)
- Kowalski LA; Tsang SS; Davison AJ
Vanadate enhances transformation of bovine papillomavirus DNA-transfected C3H/10T1/2 cells.
Cancer Lett 64: 83-90 (1992)
- Kulke R; Horwitz BH; Zibello T; DiMaio D
The central hydrophobic domain of the bovine papillomavirus E5 transforming protein can be functionally replaced by many hydrophobic amino acid sequences containing a glutamine.
J Virol 66: 505-11 (1992)
- Lai CH; Hsueh S; Lin CY; Huang MY; You GB; Chang HC; Pao CC
Human papillomavirus in benign and malignant ovarian and endometrial tissues.
Int J Gynecol Pathol 11: 210-5 (1992)
- Lazo PA; Gallego MI; Ballester S; Feduchi E
Genetic alterations by human papillomaviruses in oncogenesis.
FEBS Lett 300: 109-13 (1992)
- Lechner MS; Mack DH; Finicle AB; Crook T; Vousden KH; Laimins LA
Human papillomavirus E6 proteins bind p53 in vivo and abrogate p53-mediated repression of transcription.
EMBO J 11: 3045-52 (1992)
- Leechanachai P; Banks L; Moreau F; Matlashewski G
The E5 gene from human papillomavirus type 16 is an oncogene which enhances growth factor-mediated signal transduction to the nucleus.
Oncogene 7: 19-25 (1992)
- Lehtinen M; Leminen A; Kuoppala T; Tiikkainen M; Lehtinen T; Lehtovirta P; Punnonen R; Vesterinen E; Paavonen J
Pre- and posttreatment serum antibody responses to HPV 16 E2 and HSV 2 ICP8 proteins in women with cervical carcinoma.
J Med Virol 37: 180-6 (1992)
- Lehtinen M; Leminen A; Paavonen J; Lehtovirta P; Hyoty H; Vesterinen E; Dillner J
Predominance of serum antibodies to synthetic peptide stemming from HPV 18 open reading frame E2 in cervical adenocarcinoma.
J Clin Pathol 45: 494-7 (1992)
- Lehtinen M; Stellato G; Hyoty H; Nieminen P; Vesterinen E; Paavonen J
A T-helper cell epitope overlaps a major B-cell epitope in human papillomavirus type 18 E2 protein.
APMIS 100: 1022-6 (1992)

- Li SL; Kim MS; Cherrick HM; Doniger J; Park NH
Sequential combined tumorigenic effect of HPV-16 and chemical carcinogens.
Carcinogenesis 13: 1981-7 (1992)
- Li SL; Kim MS; Cherrick HM; Park NH
Low p53 level in immortal, non-tumorigenic oral keratinocytes harboring HPV-16 DNA.
Eur J Cancer B Oral Oncol 28B: 129-34 (1992)
- Lin YL; Borenstein LA; Selvakumar R; Ahmed R; Wettstein FO
Effective vaccination against papilloma development by immunization with L1 or L2 structural protein of cottontail rabbit papillomavirus.
Virology 187: 612-9 (1992)
- Liu W
[Detection of human papillomavirus (HPV) DNA in carcinoma of uterine cervix and genital tract diseases in Jiangsu province, China]
Chung Hua Chung Liu Tsa Chih 13: 406-8 (1992)
- Lorincz A
Detection of human papillomavirus DNA without amplification: prospects for clinical utility.
IARC Sci Publ 96: 135-45 (1992)
- Lungu O; Sun XW; Felix J; Richart RM; Silverstein S; Wright TC Jr
Relationship of human papillomavirus type to grade of cervical intraepithelial neoplasia.
JAMA 267: 2493-6 (1992)
- Madej M; Conway MJ; Morgan AJ; Sweet J; Wallace L; Qualtiere LF; Arrand JR; Mackett M
Purification and characterization of Epstein-Barr virus gp340/220 produced by a bovine papillomavirus virus expression vector system.
Vaccine 10: 777-82 (1992)
- Malejczyk J; Malejczyk M; Kock A; Urbanski A; Majewski S; Hunzelmann N; Jablonska S; Orth G; Luger TA
Autocrine growth limitation of human papillomavirus type 16-harboring keratinocytes by constitutively released tumor necrosis factor- α .
J Immunol 149: 2702-8 (1992)
- Manoharan M; Johnson LK; McGee DP; Guinasso CJ; Ramasamy K; Springer RH; Bennett CF; Ecker DJ; Vickers T; Cowsert L; et al
Chemical modifications to improve uptake and bioavailability of antisense oligonucleotides.
Ann N Y Acad Sci 660: 306-9 (1992)
- Marich JE; Pontsler AV; Rice SM; McGraw KA; Dubensky TW
The phylogenetic relationship and complete nucleotide sequence of human papillomavirus type 35.
Virology 186: 770-6 (1992)
- Masih AS; Stoler MH; Farrow GM; Wooldridge TN; Johansson SL
Penile verrucous carcinoma: a clinicopathologic, human papillomavirus typing and flow cytometric analysis.
Mod Pathol 5: 48-55 (1992)
- Masuda A; Kizaka-Kondoh S; Miwatani H; Terada Y; Nojima H; Okayama H
Signal transduction cascade shared by epidermal growth factor and platelet-derived growth factor is a major pathway for oncogenic transformation in NRK cells.
New Biol 4: 489-503 (1992)
- Matsukura T; Iwasaki T; Kawashima M
Molecular cloning of a novel human papillomavirus (type 60) from a plantar cyst with characteristic pathological changes.
Virology 190: 561-4 (1992)
- McBride AA; Klausner RD; Howley PM
Conserved cysteine residue in the DNA-binding domain of the bovine papillomavirus type 1 E2 protein confers redox regulation of the DNA-binding activity in vitro.
Proc Natl Acad Sci U S A 89: 7531-5 (1992)
- McGlennen RC; Ghai J; Ostrow RS; LaBresh K; Schneider JF; Faras AJ
Cellular transformation by a unique isolate of human papillomavirus type 11.
Cancer Res 52: 5872-8 (1992)
- McNicol P; Guijon F; Brunham R; Gray M; Paraskevas M
Laboratory diagnosis of latent human papillomavirus infection.
Diagn Microbiol Infect Dis 15: 679-83 (1992)
- Merrick DT; Blanton RA; Gown AM; McDougall JK
Altered expression of proliferation and differentiation markers in human papillomavirus 16 and 18 immortalized epithelial cells grown in organotypic culture.
Am J Pathol 140: 167-77 (1992)
- Meyers C; Harry J; Lin YL; Wettstein FO
Identification of three transforming proteins encoded by cottontail rabbit papillomavirus.
J Virol 66: 1655-64 (1992)

References (1992)

- Mietz JA; Unger T; Huibregtse JM; Howley PM
The transcriptional transactivation function of wild-type p53 is inhibited by SV40 large T-antigen and by HPV-16 E6 oncoprotein.
EMBO J 11: 5013-20 (1992)
- Mincheva A; Karag'ozov I
[Interferon, interferon inducers, human papillomavirus infection]
Akush Ginekol (Sofia) 31: 28-30 (1992)
- Mitrani-Rosenbaum S; Tsvieli R
Differential cooperation of a carcinogen with human papillomavirus type 6 and 16 DNAs in in vitro oncogenic transformation.
Intervirology 33: 76-85 (1992)
- Moreau F; Matlashewski G
Molecular analysis of different allelic variants of wild-type human p53.
Biochem Cell Biol 70: 1014-9 (1992)
- Morgan DM; Pecoraro G; Rosenberg I; Defendi V
Transformation by human papillomavirus type 16 (HPV16) DNA but not HPV6b DNA is enhanced by addition of the human cytomegalovirus enhancer.
Virology 189: 687-94 (1992)
- Morre DJ; Morre DM; Paulik M; Batova A; Broome AM; Pirisi L; Creek KE
Retinoic acid and calcitriol inhibition of growth and NADH oxidase of normal and immortalized human keratinocytes.
Biochim Biophys Acta 1134: 217-22 (1992)
- Morrison EA; Goldberg GL; Kadish AS; Burk RD
Polymerase chain reaction detection of human papillomavirus: quantitation may improve clinical utility.
J Clin Microbiol 30: 2539-43 (1992)
- Muller M; Viscidi RP; Sun Y; Guerrero E; Hill PM; Shah F; Bosch FX; Munoz N; Gissmann L; Shah KV
Antibodies to HPV-16 E6 and E7 proteins as markers for HPV-16-associated invasive cervical cancer.
Virology 187: 508-14 (1992)
- Mumm SR; Hippenmeyer PJ; Grandgenett DP
Characterization of a stable eukaryotic cell line expressing the Rous sarcoma virus integrase.
Virology 189: 500-10 (1992)
- Mungal S; Steinberg BM; Taichman LB
Replication of plasmid-derived human papillomavirus type 11 DNA in cultured keratinocytes.
J Virol 66: 3220-4 (1992)
- Munger K; Scheffner M; Huibregtse JM; Howley PM
Interactions of HPV E6 and E7 oncoproteins with tumour suppressor gene products.
Cancer Surv 12: 197-217 (1992)
- Nair BS; Pillai R
Oncogenesis of squamous carcinoma of the uterine cervix.
Int J Gynecol Pathol 11: 47-57 (1992)
- Nakazawa A; Inoue M; Saito J; Sasagawa T; Ueda G; Tanizawa O
Detection of human papillomavirus types 16 and 18 in the exfoliated cervical cells using the polymerase chain reaction.
Int J Gynaecol Obstet 37: 13-8 (1992)
- Novelli G; Gennarelli M; De Santis L; Angeloni P; Dallapiccola B
Inosine-containing primers in human papillomavirus detection by polymerase chain reaction.
Biomed Pharmacother 46: 167-9 (1992)
- O'Banion MK; Jacobson ER; Sundberg JP
Molecular cloning and partial characterization of a parrot papillomavirus.
Intervirology 33: 91-6 (1992)
- O'Banion MK; Levenson RM; Brinckmann UG; Young DA
Glucocorticoid modulation of transformed cell proliferation is oncogene specific and correlates with effects on c-myc levels.
Mol Endocrinol 6: 1371-80 (1992)
- Ogura H; Kawakami T; Fujiwara T; Sakai A; Saito R; Watanabe S; Masuda Y; Yabe Y
Detection of human papillomavirus type 6f genome in nasal papillomatosis.
Acta Otolaryngol (Stockh) 112: 115-9 (1992)
- Ostrow RS; Forslund KM; McGlennen RC; Shaw DP; Schlievert PM; Ussery MA; Huggins JW; Faras AJ
Ribavirin mitigates wart growth in rabbits at early stages of infection with cottontail rabbit papillomavirus.
Antiviral Res 17: 99-113 (1992)
- Pagano M; Durst M; Joswig S; Draetta G; Jansen-Durr P
Binding of the human E2F transcription factor to the retinoblastoma protein but not to cyclin A is abolished in HPV-16-immortalized cells.
Oncogene 7: 1681-6 (1992)
- Park NH; Li SL; Xie JF; Cherrick HM
In vitro and animal studies of the role of viruses in oral carcinogenesis.
Eur J Cancer B Oral Oncol 28B: 145-52 (1992)

- Pater A; Belaguli NS; Gardner HA; Mithal A; Pater MM
Glucocorticoid-dependent transformation by human papillomavirus type 16 E7 coding and 3' noncoding sequences.
Virology 188: 369-72 (1992)
- Pater MM; Nakshatri H; Kisaka C; Pater A
The first 124 nucleotides of the E7 coding sequences of HPV16 can render the HPV11 genome transformation competent.
Virology 186: 348-51 (1992)
- Patrick DR; Zhang K; Defeo-Jones D; Vuocolo GR; Maigetter RZ; Sardana MK; Oliff A; Heimbrook DC
Characterization of functional HPV-16 E7 protein produced in *Escherichia coli*.
J Biol Chem 267: 6910-5 (1992)
- Paz-y-Mino C; Ocampo L; Narvaez R; Narvaez L
Chromosome fragility in lymphocytes of women with cervical uterine lesions produced by human papillomavirus.
Cancer Genet Cytogenet 59: 173-6 (1992)
- Pelisson I; Soler C; Chignol MC; Euvrard S; Chardonnet Y
[Human papillomaviruses and cellular oncogenes (c-myc, c-Ha-ras) in cutaneous and mucosal lesions in transplantation recipients]
Bull Cancer (Paris) 79: 471-82 (1992)
- Pelisson I; Soler C; Pechoux C; Chignol MC; Viac J; Euvrard S; Chardonnet Y
c-myc and c-Ha-ras cellular oncogenes and human papillomaviruses in benign and malignant cutaneous lesions.
J Dermatol Sci 3: 56-67 (1992)
- Penneys NS; Friend A; Zhu WY; Leonardi C
Swollen keratinocytes: a histologic marker of unusual human papillomavirus-type infection and immunosuppression.
J Cutan Pathol 19: 217-20 (1992)
- Pennie WD; Campo MS
Synergism between bovine papillomavirus type 4 and the flavonoid quercetin in cell transformation in vitro.
Virology 190: 861-5 (1992)
- Perez-Reyes N; Halbert CL; Smith PP; Benditt EP; McDougall JK
Immortalization of primary human smooth muscle cells.
Proc Natl Acad Sci U S A 89: 1224-8 (1992)
- Petti L; DiMaio D
Stable association between the bovine papillomavirus E5 transforming protein and activated platelet-derived growth factor receptor in transformed mouse cells.
Proc Natl Acad Sci U S A 89: 6736-40 (1992)
- Phelps WC; Munger K; Yee CL; Barnes JA; Howley PM
Structure-function analysis of the human papillomavirus type 16 E7 oncoprotein.
J Virol 66: 2418-27 (1992)
- Philipp W; Honore N; Sapp M; Cole ST; Streeck RE
Human papillomavirus type 42: new sequences, conserved genome organization.
Virology 186: 331-4 (1992)
- Pim D; Collins M; Banks L
Human papillomavirus type 16 E5 gene stimulates the transforming activity of the epidermal growth factor receptor.
Oncogene 7: 27-32 (1992)
- Pirisi L; Batova A; Jenkins GR; Hodam JR; Creek KE
Increased sensitivity of human keratinocytes immortalized by human papillomavirus type 16 DNA to growth control by retinoids.
Cancer Res 52: 187-93 (1992)
- Pozzi G; Contorni M; Oggioni MR; Manganeli R; Tommasino M; Cavalieri F; Fischetti VA
Delivery and expression of a heterologous antigen on the surface of streptococci.
Infect Immun 60: 1902-7 (1992)
- Prakash SS; Grossman SR; Pepinsky RB; Laimins LA; Androphy EJ
Amino acids necessary for DNA contact and dimerization imply novel motifs in the papillomavirus E2 transactivator.
Genes Dev 6: 105-16 (1992)
- Ravnan JB; Gilbert DM; Ten Hagen KG; Cohen SN
Random-choice replication of extrachromosomal bovine papillomavirus (BPV) molecules in heterogeneous, clonally derived BPV-infected cell lines.
J Virol 66: 6946-52 (1992)
- Reiss M; Brash DE; Munoz-Antonia T; Simon JA; Ziegler A; Vellucci VF; Zhou ZL
Status of the p53 tumor suppressor gene in human squamous carcinoma cell lines.
Oncol Res 4: 349-57 (1992)
- Remm M; Brain R; Jenkins JR
The E2 binding sites determine the efficiency of replication for the origin of human papillomavirus type 18.
Nucleic Acids Res 20: 6015-21 (1992)

References (1992)

- Riou G; Le MG; Favre M; Jeannel D; Bourhis J; Orth G
Human papillomavirus-negative status and c-myc gene overexpression: independent prognostic indicators of distant metastasis for early-stage invasive cervical cancers.
J Natl Cancer Inst 84: 1525-6 (1992)
- Rogel-Gaillard C; Breitburd F; Orth G
Human papillomavirus type 1 E4 proteins differing by their N-terminal ends have distinct cellular localizations when transiently expressed in vitro.
J Virol 66: 816-23 (1992)
- Romanczuk H; Howley PM
Disruption of either the E1 or the E2 regulatory gene of human papillomavirus type 16 increases viral immortalization capacity.
Proc Natl Acad Sci U S A 89: 3159-63 (1992)
- Rose BR; Thompson CH; Chantrill LA; Tattersall MH; Cossart YE
Prevalence and distribution of human papillomavirus type-16 DNA in pelvic lymph nodes of patients with cervical cancer and in women with no history of cervical abnormality.
Int J Cancer 52: 225-8 (1992)
- Roth EJ; Kurz B; Liang L; Hansen CL; Dameron CT; Winge DR; Smotkin D
Metal thiolate coordination in the E7 proteins of human papilloma virus 16 and cottontail rabbit papilloma virus as expressed in *Escherichia coli*.
J Biol Chem 267: 16390-5 (1992)
- Rubben A; Beaudenon S; Favre M; Schmitz W; Spelten B; Grussendorf-Conen EI
Rearrangements of the upstream regulatory region of human papillomavirus type 6 can be found in both Buschke-Lowenstein tumours and in condylomata acuminata.
J Gen Virol 73 (Pt 12): 3147-53 (1992)
- Sang BC; Barbosa MS
Increased E6/E7 transcription in HPV 18-immortalized human keratinocytes results from inactivation of E2 and additional cellular events.
Virology 189: 448-55 (1992)
- Sang BC; Barbosa MS
Single amino acid substitutions in "low-risk" human papillomavirus (HPV) type 6 E7 protein enhance features characteristic of the "high-risk" HPV E7 oncoproteins.
Proc Natl Acad Sci U S A 89: 8063-7 (1992)
- Sarkar FH; Miles BJ; Plieth DH; Crissman JD
Detection of human papillomavirus in squamous neoplasm of the penis.
J Urol 147: 389-92 (1992)
- Sarkar FH; Visscher DW; Kintanar EB; Zarbo RJ; Crissman JD
Sinonasal Schneiderian papillomas: human papillomavirus typing by polymerase chain reaction.
Mod Pathol 5: 329-32 (1992)
- Sasagawa T; Inoue M; Inoue H; Yutsudo M; Tanizawa O; Hakura A
Induction of uterine cervical neoplasias in mice by human papillomavirus type 16 E6/E7 genes.
Cancer Res 52: 4420-6 (1992)
- Sasagawa T; Inoue M; Tanizawa O; Yutsudo M; Hakura A
Identification of antibodies against human papillomavirus type 16 E6 and E7 proteins in sera of patients with cervical neoplasias.
Jpn J Cancer Res 83: 705-13 (1992)
- Sastre-Garau X; Couturier J; Orth G
[Chromosome sites of papillomavirus genome integration in genital tumors]
Ann Pathol 12: 55-8 (1992)
- Scheffner M; Munger K; Huibregtse JM; Howley PM
Targeted degradation of the retinoblastoma protein by human papillomavirus E7-E6 fusion proteins.
EMBO J 11: 2425-31 (1992)
- Scheffner M; Takahashi T; Huibregtse JM; Minna JD; Howley PM
Interaction of the human papillomavirus type 16 E6 oncoprotein with wild-type and mutant human p53 proteins.
J Virol 66: 5100-5 (1992)
- Scinicariello F; Sato T; Lee CS; Hsu HC; Chan TS; Tyring SK
Detection of human papillomavirus in primary hepatocellular carcinoma.
Anticancer Res 12: 763-6 (1992)
- Sedman SA; Hubbert NL; Vass WC; Lowy DR; Schiller JT
Mutant p53 can substitute for human papillomavirus type 16 E6 in immortalization of human keratinocytes but does not have E6-associated trans-activation or transforming activity.
J Virol 66: 4201-8 (1992)
- Selvey LA; Dunn LA; Murray B; Tindle RW; Frazer IH
An ELISA capture assay for the E7 transforming proteins of HPV16 and HPV18.
J Virol Methods 37: 119-27 (1992)

- Sherman L; Alloul N
Human papillomavirus type 16 expresses a variety of alternatively spliced mRNAs putatively encoding the E2 protein.
Virology 191: 953-9 (1992)
- Sherman L; Alloul N; Golan I; Durst M; Baram A
Expression and splicing patterns of human papillomavirus type-16 mRNAs in pre-cancerous lesions and carcinomas of the cervix, in human keratinocytes immortalized by HPV 16, and in cell lines established from cervical cancers.
Int J Cancer 50: 356-64 (1992)
- Shibata D; Hawes D; Li ZH; Hernandez AM; Spruck CH; Nichols PW
Specific genetic analysis of microscopic tissue after selective ultraviolet radiation fractionation and the polymerase chain reaction.
Am J Pathol 141: 539-43 (1992)
- Shillitoe EJ; Steele C
Inhibition of the transformed phenotype of carcinoma cells that contain human papillomavirus.
Ann N Y Acad Sci 660: 286-7 (1992)
- Shimada M; Fujinaga K
[Detection of human papillomavirus DNA in cervical carcinoma by the polymerase chain reaction]
Nippon Rinsho 50 Suppl: 193-8 (1992)
- Shindoh M; Sawada Y; Kohgo T; Amemiya A; Fujinaga K
Detection of human papillomavirus DNA sequences in tongue squamous-cell carcinoma utilizing the polymerase chain reaction method.
Int J Cancer 50: 167-71 (1992)
- Shroyer KR; Kim JG; Manos MM; Greer CE; Pearlman NW; Franklin WA
Papillomavirus found in anorectal squamous carcinoma, not in colon adenocarcinoma.
Arch Surg 127: 741-4 (1992)
- Shu LL; Feng HM; Jin Q; Hou YD
Molecular cloning of a new variant of human papillomavirus type 6 isolated from a Chinese woman and expression of its L1 gene in *E. coli*-molecular cloning & expression of HPV6 gene.
Sci China [B] 35: 585-91 (1992)
- Sircar S; Horvath J; Roberge D; Diouri M; Weber JM
Adenovirus transformation revertant resistant to retransformation by E1 but not by SV40-T and HPV16-E7 oncogenes.
Virology 191: 187-92 (1992)
- Smith PP; Friedman CL; Bryant EM; McDougall JK
Viral integration and fragile sites in human papillomavirus-immortalized human keratinocyte cell lines.
Genes Chromosom Cancer 5: 150-7 (1992)
- Smits HL; Tieben LM; Tjong-A-Hung SP; Jebbink MF; Minnaar RP; Jansen CL; ter Schegget J
Detection and typing of human papillomaviruses present in fixed and stained archival cervical smears by a consensus polymerase chain reaction and direct sequence analysis allow the identification of a broad spectrum of human papillomavirus types.
J Gen Virol 73 (Pt 12): 3263-8 (1992)
- Smits PH; de Ronde A; Smits HL; Minnaar RP; van der Noordaa J; ter Schegget J
Modulation of the human papillomavirus type 16 induced transformation and transcription by deletion of loci on the short arm of human chromosome 11 can be mimicked by SV40 small t.
Virology 190: 40-4 (1992)
- Snijders PJ; Cromme FV; van den Brule AJ; Schrijnemakers HF; Snow GB; Meijer CJ; Walboomers JM
Prevalence and expression of human papillomavirus in tonsillar carcinomas, indicating a possible viral etiology.
Int J Cancer 51: 845-50 (1992)
- Snijders PJ; Meijer CJ; van den Brule AJ; Schrijnemakers HF; Snow GB; Walboomers JM
Human papillomavirus (HPV) type 16 and 33 E6/E7 region transcripts in tonsillar carcinomas can originate from integrated and episomal HPV DNA.
J Gen Virol 73 (Pt 8): 2059-66 (1992)
- Snijders PJ; van den Brule AJ; Schrijnemakers HF; Raaphorst PM; Meijer CJ; Walboomers JM
Human papillomavirus type 33 in a tonsillar carcinoma generates its putative E7 mRNA via two E6* transcript species which are terminated at different early region poly(A) sites.
J Virol 66: 3172-8 (1992)
- Soler C; Chardonnet Y; Allibert P; Euvrard S; Mandrand B; Thivolet J
Detection of multiple types of human papillomavirus in a giant condyloma from a grafted patient. Analysis by immunohistochemistry, in situ hybridisation, Southern blot and polymerase chain reaction.
Virus Res 23: 193-208 (1992)

References (1992)

- Spitkovsky DD; Royer-Pokora B; Delius H; Kisseljov F; Jenkins NA; Gilbert DJ; Copeland NG; Royer HD
Tissue restricted expression and chromosomal localization of the YB-1 gene encoding a 42 kD nuclear CCAAT binding protein.
Nucleic Acids Res 20: 797-803 (1992)
- Stacey SN; Bartholomew JS; Ghosh A; Stern PL; Mackett M; Arrand JR
Expression of human papillomavirus type 16 E6 protein by recombinant baculovirus and use for detection of anti-E6 antibodies in human sera.
J Gen Virol 73 (Pt 9): 2337-45 (1992)
- Steele C; Sacks PG; Adler-Storthz K; Shillitoe EJ
Effect on cancer cells of plasmids that express antisense RNA of human papillomavirus type 18.
Cancer Res 52: 4706-11 (1992)
- Steger G; Pfister H
In vitro expressed HPV 8 E6 protein does not bind p53.
Arch Virol 125: 355-60 (1992)
- Steuer B; Stoppler H; Breuer B; Alonso A
Transcription of BPV-1 genes in transfected F9 cells.
Oncogene 7: 1875-8 (1992)
- Stirdivant SM; Ahern JD; Oliff A; Heimbrook DC
Retinoblastoma protein binding properties are dependent on 4 cysteine residues in the protein binding pocket.
J Biol Chem 267: 14846-51 (1992)
- Stirdivant SM; Huber HE; Patrick DR; Defeo-Jones D; McAvoy EM; Garsky VM; Oliff A; Heimbrook DC
Human papillomavirus type 16 E7 protein inhibits DNA binding by the retinoblastoma gene product.
Mol Cell Biol 12: 1905-14 (1992)
- Stoler MH; Rhodes CR; Whitbeck A; Wolinsky SM; Chow LT; Broker TR
Human papillomavirus type 16 and 18 gene expression in cervical neoplasias.
Hum Pathol 23: 117-28 (1992)
- Storey A; Greenfield I; Banks L; Pim D; Crook T; Crawford L; Stanley M
Lack of immortalizing activity of a human papillomavirus type 16 variant DNA with a mutation in the E2 gene isolated from normal human cervical keratinocytes.
Oncogene 7: 459-65 (1992)
- Stubenrauch F; Malejczyk J; Fuchs PG; Pfister H
Late promoter of human papillomavirus type 8 and its regulation.
J Virol 66: 3485-93 (1992)
- Sugase M
[Uterine cervical carcinoma and human papillomaviruses]
Hum Cell 5: 143-9 (1992)
- Sundberg JP; Kenty GA; Beamer WG; Adkison DL
Forestomach papillomas in flaky skin and steel-Dickie mutant mice.
J Vet Diagn Invest 4: 312-7 (1992)
- Syrjanen S; Andersson B; Juntunen L; Syrjanen K
Polymerase chain reaction for producing biotinylated human papillomavirus DNA probes for in situ hybridization.
Sex Transm Dis 19: 140-5 (1992)
- Takahara Y; Hamada K; Housman DE
A new retrovirus packaging cell for gene transfer constructed from amplified long terminal repeat-free chimeric proviral genes.
J Virol 66: 3725-32 (1992)
- Takami Y; Sasagawa T; Sudiro TM; Yutsudo M; Hakura A
Determination of the functional difference between human papillomavirus type 6 and 16 E7 proteins by their 30 N-terminal amino acid residues.
Virology 186: 489-95 (1992)
- Tan SH; Gloss B; Bernard HU
During negative regulation of the human papillomavirus-16 E6 promoter, the viral E2 protein can displace Sp1 from a proximal promoter element.
Nucleic Acids Res 20: 251-6 (1992)
- Tanimoto H
Occurrence and expression of human papillomavirus type 16 genes in uterine cervical carcinomas.
Hiroshima J Med Sci 41: 71-7 (1992)
- Tay SK; Chow VT; Tham KM; Lim-Tan SK; Bernard HU
Patterns and implications of subclinical vulval human papillomavirus infection: the impact of PCR analysis.
Asia Oceania J Obstet Gynaecol 18: 65-71 (1992)
- Thierry F; Spyrou G; Yaniv M; Howley P
Two AP1 sites binding JunB are essential for human papillomavirus type 18 transcription in keratinocytes.
J Virol 66: 3740-8 (1992)
- Tinsley JM; Fisher C; Searle PF
Abnormalities of epidermal differentiation associated with expression of the human papillomavirus type 1 early region in transgenic mice.
J Gen Virol 73 (Pt 5): 1251-60 (1992)

References (1992)

- Toh Y; Kuwano H; Tanaka S; Baba K; Matsuda H; Sugimachi K; Mori R
Detection of human papillomavirus DNA in esophageal carcinoma in Japan by polymerase chain reaction.
Cancer 70: 2234-8 (1992)
- Tommasino M; Contorni M; Cavalieri F
HPV16 E7 phosphorylation in fission yeast: characterization and biological effects.
Gene 111: 93-8 (1992)
- Tornesello ML; Buonaguro FM; Beth-Giraldo E; Kyalwazi SK; Giraldo G
Human papillomavirus (HPV) DNA in penile carcinomas and in two cell lines from high-incidence areas for genital cancers in Africa.
Int J Cancer 51: 587-92 (1992)
- Tosi P; Cintorino M; Santopietro R; Lio R; Barbini P; Ji H; Chang F; Kataja V; Syrjanen S; Syrjanen K
Prognostic factors in invasive cervical carcinomas associated with human papillomavirus (HPV). Quantitative data and cytokeratin expression.
Pathol Res Pract 188: 866-73 (1992)
- Tseng CJ; Lin CY; Wang RL; Chen LJ; Chang YL; Hsieh TT; Pao CC
Possible transplacental transmission of human papillomaviruses.
Am J Obstet Gynecol 166: 35-40 (1992)
- Tsuda H; Hirohashi S
Frequent occurrence of p53 gene mutations in uterine cancers at advanced clinical stage and with aggressive histological phenotypes.
Jpn J Cancer Res 83: 1184-91 (1992)
- Tsutsumi K; Belaguli N; Qi S; Michalak TI; Gulliver WP; Pater A; Pater MM
Human papillomavirus 16 DNA immortalizes two types of normal human epithelial cells of the uterine cervix.
Am J Pathol 140: 255-61 (1992)
- Uemura N; Ozawa K; Tojo A; Takahashi K; Okano A; Karasuyama H; Tani K; Asano S
Acquisition of interleukin-3 independence in FDC-P2 cells after transfection with the activated c-H-ras gene using a bovine papillomavirus-based plasmid vector.
Blood 80: 3198-204 (1992)
- Van Ranst M; Fuse A; Fiten P; Beuken E; Pfister H; Burk RD; Opendakker G
Human papillomavirus type 13 and pygmy chimpanzee papillomavirus type 1: comparison of the genome organizations.
Virology 190: 587-96 (1992)
- Van Ranst M; Tachezy R; Pruss J; Burk RD
Primary structure of the E6 protein of *Micromys minutus* papillomavirus and *Mastomys natalensis* papillomavirus.
Nucleic Acids Res 20: 2889 (1992)
- Vande Pol SB; Howley PM
The bovine papillomavirus constitutive enhancer is essential for viral transformation, DNA replication, and the maintenance of latency.
J Virol 66: 2346-58 (1992)
- van den Brule AJ; Snijders PJ; Raaphorst PM; Schrijnemakers HF; Delius H; Gissmann L; Meijer CJ; Walboomers JM
General primer polymerase chain reaction in combination with sequence analysis for identification of potentially novel human papillomavirus genotypes in cervical lesions.
J Clin Microbiol 30: 1716-21 (1992)
- Villa LL; Vieira KB; Pei XF; Schlegel R
Differential effect of tumor necrosis factor on proliferation of primary human keratinocytes and cell lines containing human papillomavirus types 16 and 18.
Mol Carcinog 6: 5-9 (1992)
- von Knebel Doeberitz M
Papillomaviruses in human disease: Part II. Molecular biology and immunology of papillomavirus infections and carcinogenesis.
Eur J Med 1: 485-91 (1992)
- von Knebel Doeberitz M; Rittmuller C; zur Hausen H; Durst M
Inhibition of tumorigenicity of cervical cancer cells in nude mice by HPV E6-E7 anti-sense RNA [letter]
Int J Cancer 51: 831-4 (1992)
- Vormwald-Dogan V; Fischer B; Bludau H; Freese UK; Gissmann L; Glitz D; Schwartz E; Durst M
Sense and antisense transcripts of human papillomavirus type 16 in cervical cancers.
J Gen Virol 73 (Pt 7): 1833-8 (1992)
- Waldenstrom M; Schenstrom K; Sollerbrant K; Hansson L
Replication of bovine papillomavirus vectors in murine cells.
Gene 120: 175-81 (1992)

References (1992)

- Watanabe S; Sato H; Furuno A; Yoshiike K
Changing the spacing between metal-binding motifs decreases stability and transforming activity of the human papillomavirus type 18 E7 oncoprotein.
Virology 190: 872-5 (1992)
- Watanabe S; Sato H; Komiyama N; Kanda T; Yoshiike K
The E7 functions of human papillomaviruses in rat 3Y1 cells.
Virology 187: 107-14 (1992)
- Wiener JS; Effert PJ; Humphrey PA; Yu L; Liu ET; Walther PJ
Prevalence of human papillomavirus types 16 and 18 in squamous-cell carcinoma of the penis: a retrospective analysis of primary and metastatic lesions by differential polymerase chain reaction.
Int J Cancer 50: 694-701 (1992)
- Wiener JS; Liu ET; Walther PJ
Oncogenic human papillomavirus type 16 is associated with squamous cell cancer of the male urethra.
Cancer Res 52: 5018-23 (1992)
- Wikstrom A; Eklund C; Von Krogh G; Lidbrink P; Dillner J
Levels of immunoglobulin G antibodies against defined epitopes of the L1 and L2 capsid proteins of human papillomavirus type 6 are elevated in men with a history of condylomata acuminata.
J Clin Microbiol 30: 1795-800 (1992)
- Wilson VG; Ludes-Meyers J
Partite expression of the bovine papillomavirus E1 open reading frame in *Escherichia coli*.
Biochim Biophys Acta 1129: 215-8 (1992)
- Winokur PL; McBride AA
Separation of the transcriptional activation and replication functions of the bovine papillomavirus-1 E2 protein.
EMBO J 11: 4111-8 (1992)
- Woodworth CD; Cheng S; Simpson S; Hamacher L; Chow LT; Broker TR; DiPaolo JA
Recombinant retroviruses encoding human papillomavirus type 18 E6 and E7 genes stimulate proliferation and delay differentiation of human keratinocytes early after infection.
Oncogene 7: 619-26 (1992)
- Woodworth CD; Lichti U; Simpson S; Evans CH; DiPaolo JA
Leukoregulin and gamma-interferon inhibit human papillomavirus type 16 gene transcription in human papillomavirus-immortalized human cervical cells.
Cancer Res 52: 456-63 (1992)
- Wrede D; Luqmani YA; Coombes RC; Vousden KH
Absence of HPV 16 and 18 DNA in breast cancer.
Br J Cancer 65: 891-4 (1992)
- Yaegashi N; Jenison SA; Batra M; Galloway DA
Human antibodies recognize multiple distinct type-specific and cross-reactive regions of the minor capsid proteins of human papillomavirus types 6 and 11.
J Virol 66: 2008-19 (1992)
- Yokota H; Yoshikawa H; Mizuno M
[Amplification and typing of human papillomavirus (HPV) DNA from female genitalia by polymerase chain reaction using a consensus primer]
Nippon Rinsho 50 Suppl: 199-202 (1992)
- Zhou J; Sun XY; Davies H; Crawford L; Park D; Frazer IH
Definition of linear antigenic regions of the HPV16 L1 capsid protein using synthetic virion-like particles.
Virology 189: 592-9 (1992)
- Zhu WY; Leonardi C; Penneys NS
Detection of human papillomavirus DNA in seborrheic keratosis by polymerase chain reaction.
J Dermatol Sci 4: 166-71 (1992)

- Agrawal R; Pelkonen J; Mantyjarvi RA
Effect of gamma-interferon or cycloheximide treatment on viral and c-myc transcripts in bovine-papillomavirus-type-1-transformed primary mouse fibroblasts.
Intervirology 36: 44-52 (1993)
- Alejo M; Macedo I; Matias-Guiu X; Prat J
Adenocarcinoma in situ of the uterine cervix: clinicopathological study of nine cases with detection of human papillomavirus DNA by in situ hybridization and polymerase chain reaction.
Int J Gynecol Pathol 12: 219-23 (1993)
- Andrews EM; DiMaio D
Hierarchy of polyadenylation site usage by bovine papillomavirus in transformed mouse cells.
J Virol 67: 7705-10 (1993)
- Anwar K; Nakakuki K; Imai H; Naiki H; Inuzuka M
Over-expression of p53 protein in human laryngeal carcinoma.
Int J Cancer 53: 952-6 (1993)
- Anwar K; Nakakuki K; Naiki H; Inuzuka M
ras gene mutations and HPV infection are common in human laryngeal carcinoma.
Int J Cancer 53: 22-8 (1993)
- Apostolakos MJ; Schuermann WH; Frampton MW; Utell MJ; Willey JC
Measurement of gene expression by multiplex competitive polymerase chain reaction.
Anal Biochem 213: 277-84 (1993)
- Apt D; Chong T; Liu Y; Bernard HU
Nuclear factor I and epithelial cell-specific transcription of human papillomavirus type 16.
J Virol 67: 4455-63 (1993)
- Arany I; Rady P; Tying SK
Alterations in cytokine/antioncogene expression in skin lesions caused by "low-risk" types of human papillomaviruses.
Viral Immunol 6: 255-65 (1993)
- Arbeit JM; Munger K; Howley PM; Hanahan D
Neuroepithelial carcinomas in mice transgenic with human papillomavirus type 16 E6/E7 ORFs.
Am J Pathol 142: 1187-97 (1993)
- Arends MJ; Donaldson YK; Duvall E; Wyllie AH; Bird CC
Human papillomavirus type 18 associates with more advanced cervical neoplasia than human papillomavirus type 16.
Hum Pathol 24: 432-7 (1993)
- Armstrong DJ; Roman A
The anomalous electrophoretic behavior of the human papillomavirus type 16 E7 protein is due to the high content of acidic amino acid residues.
Biochem Biophys Res Commun 192: 1380-7 (1993)
- Arroyo M; Bagchi S; Raychaudhuri P
Association of the human papillomavirus type 16 E7 protein with the S-phase-specific E2F-cyclin A complex.
Mol Cell Biol 13: 6537-46 (1993)
- Band V; Dalal S; Delmolino L; Androphy EJ
Enhanced degradation of p53 protein in HPV-6 and BPV-1 E6-immortalized human mammary epithelial cells.
EMBO J 12: 1847-52 (1993)
- Barksdale SK; Baker CC
Differentiation-specific expression from the bovine papillomavirus type 1 P2443 and late promoters.
J Virol 67: 5605-16 (1993)
- Bavin PJ; Walker PG; Emery VC
Sequence microheterogeneity in the long control region of clinical isolates of human papillomavirus type 16.
J Med Virol 39: 267-72 (1993)
- Bohm S; Wilczynski SP; Pfister H; Iftner T
The predominant mRNA class in HPV16-infected genital neoplasias does not encode the E6 or the E7 protein.
Int J Cancer 55: 791-8 (1993)
- Brandsma JL; Xiao W
Infectious virus replication in papillomas induced by molecularly cloned cottontail rabbit papillomavirus DNA.
J Virol 67: 567-71 (1993)
- Braun L; Mikumo R; Mark HF; Lauchlan S
Analysis of the growth properties and physical state of the human papillomavirus type 16 genome in cell lines derived from primary cervical tumors.
Am J Pathol 143: 832-44 (1993)
- Bream GL; Ohmstede CA; Phelps WC
Characterization of human papillomavirus type 11 E1 and E2 proteins expressed in insect cells.
J Virol 67: 2655-63 (1993)
- Buchman GW; Schuster DM; Rashtchian A
Selective RNA amplification: a novel method using dUMP-containing primers and uracil DNA glycosylase.
PCR Methods Appl 3: 28-31 (1993)

References (1993)

- Burns JE; Baird MC; Clark LJ; Burns PA; Edington K; Chapman C; Mitchell R; Robertson G; Soutar D; Parkinson EK
Gene mutations and increased levels of p53 protein in human squamous cell carcinomas and their cell lines.
Br J Cancer 67: 1274-84 (1993)
- Butz K; Hoppe-Seyler F
Transcriptional control of human papillomavirus (HPV) oncogene expression: composition of the HPV type 18 upstream regulatory region.
J Virol 67: 6476-86 (1993)
- Carlotti F; Crawford L
Trans-activation of the adenovirus E2 promoter by human papillomavirus type 16 E7 is mediated by retinoblastoma-dependent and -independent pathways.
J Gen Virol 74 (Pt 11): 2479-86 (1993)
- Casazza S
[Expression of phosphoprotein P53 in pathology (editorial)]
Pathologica 85: 321-34 (1993)
- Cason J; Kambo PK; Jewers RJ; Chrystie IL; Best JM
Mapping of linear B cell epitopes on capsid proteins of bovine papillomavirus: identification of three external type-restricted epitopes.
J Gen Virol 74 (Pt 12): 2669-77 (1993)
- Chang F; Syrjanen S; Kurvinen K; Syrjanen K
The p53 tumor suppressor gene as a common cellular target in human carcinogenesis.
Am J Gastroenterol 88: 174-86 (1993)
- Chang F; Syrjanen S; Tervahauta A; Syrjanen K
Tumorigenesis associated with the p53 tumour suppressor gene.
Br J Cancer 68: 653-61 (1993)
- Chen BF
[Polymerase chain reaction in detection of human papillomavirus DNA in esophageal carcinoma]
Chung Hua I Hsueh Tsa Chih 73: 667-9, 701 (1993)
- Chen BF; Clejan S
Rapid preparation of tissue DNA from paraffin-embedded blocks and analysis by polymerase chain reaction.
J Histochem Cytochem 41: 765-8 (1993)
- Chen L; Ashe S; Singhal MC; Galloway DA; Hellstrom I; Hellstrom KE
Metastatic conversion of cells by expression of human papillomavirus type 16 E6 and E7 genes.
Proc Natl Acad Sci U S A 90: 6523-7 (1993)
- Chen SL; Han CP; Tsao YP
Construction of site-specific mutants of E5A protein of HPV-11 using the polymerase chain reaction and a single mutant primer.
J Virol Methods 42: 89-94 (1993)
- Chen SL; Han CP; Tsao YP; Lee JW; Yin CS
Identification and typing of human papillomavirus in cervical cancers in Taiwan.
Cancer 72: 1939-45 (1993)
- Chen TM; Chen CA; Hsieh CY; Chang DY; Chen YH; Defendi V
The state of p53 in primary human cervical carcinomas and its effects in human papillomavirus-immortalized human cervical cells.
Oncogene 8: 1511-8 (1993)
- Chen TM; Pecoraro G; Defendi V
Genetic analysis of in vitro progression of human papillomavirus-transfected human cervical cells.
Cancer Res 53: 1167-71 (1993)
- Cherny DI; Malkov VA; Volodin AA; Frank-Kamenetskii MD
Electron microscopy visualization of oligonucleotide binding to duplex DNA via triplex formation.
J Mol Biol 230: 379-83 (1993)
- Chinami M; Moriyama K; Fukumaki Y; Terada M; Shingu M
Association of RNA with human papillomavirus E7 protein of type 16 but not type 6b.
Biochem Biophys Res Commun 197: 1609-14 (1993)
- Choo CK; Rorke EA; Eckert RL
Calcium regulates the differentiation of human papillomavirus type 16 (HPV16) immortalized ectocervical epithelial cells, but not the expression of the papillomavirus E6 and E7 oncogenes.
Exp Cell Res 208: 161-9 (1993)
- Choo KB; Liew LN; Liew SJ; Cheng WT
Hypermethylation of the human papillomavirus transgenome in transgenic mice.
Proc Natl Sci Counc Repub China B 17: 159-63 (1993)
- Cid A; Auewarakul P; Garcia-Carranca A; Ovseiovich R; Gaisert H; Gissmann L
Cell-type-specific activity of the human papillomavirus type 18 upstream regulatory region in transgenic mice and its modulation by tetradecanoyl phorbol acetate and glucocorticoids.
J Virol 67: 6742-52 (1993)

- Claiborne A; Miller H; Parsonage D; Ross RP
Protein-sulfenic acid stabilization and function in enzyme catalysis and gene regulation.
FASEB J 7: 1483-90 (1993)
- Cohen BD; Goldstein DJ; Rutledge L; Vass WC; Lowy DR; Schlegel R; Schiller JT
Transformation-specific interaction of the bovine papillomavirus E5 oncoprotein with the platelet-derived growth factor receptor transmembrane domain and the epidermal growth factor receptor cytoplasmic domain.
J Virol 67: 5303-11 (1993)
- Cohen BD; Lowy DR; Schiller JT
The conserved C-terminal domain of the bovine papillomavirus E5 oncoprotein can associate with an alpha-adaptin-like molecule: a possible link between growth factor receptors and viral transformation.
Mol Cell Biol 13: 6462-8 (1993)
- Coleman N; Greenfield IM; Hare J; Kruger-Gray H; Chain BM; Stanley MA
Characterization and functional analysis of the expression of intercellular adhesion molecule-1 in human papillomavirus-related disease of cervical keratinocytes.
Am J Pathol 143: 355-67 (1993)
- Compton T
An immortalized human fibroblast cell line is permissive for human cytomegalovirus infection.
J Virol 67: 3644-8 (1993)
- Conrad M; Bubb VJ; Schlegel R
The human papillomavirus type 6 and 16 E5 proteins are membrane-associated proteins which associate with the 16-kilodalton pore-forming protein.
J Virol 67: 6170-8 (1993)
- Contorni M; Leoncini P
Typing of human papillomavirus DNAs by restriction endonuclease mapping of the PCR products.
J Virol Methods 41: 29-36 (1993)
- Corina K; Grossman SR; Barsoum J; Prakash SS; Androphy EJ; Pepinsky RB
The tryptophan bridge is a critical feature of the papillomavirus E2 DNA binding domain.
Virology 197: 391-6 (1993)
- Cossum PA; Sasmor H; Dellinger D; Truong L; Cummins L; Owens SR; Markham PM; Shea JP; Crooke S
Disposition of the 14C-labeled phosphorothioate oligonucleotide ISIS 2105 after intravenous administration to rats.
J Pharmacol Exp Ther 267: 1181-90 (1993)
- Cowsert LM; Fox MC; Zon G; Mirabelli CK
In vitro evaluation of phosphorothioate oligonucleotides targeted to the E2 mRNA of papillomavirus: potential treatment for genital warts.
Antimicrob Agents Chemother 37: 171-7 (1993)
- Cox M; Maitland N; Scully C
Human herpes simplex-1 and papillomavirus type 16 homologous DNA sequences in normal, potentially malignant and malignant oral mucosa.
Eur J Cancer B Oral Oncol 29B: 215-9 (1993)
- Cromme FV; Snijders PJ; van den Brule AJ; Kenemans P; Meijer CJ; Walboomers JM
MHC class I expression in HPV 16 positive cervical carcinomas is post-transcriptionally controlled and independent from c-myc overexpression.
Oncogene 8: 2969-75 (1993)
- Cuthill S; Sibbet GJ; Campo MS
Characterization of a nuclear factor, papilloma enhancer binding factor-1, that binds the long control region of human papillomavirus type 16 and contributes to enhancer activity.
Mol Carcinog 8: 96-104 (1993)
- Czegledy J; Rylander E; Evander M; Wadell G
Relation between the presence of human papillomavirus type 16 deoxyribonucleic acid in cervicovaginal cells and general health condition.
Am J Obstet Gynecol 169: 386-8 (1993)
- DasGupta S; Shivakumar CV; Das GC
Identification of proteins binding to the F441 locus of polyomavirus B enhancer that are required for its activity in embryonic carcinoma cells.
J Gen Virol 74 (Pt 4): 597-605 (1993)
- Davies R; Hicks R; Crook T; Morris J; Vousden K
Human papillomavirus type 16 E7 associates with a histone H1 kinase and with p107 through sequences necessary for transformation.
J Virol 67: 2521-8 (1993)
- De Geest K; Turyk ME; Hosken MI; Hudson JB; Laimins LA; Wilbanks GD
Growth and differentiation of human papillomavirus type 31b positive human cervical cell lines.
Gynecol Oncol 49: 303-10 (1993)
- De Marco F; Marcante ML
HPV-16 E6-E7 differential transcription induced in Siha cervical cancer cell line by interferons.
J Biol Regul Homeost Agents 7: 15-21 (1993)

References (1993)

- Deau MC; Favre M; Jablonska S; Rueda LA; Orth G
Genetic heterogeneity of oncogenic human papillomavirus type 5 (HPV5) and phylogeny of HPV5 variants associated with epidermodysplasia verruciformis.
J Clin Microbiol 31: 2918-26 (1993)
- Defeo-Jones D; Vuocolo GA; Haskell KM; Hanobik MG; Kiefer DM; McAvoy EM; Ivey-Hoyle M; Brandsma JL; Oliff A; Jones RE
Papillomavirus E7 protein binding to the retinoblastoma protein is not required for viral induction of warts.
J Virol 67: 716-25 (1993)
- Demeter LM; Stoler MH; Bonnez W; Corey L; Pappas P; Strussenberg J; Reichman RC
Penile intraepithelial neoplasia: clinical presentation and an analysis of the physical state of human papillomavirus DNA.
J Infect Dis 168: 38-46 (1993)
- Dhanwada KR; Garrett L; Smith P; Thompson KD; Doster A; Jones C
Characterization of human keratinocytes transformed by high risk human papillomavirus types 16 or 18 and herpes simplex virus type 2.
J Gen Virol 74 (Pt 6): 955-63 (1993)
- DiPaolo JA; Popescu NC; Alvarez L; Woodworth CD
Cellular and molecular alterations in human epithelial cells transformed by recombinant human papillomavirus DNA.
Crit Rev Oncog 4: 337-60 (1993)
- DiPaolo JA; Woodworth CD
Molecular control of human papillomavirus RNA expression in neoplasia.
Basic Life Sci 61: 239-46 (1993)
- Dillner L; Fredriksson A; Persson E; Forslund O; Hansson BG; Dillner J
Antibodies against papillomavirus antigens in cervical secretions from condyloma patients.
J Clin Microbiol 31: 192-7 (1993)
- Dodd JG; Paraskevas M; McNicol PJ
Detection of human papillomavirus 16 transcription in human prostate tissue.
J Urol 149: 400-2 (1993)
- Dollard SC; Broker TR; Chow LT
Regulation of the human papillomavirus type 11 E6 promoter by viral and host transcription factors in primary human keratinocytes.
J Virol 67: 1721-6 (1993)
- Donaldson YK; Arends MJ; Duvall E; Bird CC
PCR analysis of the upstream regulatory region of human papillomavirus genomes in cervical intraepithelial neoplasia and cervical carcinoma.
J Clin Pathol 46: 1021-3 (1993)
- Donaldson YK; Arends MJ; Duvall E; Bird CC
A PCR approach to discriminate between integrated and episomal HPV DNA in small clinical specimens.
Mol Cell Probes 7: 285-92 (1993)
- Egawa K; Delius H; Matsukura T; Kawashima M; de Villiers EM
Two novel types of human papillomavirus, HPV 63 and HPV 65: comparisons of their clinical and histological features and DNA sequences to other HPV types.
Virology 194: 789-99 (1993)
- Eriksson PO; Sahlman L
1H NMR studies of the mercuric ion binding protein MerP: sequential assignment, secondary structure and global fold of oxidized MerP.
J Biomol NMR 3: 613-26 (1993)
- Espy MJ; Smith TF; Persing DH
Dependence of polymerase chain reaction product inactivation protocols on amplicon length and sequence composition.
J Clin Microbiol 31: 2361-5 (1993)
- Falcinelli C; Luzi P; Alberti P; Cosmi EV; Anceschi MM
Human papilloma virus infection and Ki-ras oncogene in paraffin-embedded squamous carcinomas of the cervix.
Gynecol Obstet Invest 36: 185-8 (1993)
- Falcinelli C; van Belkum A; Schrauwen L; Seldenrijk K; Quint WG
Absence of human papillomavirus type 16 E6 transcripts in HPV 16-infected, cytologically normal cervical scrapings.
J Med Virol 40: 261-5 (1993)
- Fang BS; Guedes AC; Munoz LC; Villa LL
Human papillomavirus type 16 variants isolated from vulvar Bowenoid papulosis.
J Med Virol 41: 49-54 (1993)
- Farghaly SA
Correlation between HPV positivity and state of the p53 gene in cervical carcinoma cell lines [letter]
Gynecol Oncol 51: 288-9 (1993)

- Fattaey AR; Helin K; Dembski MS; Dyson N; Harlow E; Vuocolo GA; Hanobik MG; Haskell KM; Oliff A; Defeo-Jones D; et al
Characterization of the retinoblastoma binding proteins RBP1 and RBP2.
Oncogene 8: 3149-56 (1993)
- Faulkner-Jones BE; Tabrizi SN; Borg AJ; Roche PJ; Haralambidis J; Coghlan JP; Garland SM
Detection of human papillomavirus DNA and mRNA using synthetic, type-specific oligonucleotide probes.
J Virol Methods 41: 277-96 (1993)
- Felix DH; Jalal H; Cubie HA; Southam JC; Wray D; Maitland NJ
Detection of Epstein-Barr virus and human papillomavirus type 16 DNA in hairy leukoplakia by in situ hybridisation and the polymerase chain reaction.
J Oral Pathol Med 22: 277-81 (1993)
- Feltkamp MC; Smits HL; Vierboom MP; Minnaar RP; de Jongh BM; Drijfhout JW; ter Schegget J; Melief CJ; Kast WM
Vaccination with cytotoxic T lymphocyte epi-tope-containing peptide protects against a tumor induced by human papillomavirus type 16-transformed cells.
Eur J Immunol 23: 2242-9 (1993)
- Figge J; Breese K; Vajda S; Zhu QL; Eisele L; Andersen TT; MacColl R; Friedrich T; Smith TF
The binding domain structure of retinoblastoma-binding proteins.
Protein Sci 2: 155-64 (1993)
- Fimiani M; Mazzatenta C; Biagioli M; Andreassi L
Vulvar squamous papillomatosis and human papillomavirus infection. A polymerase chain reaction study.
Arch Dermatol Res 285: 250-4 (1993)
- Forslund O; Hansson BG; Rymark P; Bjerre B
Human papillomavirus DNA in urine samples compared with that in simultaneously collected urethra and cervix samples.
J Clin Microbiol 31: 1975-9 (1993)
- Franchini G; Mulloy JC; Koralknik IJ; Lo Monaco A; Sparkowski JJ; Andresson T; Goldstein DJ; Schlegel R
The human T-cell leukemia/lymphotropic virus type I p12I protein cooperates with the E5 oncoprotein of bovine papillomavirus in cell transformation and binds the 16-kilodalton subunit of the vacuolar H⁺ ATPase.
J Virol 67: 7701-4 (1993)
- Frazer IH; Leonard JH; Schonrock J; Wright RG; Kearsley JH
HPV DNA in oropharyngeal squamous cell cancers: comparison of results from four DNA detection methods.
Pathology 25: 138-43 (1993)
- Furihata M; Inoue K; Ohtsuki Y; Hashimoto H; Terao N; Fujita Y
High-risk human papillomavirus infections and overexpression of p53 protein as prognostic indicators in transitional cell carcinoma of the urinary bladder.
Cancer Res 53: 4823-7 (1993)
- Furihata M; Ohtsuki Y; Ogoshi S; Takahashi A; Tamiya T
The detection of either human papillomavirus genomes (types 16 and 18) or overexpressed p53 protein in esophageal cancers patients from Kochi, Japan [letter]
Am J Gastroenterol 88: 471-2 (1993)
- Furihata M; Ohtsuki Y; Ogoshi S; Takahashi A; Tamiya T; Ogata T
Prognostic significance of human papillomavirus genomes (type-16, -18) and aberrant expression of p53 protein in human esophageal cancer.
Int J Cancer 54: 226-30 (1993)
- Garcia-Carranca A; Gariglio PV
[Molecular aspects of human papillomaviruses and their relation to uterine cervix cancer]
Rev Invest Clin 45: 85-92 (1993)
- Gerber H; Antczak DF
Viruses, tumours and the MHC [editorial; comment]
Equine Vet J 25: 395 (1993)
- Ghim SJ; Jenson AB
Identification of conformational epitopes of the BPV-1 capsid recognized by competitive inhibition of sera from infected or immunized animals.
Pathobiology 61: 138-44 (1993)
- Gilles C; Piette J; Rombouts S; Laurent C; Foidart JM
Immortalization of human cervical keratinocytes by human papillomavirus type 33.
Int J Cancer 53: 872-9 (1993)
- Girardi F; Pickel H; Beyer-Finkler E; Pfister H
Specific rearrangements of human papillomavirus DNA provide molecular evidence for genetic heterogeneity of primary cervical cancers, recurrences, and lymph node metastases in two patients.
Gynecol Oncol 51: 281-6 (1993)

References (1993)

- Glew SS; Duggan-Keen M; Ghosh AK; Ivinson A; Sinnott P; Davidson J; Dyer PA; Stern PL
Lack of association of HLA polymorphisms with human papillomavirus-related cervical cancer.
Hum Immunol 37: 157-64 (1993)
- Griep AE; Herber R; Jeon S; Lohse JK; Dubielzig RR; Lambert PF
Tumorigenicity by human papillomavirus type 16 E6 and E7 in transgenic mice correlates with alterations in epithelial cell growth and differentiation.
J Virol 67: 1373-84 (1993)
- Hagensee ME; Yaegashi N; Galloway DA
Self-assembly of human papillomavirus type 1 capsids by expression of the L1 protein alone or by coexpression of the L1 and L2 capsid proteins.
J Virol 67: 315-22 (1993)
- Haskell KM; Vuocolo GA; Defeo-Jones D; Jones RE; Ivey-Hoyle M
Comparison of the binding of the human papillomavirus type 16 and cottontail rabbit papillomavirus E7 proteins to the retinoblastoma gene product.
J Gen Virol 74 (Pt 1): 115-9 (1993)
- Hasuo Y; Chua KL; Hjerpe A
Prevalence in Sweden of human papillomavirus type 16 infection in CIN II-III cases assessed by a PCR method.
Kurume Med J 40: 147-52 (1993)
- He YK; Lu CD; Qi GR
In vitro cleavage of HPV16 E6 and E7 RNA fragments by synthetic ribozymes and transcribed ribozymes from RNA-trimming plasmids.
FEBS Lett 322: 21-4 (1993)
- Heino P; Goldman S; Lagerstedt U; Dillner J
Molecular and serological studies of human papillomavirus among patients with anal epidermoid carcinoma.
Int J Cancer 53: 377-81 (1993)
- Helland A; Holm R; Kristensen G; Kaern J; Karlsen F; Trophe C; Nesland JM; Borresen AL
Genetic alterations of the TP53 gene, p53 protein expression and HPV infection in primary cervical carcinomas.
J Pathol 171: 105-14 (1993)
- Hillman RJ; Ryait BK; Botcherby M; Taylor-Robinson D
Human papillomavirus DNA in the urogenital tracts of men with gonorrhoea, penile warts or genital dermatoses.
Genitourin Med 69: 187-92 (1993)
- Hippelainen M; Eskelinen M; Lipponen P; Chang F; Syrjanen K
Mitotic activity index, volume corrected mitotic index and human papilloma-virus suggestive morphology are not prognostic factors in carcinoma of the oesophagus.
Anticancer Res 13: 677-81 (1993)
- Ho L; Chan SY; Burk RD; Das BC; Fujinaga K; Icenogle JP; Kahn T; Kiviat N; Lancaster W; Mavromara-Nazos P; et al
The genetic drift of human papillomavirus type 16 is a means of reconstructing prehistoric viral spread and the movement of ancient human populations.
J Virol 67: 6413-23 (1993)
- Ho L; Tay SK; Chan SY; Bernard HU
Sequence variants of human papillomavirus type 16 from couples suggest sexual transmission with low infectivity and polyclonality in genital neoplasia.
J Infect Dis 168: 803-9 (1993)
- Hoppe-Seyley F; Butz K
A novel cis-stimulatory element maps to the 5' portion of the human papillomavirus type 18 upstream regulatory region and is functionally dependent on a sequence-aberrant Sp1 binding site.
J Gen Virol 74 (Pt 2): 281-6 (1993)
- Hoppe-Seyley F; Butz K
Repression of endogenous p53 transactivation function in HeLa cervical carcinoma cells by human papillomavirus type 16 E6, human mdm-2, and mutant p53.
J Virol 67: 3111-7 (1993)
- Hording U; Kringsholm B; Andreasson B; Visfeldt J; Daugaard S; Bock JE
Human papillomavirus in vulvar squamous-cell carcinoma and in normal vulvar tissues: a search for a possible impact of HPV on vulvar cancer prognosis.
Int J Cancer 55: 394-6 (1993)
- Hording U; Rygaard C; Ruge S; Felding C; Lundvall F; Junge J
Cervical koilocytosis and high risk HPV types: the benefit of laser vaporization.
Eur J Obstet Gynecol Reprod Biol 51: 125-30 (1993)
- Horn S; Pfister H; Fuchs PG
Constitutive transcriptional activator of Epidermodysplasia verruciformis-associated human papillomavirus 8.
Virology 196: 674-81 (1993)
- Hsu EM; McNicol PJ; Guijon FB; Paraskevas M
Quantification of HPV-16 E6-E7 transcription in cervical intraepithelial neoplasia by reverse transcriptase polymerase chain reaction.
Int J Cancer 55: 397-401 (1993)

- Hsu WP; Ho ES; Yang CH; Lee YM; Chang CM; Liu WT
Prognostic significance of human papillomavirus detected with polymerase chain reaction in cervical cancer.
Chung Hua I Hsueh Tsa Chih 51: 97-102 (1993)
- Huang PS; Patrick DR; Edwards G; Goodhart PJ; Huber HE; Miles L; Garsky VM; Oliff A; Heimbrook DC
Protein domains governing interactions between E2F, the retinoblastoma gene product, and human papillomavirus type 16 E7 protein.
Mol Cell Biol 13: 953-60 (1993)
- Huang SL; Liu ZQ; Zhang XN; Li MJ
Human papillomavirus, human cytomegalovirus and onco-gene C-myc in cervical carcinoma and cervicitis.
Chin Med J (Engl) 106: 208-10 (1993)
- Hubert WG; Lambert PF
The 23-kilodalton E1 phosphoprotein of bovine papillomavirus type 1 is nonessential for stable plasmid replication in murine C127 cells.
J Virol 67: 2932-7 (1993)
- Hughes FJ; Romanos MA
E1 protein of human papillomavirus is a DNA helicase/ATPase.
Nucleic Acids Res 21: 5817-23 (1993)
- Huibregtse JM; Scheffner M; Howley PM
Localization of the E6-AP regions that direct human papillomavirus E6 binding, association with p53, and ubiquitination of associated proteins.
Mol Cell Biol 13: 4918-27 (1993)
- Huibregtse JM; Scheffner M; Howley PM
Cloning and expression of the cDNA for E6-AP, a protein that mediates the interaction of the human papillomavirus E6 oncoprotein with p53.
Mol Cell Biol 13: 775-84 (1993)
- Hwang ES; Riese DJ 2d; Settleman J; Nilson LA; Honig J; Flynn S; DiMaio D
Inhibition of cervical carcinoma cell line proliferation by the introduction of a bovine papillomavirus regulatory gene.
J Virol 67: 3720-9 (1993)
- Ibaraki T; Satake M; Kurai N; Ichijo M; Ito Y
Transacting activities of the E7 genes of several types of human papillomavirus.
Virus Genes 7: 187-96 (1993)
- Inoue H; Miyasaka M; Inoue T; Hakura A
Suppression of viral transformation in rat embryo fibroblast cells immortalized with transforming genes of human papillomavirus type 16.
Virology 195: 757-61 (1993)
- Inoue M
[Genetic alterations and transformations in development and establishment of uterine endometrial carcinomas]
Nippon Sanka Fujinka Gakkai Zasshi 45: 751-62 (1993)
- Iwasaka T; Oh-uchida M; Matsuo N; Yokoyama M; Fukuda K; Hara K; Fukuyama K; Hori K; Sugimori H
Correlation between HPV positivity and state of the p53 gene in cervical carcinoma cell lines.
Gynecol Oncol 48: 104-9 (1993)
- Iwasaka T; Yokoyama M; Hayashi Y; Sugimori H
Human papillomavirus 16 and 18 DNA can solely induce oncogenic transformation of mammalian cells in primary culture.
Acta Obstet Gynecol Scand 72: 81-6 (1993)
- Iwasawa A; Kumamoto Y; Fujinaga K
Detection of human papillomavirus deoxyribonucleic acid in penile carcinoma by polymerase chain reaction and in situ hybridization.
J Urol 149: 59-63 (1993)
- Jackson ME; Campo MS; Gaukroger JM
Cooperation between papillomavirus and chemical cofactors in oncogenesis.
Crit Rev Oncog 4: 277-91 (1993)
- Jackson P; Bos E; Braithwaite AW
Wild-type mouse p53 down-regulates transcription from different virus enhancer/promoters.
Oncogene 8: 589-97 (1993)
- Jakate SM; Saclarides TJ
Immunohistochemical detection of mutant P53 protein and human papillomavirus-related E6 protein in anal cancers.
Dis Colon Rectum 36: 1026-9 (1993)
- Jee SH; Kuo YF; Chen Y; Cheung SH; Chiu HC; Lee SC
Characterization of a human keratinocyte cell line immortalized by human papillomavirus 16 DNA.
J Formos Med Assoc 92: 876-83 (1993)
- Jiang D; Srinivasan A; Lozano G; Robbins PD
SV40 T antigen abrogates p53-mediated transcriptional activity.
Oncogene 8: 2805-12 (1993)

References (1993)

- Jochmus I; Durst M; Reid R; Altmann A; Bijward KE; Gissmann L; Jenson AB
Major histocompatibility complex and human papillomavirus type 16 E7 expression in high-grade vulvar lesions. *Hum Pathol* 24: 519-24 (1993)
- Jullian EH; Dhellemmes C; Saglio O; Chavinie J; Pompidou A
Improved detection of human papillomavirus types 16 and 18 in cervical scrapes by a multiplex polymerase chain reaction: a 4% prevalence among 120 French women with normal cytology. *Lab Invest* 68: 242-7 (1993)
- Karlen S; Beard P
Identification and characterization of novel promoters in the genome of human papillomavirus type 18. *J Virol* 67: 4296-306 (1993)
- Karlen S; Offord EA; Beard P
Herpes simplex virions interfere with the expression of human papillomavirus type 18 genes. *J Gen Virol* 74 (Pt 6): 965-73 (1993)
- Kasher MS; Wakulchik M; Cook JA; Smith MC
One-step purification of recombinant human papillomavirus type 16 E7 oncoprotein and its binding to the retinoblastoma gene product. *Biotechniques* 14: 630-41 (1993)
- Kasperbauer JL; O'Halloran GL; Espy MJ; Smith TF; Lewis JE
Polymerase chain reaction (PCR) identification of human papillomavirus (HPV) DNA in verrucous carcinoma of the larynx. *Laryngoscope* 103: 416-20 (1993)
- Kast WM; Brandt RM; Drijfhout JW; Melief CJ
Human leukocyte antigen-A2.1 restricted candidate cytotoxic T lymphocyte epitopes of human papillomavirus type 16 E6 and E7 proteins identified by using the processing-defective human cell line T2. *J Immunother* 14: 115-20 (1993)
- Kenter GG; Cornelisse CJ; Jiwa NM; Aartsen EJ; Hermans J; Mooi W; Heintz AP; Fleuren GJ
Human papillomavirus type 16 in tumor tissue of low-stage squamous carcinoma of the uterine cervix in relation to ploidy grade and prognosis. *Cancer* 71: 397-401 (1993)
- Kessis TD; Slebos RJ; Han SM; Shah K; Bosch XF; Munoz N; Hedrick L; Cho KR
p53 gene mutations and MDM2 amplification are uncommon in primary carcinomas of the uterine cervix. *Am J Pathol* 143: 1398-405 (1993)
- Kessis TD; Slebos RJ; Nelson WG; Kastan MB; Plunkett BS; Han SM; Lorincz AT; Hedrick L; Cho KR
Human papillomavirus 16 E6 expression disrupts the p53-mediated cellular response to DNA damage. *Proc Natl Acad Sci U S A* 90: 3988-92 (1993)
- Khan MA; Jenkins GR; Tolleson WH; Creek KE; Pirisi L
Retinoic acid inhibition of human papillomavirus type 16-mediated transformation of human keratinocytes. *Cancer Res* 53: 905-9 (1993)
- Khan MA; Tolleson WH; Gangemi JD; Pirisi L
Inhibition of growth, transformation, and expression of human papillomavirus type 16 E7 in human keratinocytes by alpha interferons. *J Virol* 67: 3396-403 (1993)
- Kiessling U; Kube D; Janda J; Bergmann S; Johannsmeyer KD; Hauff P; Arnoldt W; Waehlte H
Properties of two epithelial cell lines derived from HPV-associated cervical and vulvar lesions. *Eur J Cancer* 29A: 1746-53 (1993)
- Kim MS; Shin KH; Baek JH; Cherrick HM; Park NH
HPV-16, tobacco-specific N-nitrosamine, and N-methyl-N'-nitro-N-nitrosoguanidine in oral carcinogenesis. *Cancer Res* 53: 4811-6 (1993)
- Kirnbauer R; Taub J; Greenstone H; Roden R; Durst M; Gissmann L; Lowy DR; Schiller JT
Efficient self-assembly of human papillomavirus type 16 L1 and L1-L2 into virus-like particles. *J Virol* 67: 6929-36 (1993)
- Klingelutz AJ; Smith PP; Garrett LR; McDougall JK
Alteration of the DCC tumor-suppressor gene in tumorigenic HPV-18 immortalized human keratinocytes transformed by nitrosomethylurea. *Oncogene* 8: 95-9 (1993)
- Koralnik IJ; Fullen J; Franchini G
The p12I, p13II, and p30II proteins encoded by human T-cell leukemia/lymphotropic virus type I open reading frames I and II are localized in three different cellular compartments. *J Virol* 67: 2360-6 (1993)
- Krajinovic M; Lazic J; Stanimirovic B; Diklic V; Savic A
The E2 region of HPV 16 in relation to different types of cervical lesions. *J Med Virol* 41: 1-5 (1993)

- Krebnak V; Pistek T; Vagner J; Suchankova A; Kanka J; Ritterova L; Vonka V
Identification of seroreactive epitopes of human papillomavirus type 18 E7 protein by synthetic peptides.
Acta Virol 37: 395-402 (1993)
- Kurz J; Mitra K; Adam R; Miao X; MacKay JS; Isa NN; Coombs DH; Krause MO
PCR detection and typing of genital papillomavirus in a New Brunswick population.
Int J Cancer 55: 604-8 (1993)
- Kyo S; Inoue M; Nishio Y; Nakanishi K; Akira S; Inoue H; Yutsudo M; Tanizawa O; Hakura A
NF-IL6 represses early gene expression of human papillomavirus type 16 through binding to the noncoding region.
J Virol 67: 1058-66 (1993)
- Laimins LA
The biology of human papillomaviruses: from warts to cancer.
Infect Agents Dis 2: 74-86 (1993)
- Lambert PF; Pan H; Pitot HC; Liem A; Jackson M; Griep AE
Epidermal cancer associated with expression of human papillomavirus type 16 E6 and E7 oncogenes in the skin of transgenic mice.
Proc Natl Acad Sci U S A 90: 5583-7 (1993)
- Lee KR; Howard P; Heintz NH; Collins CC
Low prevalence of human papillomavirus types 16 and 18 in cervical adenocarcinoma in situ, invasive adenocarcinoma, and glandular dysplasia by polymerase chain reaction.
Mod Pathol 6: 433-7 (1993)
- Lee NK; Ye YW; Chen J; Li X; Waber PG; Nisen PD
p53, retinoblastoma, and human papillomavirus in squamous cell carcinoma and adjacent normal mucosa of the upper aerodigestive tract.
Arch Otolaryngol Head Neck Surg 119: 1125-31 (1993)
- Lentz MR; Pak D; Mohr I; Botchan MR
The E1 replication protein of bovine papillomavirus type 1 contains an extended nuclear localization signal that includes a p34cdc2 phosphorylation site.
J Virol 67: 1414-23 (1993)
- Lewensohn-Fuchs I; Wester D; Bistoletti P; Elfgren K; Ohlman S; Dillner J; Dalianis T
Serological responses to human papillomavirus type 16 antigens in women before and after renal transplantation.
J Med Virol 40: 188-92 (1993)
- Li R; Botchan MR
The acidic transcriptional activation domains of VP16 and p53 bind the cellular replication protein A and stimulate in vitro BPV-1 DNA replication.
Cell 73: 1207-21 (1993)
- Lin YL; Borenstein LA; Ahmed R; Wettstein FO
Cottontail rabbit papillomavirus L1 protein-based vaccines: protection is achieved only with a full-length, nondenatured product.
J Virol 67: 4154-62 (1993)
- Lopez E; Chypre C; Alpha B; Mathis G
Europium(III) trisbipyridine cryptate label for time-resolved fluorescence detection of polymerase chain reaction products fixed on a solid support.
Clin Chem 39: 196-201 (1993)
- Lopez Ocejio O; Perea SE; Reyes A; Vigoa L; Lopez Saura P
Partial phenotypic reversion of HeLa cells by long-term interferon-alpha treatment.
J Interferon Res 13: 369-75 (1993)
- Lu JZ; Sun YN; Rose RC; Bonnez W; McCance DJ
Two E2 binding sites (E2BS) alone or one E2BS plus an A/T-rich region are minimal requirements for the replication of the human papillomavirus type 11 origin.
J Virol 67: 7131-9 (1993)
- Lucotte G; Francois MH; Petit MC; Berriche S; Reveilleau S
A multiple primer pairs polymerase chain reaction for the detection of human genital papillomavirus types.
Mol Cell Probes 7: 339-44 (1993)
- Lusky M; Hurwitz J; Seo YS
Cooperative assembly of the bovine papilloma virus E1 and E2 proteins on the replication origin requires an intact E2 binding site.
J Biol Chem 268: 15795-803 (1993)
- Maher DW; Strawn LM; Donoghue DJ
Alanine mutagenesis of conserved residues in the platelet-derived growth factor family: identification of residues necessary for dimerization and transformation.
Oncogene 8: 533-41 (1993)
- Maiorano E; Favia GF; Bufo P
Epidermal growth factor receptor expression in squamous papillomas of the oral mucosa.
Boll Soc Ital Biol Sper 69: 195-202 (1993)
- Mansur CP; Androphy EJ
Cellular transformation by papillomavirus oncoproteins.
Biochim Biophys Acta 1155: 323-45 (1993)

References (1993)

- Margall N; Matias-Guiu X; Chillon M; Coll P; Alejo M; Nunes V; Quilez M; Rabella N; Prats G; Prat J
Detection of human papillomavirus 16 and 18 DNA in epithelial lesions of the lower genital tract by in situ hybridization and polymerase chain reaction: cervical scrapes are not substitutes for biopsies.
J Clin Microbiol 31: 924-30 (1993)
- Marti E; Lazary S; Antczak DF; Gerber H
Report of the first international workshop on equine sarcoid [see comments]
Equine Vet J 25: 397-407 (1993)
- Matsuo N; Iwasaka T; Hayashi Y; Hara K; Mvula M; Sugimori H
Polymerase chain reaction analysis of human papillomavirus in adenocarcinoma and adenosquamous carcinoma of the uterine cervix.
Int J Gynaecol Obstet 41: 251-6 (1993)
- McIntyre MC; Frattini MG; Grossman SR; Laimins LA
Human papillomavirus type 18 E7 protein requires intact Cys-X-X-Cys motifs for zinc binding, dimerization, and transformation but not for Rb binding.
J Virol 67: 3142-50 (1993)
- McLean CS; Sterling JS; Mowat J; Nash AA; Stanley MA
Delayed-type hypersensitivity response to the human papillomavirus type 16 E7 protein in a mouse model.
J Gen Virol 74 (Pt 2): 239-45 (1993)
- Medcalf EA; Milner J
Targeting and degradation of p53 by E6 of human papillomavirus type 16 is preferential for the 1620+ p53 conformation.
Oncogene 8: 2847-51 (1993)
- Meissner J
TaqI is a single cut enzyme for HPV-18.
Nucleic Acids Res 21: 1041 (1993)
- Melchers W; de Mare S; Kuitert E; Galama J; Walboomers J; van den Brule AJ
Human papillomavirus and cutaneous warts in meat handlers.
J Clin Microbiol 31: 2547-9 (1993)
- Merrick DT; Gown AM; Halbert CL; Blanton RA; McDougall JK
Human papillomavirus-immortalized keratinocytes are resistant to the effects of retinoic acid on terminal differentiation.
Cell Growth Differ 4: 831-40 (1993)
- Mittal R; Pater A; Pater MM
Multiple human papillomavirus type 16 glucocorticoid response elements functional for transformation, transient expression, and DNA-protein interactions.
J Virol 67: 5656-9 (1993)
- Mittal R; Tsutsumi K; Pater A; Pater MM
Human papillomavirus type 16 expression in cervical keratinocytes: role of progesterone and glucocorticoid hormones.
Obstet Gynecol 81: 5-12 (1993)
- Monini P; Blitz IL; Cassai E
Cooperative DNA binding of the bovine papillomavirus E2 transcriptional activator is antagonized by truncated E2 polypeptides.
J Virol 67: 5668-76 (1993)
- Monini P; de Lellis L; Borgatti P; Hassan-Omran M; Cassai E
Activation of eukaryotic transcriptional promoters by the bovine papillomavirus E1-replication factor.
Intervirology 36: 245-52 (1993)
- Morris JD; Crook T; Bandara LR; Davies R; LaThangue NB; Vousden KH
Human papillomavirus type 16 E7 regulates E2F and contributes to mitogenic signalling.
Oncogene 8: 893-8 (1993)
- Morris PJ; Ring CJ; Lillycrop KA; Latchman DS
Transactivation of the human papilloma virus 16 octamer motif by the octamer binding protein Oct-2 requires both the N and C terminal activation domains.
Nucleic Acids Res 21: 4506-10 (1993)
- Morton KA; Jones BJ; Sohn MH; Datz FL; Lynch RE
Enrichment for metallothionein does not confer resistance to cisplatin in transfected NIH/3T3 cells.
J Pharmacol Exp Ther 267: 697-702 (1993)
- Mukhopadhyay D; Chatterjee R; Chakraborty RN
Correlation of mitotic abnormalities and the presence of human papillomavirus antigens in squamous cell carcinomas of the oral cavity.
Cancer Lett 74: 51-6 (1993)
- Mulhaupt H; Bruder E; Elit L; Rothblat I; Warhol M
Combined analysis of cervical smears. Cytopathology, image cytometry and in situ hybridization.
Acta Cytol 37: 373-8 (1993)
- Munger K; Phelps WC
The human papillomavirus E7 protein as a transforming and transactivating factor.
Biochim Biophys Acta 1155: 111-23 (1993)

- Murakami T; Nagai N; Ota S; Ohama K
[Comparative study of HPV infection and proliferating cell nuclear antigen of uterine cervix]
Nippon Sanka Fujinka Gakkai Zasshi 45: 967-72 (1993)
- Newfield L; Goldsmith A; Bradlow HL; Auburn K
Estrogen metabolism and human papillomavirus-induced tumors of the larynx: chemo-prophylaxis with indole-3-carbinol.
Anticancer Res 13: 337-41 (1993)
- Nilson LA; DiMaio D
Platelet-derived growth factor receptor can mediate tumorigenic transformation by the bovine papillomavirus E5 protein.
Mol Cell Biol 13: 4137-45 (1993)
- Nishikawa T; Kobayashi H; Shindoh M; Yamashita T; Fujinaga K; Ohkawara A
A case of verrucous carcinoma associated with human papillomavirus type 16 DNA.
J Dermatol 20: 483-8 (1993)
- O'Banion MK; Winn VD; Settleman J; Young DA
Genetic definition of a new bovine papillomavirus type 1 open reading frame, E5B, that encodes a hydrophobic protein involved in altering host-cell protein processing.
J Virol 67: 3427-34 (1993)
- Offord EA; Chappuis PO; Beard P
Different stability of AP1 proteins in human keratinocyte and fibroblast cells: possible role in the cell-type specific expression of human papillomavirus type 18 genes.
Carcinogenesis 14: 2447-55 (1993)
- Ogunbiyi OA; Scholefield JH; Smith JH; Polaczar SV; Rogers K; Sharp F
Immunohistochemical analysis of p53 expression in anal squamous neoplasia.
J Clin Pathol 46: 507-12 (1993)
- Okabayashi M; Angell MG; Budgeon LR; Kreider JW
Shope papilloma cell and leukocyte proliferation in regressing and progressing lesions.
Am J Pathol 142: 489-96 (1993)
- Onda T; Kanda T; Zanma S; Yasugi T; Watanabe S; Kawana T; Ueda K; Yoshikawa H; Taketani Y; Yoshiike K
Association of the antibodies against human papillomavirus 16 E4 and E7 proteins with cervical cancer positive for human papillomavirus DNA.
Int J Cancer 54: 624-8 (1993)
- Ong CK; Chan SY; Campo MS; Fujinaga K; Mavromara-Nazos P; Labropoulou V; Pfister H; Tay SK; ter Meulen J; Villa LL; et al
Evolution of human papillomavirus type 18: an ancient phylogenetic root in Africa and intratype diversity reflect coevolution with human ethnic groups.
J Virol 67: 6424-31 (1993)
- Ostrow RS; Liu Z; Schneider JF; McGlennen RC; Forslund K; Faras AJ
The products of the E5, E6, or E7 open reading frames of RhPV 1 can individually transform NIH 3T3 cells or in cotransfections with activated ras can transform primary rodent epithelial cells.
Virology 196: 861-7 (1993)
- Otten N; von Tscharnher C; Lazary S; Antczak DF; Gerber H
DNA of bovine papillomavirus type 1 and 2 in equine sarcooids: PCR detection and direct sequencing.
Arch Virol 132: 121-31 (1993)
- Pahel G; Aulabaugh A; Short SA; Barnes JA; Painter GR; Ray P; Phelps WC
Structural and functional characterization of the HPV16 E7 protein expressed in bacteria.
J Biol Chem 268: 26018-25 (1993)
- Paquette RL; Lee YY; Wilczynski SP; Karmakar A; Kizaki M; Miller CW; Koeffler HP
Mutations of p53 and human papillomavirus infection in cervical carcinoma.
Cancer 72: 1272-80 (1993)
- Park DS; Selvey LA; Kelsall SR; Frazer IH
Human papillomavirus type 16 E6, E7 and L1 and type 18 E7 proteins produced by recombinant baculoviruses.
J Virol Methods 45: 303-18 (1993)
- Park JS; Namkoong SE; Han SK; Nha DJ; Lee HY; Kim SJ
Comparison of L1 consensus primers with E6 type specific primers for detection of human papillomaviruses in paraffin sections of cervical neoplasia.
J Korean Med Sci 8: 60-7 (1993)
- Pater A; Belaguli NS; Pater MM
Glucocorticoid requirement for growth of human papillomavirus 16-transformed primary rat kidney epithelial cells: correlation of development of hormone resistance with viral RNA expression and processing.
Cancer Res 53: 4432-6 (1993)
- Pei XF; Meck JM; Greenhalgh D; Schlegel R
Cotransfection of HPV-18 and v-fos DNA induces tumorigenicity of primary human keratinocytes.
Virology 196: 855-60 (1993)

References (1993)

- Pei XF; Wu M
Clonal growth and origin of two human keratinocyte cell lines transformed by human papillomavirus type 16 DNA. *Cancer Lett* 74: 15-24 (1993)
- Peng X; Olson RO; Christian CB; Lang CM; Kreider JW
Papillomas and carcinomas in transgenic rabbits carrying EJ-ras DNA and cottontail rabbit papillomavirus DNA. *J Virol* 67: 1698-701 (1993)
- Pennie WD; Grindlay GJ; Cairney M; Campo MS
Analysis of the transforming functions of bovine papillomavirus type 4. *Virology* 193: 614-20 (1993)
- Pilotti S; Donghi R; D'Amato L; Giarola M; Longoni A; Della Torre G; De Palo G; Pierotti MA; Rilke F
HPV detection and p53 alteration in squamous cell verrucous malignancies of the lower genital tract. *Diagn Mol Pathol* 2: 248-56 (1993)
- Poljak M; Cerar A
Human papillomavirus type 16 DNA in oesophageal squamous cell carcinoma. *Anticancer Res* 13: 2113-6 (1993)
- Pollanen R; Soini Y; Vahakangas K; Paakko P; Lehto VP
Aberrant p53 protein expression in cervical intra-epithelial neoplasia. *Histopathology* 23: 471-4 (1993)
- Prussia PR; ter Schegget J; Smits HL
Detection of oncogenic HPV DNA by a consensus polymerase chain reaction method in genital carcinomas in twenty women in Barbados. *West Indian Med J* 42: 144-6 (1993)
- Purdie KJ; Sexton CJ; Proby CM; Glover MT; Williams AT; Stables JN; Leigh IM
Malignant transformation of cutaneous lesions in renal allograft patients: a role for human papillomavirus. *Cancer Res* 53: 5328-33 (1993)
- Rady PL; Chin R; Arany I; Hughes TK; Tying SK
Direct sequencing of consensus primer generated PCR fragments of human papillomaviruses. *J Virol Methods* 43: 335-50 (1993)
- Reid R
Biology and colposcopic features of human papillomavirus-associated cervical disease. *Obstet Gynecol Clin North Am* 20: 123-51 (1993)
- Reinhartz A; Alajem S; Samson A; Herzberg M
A novel rapid hybridization technique: paper chromatography hybridization assay (PACHA). *Gene* 136: 221-6 (1993)
- Rho J; Park SD; Choe J
P3 promoter element of bovine papillomavirus. *Cell Mol Biol (Noisy-le-grand)* 39: 205-11 (1993)
- Roberts S; Ashmole I; Johnson GD; Kreider JW; Gallimore PH
Cutaneous and mucosal human papillomavirus E4 proteins form intermediate filament-like structures in epithelial cells. *Virology* 197: 176-87 (1993)
- Robertson DI; Paslawski D; Duggan MA; Stuart GC; Nation JG
Estrogen and progesterone receptor, human papillomavirus, and DNA ploidy analysis in invasive carcinoma of the cervix in pregnancy. *Am J Clin Pathol* 100: 18-21 (1993)
- Rogel-Gaillard C; Pehau-Arnaudet G; Breitburd F; Orth G
Cytopathic effect in human papillomavirus type 1-induced inclusion warts: in vitro analysis of the contribution of two forms of the viral E4 protein. *J Invest Dermatol* 101: 843-51 (1993)
- Rose RC; Bonnez W; Reichman RC; Garcea RL
Expression of human papillomavirus type 11 L1 protein in insect cells: in vivo and in vitro assembly of viruslike particles. *J Virol* 67: 1936-44 (1993)
- Rymark P; Forslund O; Hansson BG; Lindholm K
Genital HPV infection not a local but a regional infection: experience from a female teenage group. *Genitourin Med* 69: 18-22 (1993)
- Salloukh H; Lavery C; Smiley JR; Rawls WE
Construction and preliminary characterization of a non-defective herpes simplex virus recombinant bearing the genome of human papillomavirus type 16. *Can J Microbiol* 39: 111-7 (1993)
- Sandler AB; Vande Pol SB; Spalholz BA
Repression of bovine papillomavirus type 1 transcription by the E1 replication protein. *J Virol* 67: 5079-87 (1993)
- Sassi I; Mangili F; Luchini S; Colombo G; Pezzotta MG; Candiani M; Frigerio A; Santambrogio G; Cantaboni A; Vignali M
[Correlations between the HPV type, DNA index and evolution of cervical intraepithelial neoplasia (CIN)] *Pathologica* 85: 79-84 (1993)

- Scully C
Oral cancer: new insights into pathogenesis.
Dent Update 20: 95-100 (1993)
- Seo YS; Muller F; Lusky M; Gibbs E; Kim HY; Phillips B; Hurwitz J
Bovine papilloma virus (BPV)-encoded E2 protein enhances binding of E1 protein to the BPV replication origin.
Proc Natl Acad Sci U S A 90: 2865-9 (1993)
- Seo YS; Muller F; Lusky M; Hurwitz J
Bovine papilloma virus (BPV)-encoded E1 protein contains multiple activities required for BPV DNA replication.
Proc Natl Acad Sci U S A 90: 702-6 (1993)
- Sexton CJ; Proby CM; Banks L; Stables JN; Powell K; Navsaria H; Leigh IM
Characterization of factors involved in human papillomavirus type 16-mediated immortalization of oral keratinocytes.
J Gen Virol 74 (Pt 4): 755-61 (1993)
- Shay JW; Wright WE; Brasiskyte D; Van der Haegen BA
E6 of human papillomavirus type 16 can overcome the M1 stage of immortalization in human mammary epithelial cells but not in human fibroblasts.
Oncogene 8: 1407-13 (1993)
- Shervington A
T2G4 telomere repeats does not provide telomere function to a BPV-1 containing DNA fragment in mouse C127 cells.
Biochem Mol Biol Int 29: 661-72 (1993)
- Shikano K; Sasano H; Ito K; Ozawa N; Tase T; Sato S; Yajima A
The correlation between HPV infection and cell proliferative activity in uterine cervical adenocarcinoma.
Tohoku J Exp Med 169: 279-88 (1993)
- Sizemore N; Rorke EA
Human papillomavirus 16 immortalization of normal human ectocervical epithelial cells alters retinoic acid regulation of cell growth and epidermal growth factor receptor expression.
Cancer Res 53: 4511-7 (1993)
- Smith LH; Foster C; Hitchcock ME; Isseroff R
In vitro HPV-11 infection of human foreskin.
J Invest Dermatol 101: 292-5 (1993)
- Smith PR; Gao Y; Karran L; Jones MD; Snudden D; Griffin BE
Complex nature of the major viral polyadenylated transcripts in Epstein-Barr virus-associated tumors.
J Virol 67: 3217-25 (1993)
- Smits PH; Smits HL; Minnaar RP; ter Schegget J
Regulation of human papillomavirus type 16 (HPV-16) transcription by loci on the short arm of chromosome 11 is mediated by the TATAAAA motif of the HPV-16 promoter.
J Gen Virol 74 (Pt 1): 121-4 (1993)
- Soler C; Chardonnet Y; Allibert P; Euvrard S; Schmitt D; Mandrand B
Detection of mucosal human papillomavirus types 6/11 in cutaneous lesions from transplant recipients.
J Invest Dermatol 101: 286-91 (1993)
- Soyer HP; Schadendorf D; Cerroni L; Kerl H
Verrucous cysts: histopathologic characterization and molecular detection of human papillomavirus-specific DNA.
J Cutan Pathol 20: 411-7 (1993)
- Spalholz BA
Importance of the bovine papillomavirus P2443 promoter in the regulation of E2 and E5 expression.
J Virol 67: 6278-84 (1993)
- Spalholz BA; McBride AA; Sarafi T; Quintero J
Binding of bovine papillomavirus E1 to the origin is not sufficient for DNA replication.
Virology 193: 201-12 (1993)
- Spitkovskii DD; Roier GD; Mazurenko NN; Mikhaleva II; Prudchenko IA; Korbukh IA; Sukhova NM; Kiselev FL
[In vivo identification of YB-1 protein, interacting with the enhancer of human papillomavirus (HPV) type 18, using antibodies to a synthetic peptide]
Mol Biol (Mosk) 27: 81-91 (1993)
- Stacey SN; Ghosh A; Bartholomew JS; Tindle RW; Stern PL; Mackett M; Arrand JR
Expression of human papillomavirus type 16 E7 protein by recombinant baculovirus and use for the detection of E7 antibodies in sera from cervical carcinoma patients.
J Med Virol 40: 14-21 (1993)
- Steele C; Cowsert LM; Shillitoe EJ
Effects of human papillomavirus type 18-specific antisense oligonucleotides on the transformed phenotype of human carcinoma cell lines.
Cancer Res 53: 2330-7 (1993)

References (1993)

- Steele JC; Stankovic T; Gallimore PH
Production and characterization of human proliferative T-cell clones specific for human papillomavirus type 1 E4 protein.
J Virol 67: 2799-806 (1993)
- Storey A; Banks L
Human papillomavirus type 16 E6 gene cooperates with EJ-ras to immortalize primary mouse cells.
Oncogene 8: 919-24 (1993)
- Straight SW; Hinkle PM; Jewers RJ; McCance DJ
The E5 oncoprotein of human papillomavirus type 16 transforms fibroblasts and effects the downregulation of the epidermal growth factor receptor in keratinocytes.
J Virol 67: 4521-32 (1993)
- Sun Q; Tsutsumi K; Yokoyama M; Pater MM; Pater A
In vivo cytokeratin-expression pattern of stratified squamous epithelium from human papillomavirus-type-16-immortalized ectocervical and foreskin keratinocytes.
Int J Cancer 54: 656-62 (1993)
- Tamada Y
[The biological activities of the human papillomavirus type 16 E6/E7 cDNAs cloned from the SiHa cervical carcinoma cell line]
Hokkaido Igaku Zasshi 68: 849-61 (1993)
- Tamayo Legorreta EM; Echaniz Aviles G; Cruz Valdez A; Camacho Alcantara G; Calderon Jaimes E
[Human papillomavirus infection in women with and without abnormal cervical cytology]
Ginecol Obstet Mex 61: 27-34 (1993)
- Taniguchi A; Kikuchi K; Nagata K; Yasumoto S
A cell-type-specific transcription enhancer of type 16 human papillomavirus (HPV 16)-P97 promoter is defined with HPV-associated cellular events in human epithelial cell lines.
Virology 195: 500-10 (1993)
- ter Meulen J; Schweigler AC; Eberhardt HC; Luande J; Mgaya HN; Muller M; Bleul C; Ulken V; Ikenberg H; Pawlita M; et al
Sequence variation in the E7 gene of human papillomavirus type 18 in tumor and non-tumor patients and antibody response to a conserved seroreactive epitope.
Int J Cancer 53: 257-9 (1993)
- Tervahauta AI; Syrjanen SM; Vayrynen M; Saastamoinen J; Syrjanen KJ
Expression of p53 protein related to the presence of human papillomavirus (HPV) DNA in genital carcinomas and precancer lesions.
Anticancer Res 13: 1107-11 (1993)
- Thorner LK; Lim DA; Botchan MR
DNA-binding domain of bovine papillomavirus type 1 E1 helicase: structural and functional aspects.
J Virol 67: 6000-14 (1993)
- Tieben LM; ter Schegget J; Minnaar RP; Bouwes Bavink JN; Berkhout RJ; Vermeer BJ; Jebbink MF; Smits HL
Detection of cutaneous and genital HPV types in clinical samples by PCR using consensus primers.
J Virol Methods 42: 265-79 (1993)
- Tomita Y; Simizu B
Translational properties of the human papillomavirus type-6 L1-coding mRNA.
Gene 133: 223-5 (1993)
- Tommasino M; Adamczewski JP; Carlotti F; Barth CF; Manetti R; Contorni M; Cavalieri F; Hunt T; Crawford L
HPV16 E7 protein associates with the protein kinase p33CDK2 and cyclin A.
Oncogene 8: 195-202 (1993)
- Tornesello ML; Buonaguro FM; Beth-Giraldo E; Giraldo G
Human immunodeficiency virus type 1 tat gene enhances human papillomavirus early gene expression.
Intervirology 36: 57-64 (1993)
- Tosi P; Pallini V; Cintorino M; Bugnoli M; Covacci A; Petracca R; Syrjanen S; Del Vecchio MT; Mantyljarvi R; Ruggiero P; et al
Use of antibodies against a synthetic peptide of the E6 protein of human papillomavirus (HPV) type 16 for the diagnosis of genital HPV lesions.
Cytopathology 4: 3-15 (1993)
- Tsutsumi K; Hoshikawa T; Suzuki T; Takeyama I
[Expression of human papillomavirus (HPV) gene in HPV-positive laryngeal tumors and activity of the HPV long control region in cultured normal laryngeal epithelial cells]
Nippon Jibiinkoka Gakkai Kaiho 96: 767-73 (1993)
- Tucker RA; Johnson PR; Reeves WC; Icenogle JP
Using the polymerase chain reaction to genotype human papillomavirus DNAs in samples containing multiple HPV's may produce inaccurate results.
J Virol Methods 43: 321-33 (1993)

- Tyan YS; Liu ST; Ong WR; Chen ML; Shu CH; Chang YS
Detection of Epstein-Barr virus and human papillomavirus in head and neck tumors.
J Clin Microbiol 31: 53-6 (1993)
- Ustav E; Ustav M; Szymanski P; Stenlund A
The bovine papillomavirus origin of replication requires a binding site for the E2 transcriptional activator.
Proc Natl Acad Sci U S A 90: 898-902 (1993)
- Vambutas A; Di Lorenzo TP; Steinberg BM
Laryngeal papilloma cells have high levels of epidermal growth factor receptor and respond to epidermal growth factor by a decrease in epithelial differentiation.
Cancer Res 53: 910-4 (1993)
- Ventura AM; Villa LL
Plasmid pSH: a strong mammalian expression vector.
Biochem Biophys Res Commun 192: 867-9 (1993)
- Vernon SD; Hart CE; Reeves WC; Icenogle JP
The HIV-1 tat protein enhances E2-dependent human papillomavirus 16 transcription.
Virus Res 27: 133-45 (1993)
- Volpers C; Sapp M; Komly CA; Richalet-Secordel P; Streeck RE
Development of type-specific and cross-reactive serological probes for the minor capsid protein of human papillomavirus type 33.
J Virol 67: 1927-35 (1993)
- Vousden K
Interactions of human papillomavirus transforming proteins with the products of tumor suppressor genes.
FASEB J 7: 872-9 (1993)
- Vousden KH; Crook T; Farrell PJ
Biological activities of p53 mutants in Burkitt's lymphoma cells.
J Gen Virol 74 (Pt 5): 803-10 (1993)
- Vousden KH; Vojtesek B; Fisher C; Lane D
HPV-16 E7 or adenovirus E1A can overcome the growth arrest of cells immortalized with a temperature-sensitive p53.
Oncogene 8: 1697-702 (1993)
- Wang ZM
[Condyloma acuminatum: histopathology and detection of human papilloma virus]
Chung-hua Ping Li Hsueh Tsa Chih 22: 160-2 (1993)
- Watanabe S; Kanda T; Yoshiike K
Growth dependence of human papillomavirus 16 DNA-positive cervical cancer cell lines and human papillomavirus 16-transformed human and rat cells on the viral oncoproteins.
Jpn J Cancer Res 84: 1043-9 (1993)
- Watanabe S; Ogura H; Fukushima K; Yabe Y
Comparison of Virapap filter hybridization with polymerase chain reaction and Southern blot hybridization methods for detection of human papillomavirus in tonsillar and pharyngeal cancers.
Eur Arch Otorhinolaryngol 250: 115-9 (1993)
- Weinberg E; Hoisington S; Eastman AY; Rice DK; Malfetano J; Ross JS
Uterine cervical lymphoepithelial-like carcinoma. Absence of Epstein-Barr virus genomes.
Am J Clin Pathol 99: 195-9 (1993)
- Wilczynski SP; Oft M; Cook N; Liao SY; Iftner T
Human papillomavirus type 6 in squamous cell carcinoma of the bladder and cervix.
Hum Pathol 24: 96-102 (1993)
- Willey JC; Hei TK; Piao CQ; Madrid L; Willey JJ; Apostolakis MJ; Hukku B
Radiation-induced deletion of chromosomal regions containing tumor suppressor genes in human bronchial epithelial cells.
Carcinogenesis 14: 1181-8 (1993)
- Willis G; Jennings B; Ball RY; New NE; Gibson I
Analysis of ras point mutations and human papillomavirus 16 and 18 in cervical carcinomata and their metastases.
Gynecol Oncol 49: 359-64 (1993)
- Wisotzkey JD; Krizenoskas A; DiAngelo S; Kreider JW
Cleavage of cottontail rabbit papillomavirus E7 RNA with an anti-E7 ribozyme.
Biochem Biophys Res Commun 192: 833-9 (1993)
- Wong WS; Wong YF; Tam OS; Tam JS
Detection of human papilloma virus (HPV) infection in paraffin-embedded tissues of endometrial carcinoma.
Aust N Z J Obstet Gynaecol 33: 180-2 (1993)
- Woods KV; Shillitoe EJ; Spitz MR; Schantz SP; Adler-Storthz K
Analysis of human papillomavirus DNA in oral squamous cell carcinomas.
J Oral Pathol Med 22: 101-8 (1993)

References (1993)

- Woodworth CD; Wang H; Simpson S; Alvarez-Salas LM; Notario V
Overexpression of wild-type p53 alters growth and differentiation of normal human keratinocytes but not human papillomavirus-expressing cell lines.
Cell Growth Differ 4: 367-76 (1993)
- Wu EW; Clemens KE; Heck DV; Munger K
The human papillomavirus E7 oncoprotein and the cellular transcription factor E2F bind to separate sites on the retinoblastoma tumor suppressor protein.
J Virol 67: 2402-7 (1993)
- Xi LF; Demers W; Kiviat NB; Kuypers J; Beckmann AM; Galloway DA
Sequence variation in the noncoding region of human papillomavirus type 16 detected by single-strand conformation polymorphism analysis.
J Infect Dis 168: 610-7 (1993)
- Xu Q
Molecular biology of human papillomavirus.
Chin Med Sci J 8: 111-7 (1993)
- Xu YF; Meyer AN; Webster MK; Lee BA; Donoghue DJ
The v-sis protein retains biological activity as a type II membrane protein when anchored by various signal-anchor domains, including the hydrophobic domain of the bovine papilloma virus E5 oncoprotein.
J Cell Biol 123: 549-60 (1993)
- Yaegashi N; Xi L; Batra M; Galloway DA
Sequence and antigenic diversity in two immunodominant regions of the L2 protein of human papillomavirus types 6 and 16.
J Infect Dis 168: 743-7 (1993)
- Yamashita T; Segawa K; Fujinaga Y; Nishikawa T; Fujinaga K
Biological and biochemical activity of E7 genes of the cutaneous human papillomavirus type 5 and 8.
Oncogene 8: 2433-41 (1993)
- Yang L; Mohr I; Fouts E; Lim DA; Nohaile M; Botchan M
The E1 protein of bovine papilloma virus 1 is an ATP-dependent DNA helicase.
Proc Natl Acad Sci U S A 90: 5086-90 (1993)
- Yankaskas JR; Haizlip JE; Conrad M; Koval D; Lazarowski E; Paradiso AM; Rinehart CA Jr; Sarkadi B; Schlegel R; Boucher RC
Papilloma virus immortalized tracheal epithelial cells retain a well-differentiated phenotype.
Am J Physiol 264: C1219-30 (1993)
- Yu ST; Wu MM; Li LM
Prevalence of human papillomaviruses 16 and 18 in transitional cell carcinoma of bladder.
Chin Med J (Engl) 106: 494-6 (1993)
- Zhang W
[Cloning and characterization of the genome of human papilloma virus type 16 in China]
Chung Hua I Hsueh Tsa Chih 73: 229-31, 254-5 (1993)
- Zhang W; Liu Y; Jin Y; Wang X; Jin S; Liu F; Li J
Detection of the HPV16 E6 transforming gene by PCR in tissue samples from normal cervix and from cervix with precancerous lesions and carcinomas.
Chin Med Sci J 8: 139-42 (1993)
- Zhou J; Stenzel DJ; Sun XY; Frazer IH
Synthesis and assembly of infectious bovine papillomavirus particles in vitro.
J Gen Virol 74 (Pt 4): 763-8 (1993)
- Zhou J; Sun XY; Frazer IH
Glycosylation of human papillomavirus type 16 L1 protein.
Virology 194: 210-8 (1993)
- Zhu WY; Leonardi C; Blauvelt A; Serfling U; Penneys NS
Human papillomavirus DNA in the dermis of condyloma acuminatum.
J Cutan Pathol 20: 447-50 (1993)
- Zhu WY; Leonardi C; Penneys NS
Polymerase chain reaction in detection of human papillomavirus DNA and types of condyloma acuminata.
Chin Med J (Engl) 106: 141-4 (1993)

- Agarwal C; Hembree JR; Rorke EA; Eckert RL
Interferon and retinoic acid suppress the growth of human papillomavirus type 16 immortalized cervical epithelial cells, but only interferon suppresses the level of the human papillomavirus transforming oncogenes.
Cancer Res 54: 2108-12 (1994)
- Ansink AC; Krul MR; De Weger RA; Kleyne JA; Pijpers H; Van Tinteren H; De Kraker EW; Helmerhorst TJ; Heintz AP
Human papillomavirus, lichen sclerosus, and squamous cell carcinoma of the vulva: detection and prognostic significance.
Gynecol Oncol 52: 180-4 (1994)
- Baccard-Longere M; Alpha-Bazin B; Chypre C; Sauvaigo S; Teoule R; Bernard P; Seigneurin JM
Fast solid support detection of human papillomavirus in vitro amplified DNA using a DNP-anti DNP monoclonal antibody couple.
J Virol Methods 46: 29-38 (1994)
- Berchuck A; Kohler MF; Marks JR; Wiseman R; Boyd J; Bast RC Jr
The p53 tumor suppressor gene frequently is altered in gynecologic cancers.
Am J Obstet Gynecol 170: 246-52 (1994)
- Bernard HU; Apt D
Transcriptional control and cell type specificity of HPV gene expression.
Arch Dermatol 130: 210-5 (1994)
- Berumen J; Casas L; Segura E; Amezcua JL; Garcia-Carranca A
Genome amplification of human papillomavirus types 16 and 18 in cervical carcinomas is related to the retention of E1/E2 genes.
Int J Cancer 56: 640-5 (1994)
- Bristow RG; Jang A; Peacock J; Chung S; Benchimol S; Hill RP
Mutant p53 increases radioresistance in rat embryo fibroblasts simultaneously transfected with HPV16-E7 and/or activated H-ras.
Oncogene 9: 1527-36 (1994)
- Brown DR; Fan L; Jones J; Bryan J
Colocalization of human papillomavirus type 11 E1 [symbol: see text] E4 and L1 proteins in human foreskin implants grown in athymic mice.
Virology 201: 46-54 (1994)
- Burnett AF; Grendys EC; Willett GD; Johnson JC; Barter JF; Barnes WA
Preservation of multiple oncogenic human papillomavirus types in recurrences of early-stage cervical cancers.
Am J Obstet Gynecol 170: 1230-3 (1994)
- Busby-Earle RM; Steel CM; Williams AR; Cohen B; Bird CC
p53 mutations in cervical carcinogenesis—low frequency and lack of correlation with human papillomavirus status.
Br J Cancer 69: 732-7 (1994)
- Chambers MA; Wei Z; Coleman N; Nash AA; Stanley MA
“Natural” presentation of human papillomavirus type-16 E7 protein to immunocompetent mice results in antigen-specific sensitization or sustained unresponsiveness.
Eur J Immunol 24: 738-45 (1994)
- Chan PJ; Su BC; Kalugdan T; Seraj IM; Tredway DR; King A
Human papillomavirus gene sequences in washed human sperm deoxyribonucleic acid.
Fertil Steril 61: 982-5 (1994)
- Chan SY; Tan CH; Delius H; Bernard HU
Human papillomavirus type 2c is identical to human papillomavirus type 27.
Virology 201: 397-8 (1994)
- Chandrachud LM; O’Neil BW; Jarrett WF; Grindlay GJ; McGarvie GM; Campo MS
Humoral immune response to the E7 protein of bovine papillomavirus type 4 and identification of B-cell epitopes.
Virology 200: 98-104 (1994)
- Chen M; Popescu N; Woodworth C; Berneman Z; Corbellino M; Lusso P; Ablashi DV; DiPaolo JA
Human herpesvirus 6 infects cervical epithelial cells and transactivates human papillomavirus gene expression.
J Virol 68: 1173-8 (1994)
- Chen TM; Chen CA; Wu CC; Huang SC; Chang CF; Hsieh CY
The genotypes and prognostic significance of human papillomaviruses in cervical cancer.
Int J Cancer 57: 181-4 (1994)
- Chinami M; Sasaki S; Hachiya N; Yuge K; Ohsugi T; Maeda H; Shingu M
Functional oligomerization of purified human papillomavirus types 16 and 6b E7 proteins expressed in *Escherichia coli*.
J Gen Virol 75 (Pt 2): 277-81 (1994)

References (1994)

- Choo CK; Rorke EA; Eckert RL
Differentiation-independent constitutive expression of the human papillomavirus type 16 E6 and E7 oncogenes in the CaSki cervical tumour cell line.
J Gen Virol 75 (Pt 5): 1139-47 (1994)
- Ciechanover A; Shkedy D; Oren M; Bercovich B
Degradation of the tumor suppressor protein p53 by the ubiquitin-mediated proteolytic system requires a novel species of ubiquitin-carrier protein, E2.
J Biol Chem 269: 9582-9 (1994)
- Cleveland DB; Chen SY; Allen CM; Ahing SI; Svirsky JA
Adult rhabdomyoma. A light microscopic, ultrastructural, virologic, and immunologic analysis.
Oral Surg Oral Med Oral Pathol 77: 147-53 (1994)
- Crook T; Fisher C; Masterson PJ; Vousden KH
Modulation of transcriptional regulatory properties of p53 by HPV E6.
Oncogene 9: 1225-30 (1994)
- Cuzick J; Terry G; Ho L; Hollingworth T; Anderson M
Type-specific human papillomavirus DNA in abnormal smears as a predictor of high-grade cervical intraepithelial neoplasia.
Br J Cancer 69: 167-71 (1994)
- Czegledy J; Evander M; Hernadi Z; Gergely L; Wadell G
Human papillomavirus type 18 E6* mRNA in primary tumors and pelvic lymph nodes of Hungarian patients with squamous cervical cancer.
Int J Cancer 56: 182-6 (1994)
- Deb SP; Munoz RM; Brown DR; Subler MA; Deb S
Wild-type human p53 activates the human epidermal growth factor receptor promoter.
Oncogene 9: 1341-9 (1994)
- Demers GW; Foster SA; Halbert CL; Galloway DA
Growth arrest by induction of p53 in DNA damaged keratinocytes is bypassed by human papillomavirus 16 E7.
Proc Natl Acad Sci U S A 91: 4382-6 (1994)
- Demers GW; Halbert CL; Galloway DA
Elevated wild-type p53 protein levels in human epithelial cell lines immortalized by the human papillomavirus type 16 E7 gene.
Virology 198: 169-74 (1994)
- de Roda Husman AM; Walboomers JM; Meijer CJ; Risse EK; Schipper ME; Helmerhorst TM; Bleker OP; Delius H; van den Brule AJ; Snijders PJ
Analysis of cytomorphologically abnormal cervical scrapes for the presence of 27 mucosotropic human papillomavirus genotypes, using polymerase chain reaction.
Int J Cancer 56: 802-6 (1994)
- Dillner J
Enzyme immunoassay detection of induction of MHC class I expression by synthetic peptides from the E6 and E7 regions of human papillomavirus type 16.
J Immunol Methods 167: 195-205 (1994)
- Dillner J; Lenner P; Lehtinen M; Eklund C; Heino P; Wiklund F; Hallmans G; Stendahl U
A population-based seroepidemiological study of cervical cancer.
Cancer Res 54: 134-41 (1994)
- Dong G; Broker TR; Chow LT
Human papillomavirus type 11 E2 proteins repress the homologous E6 promoter by interfering with the binding of host transcription factors to adjacent elements.
J Virol 68: 1115-27 (1994)
- Eluf-Neto J; Booth M; Munoz N; Bosch FX; Meijer CJ; Walboomers JM
Human papillomavirus and invasive cervical cancer in Brazil.
Br J Cancer 69: 114-9 (1994)
- Fishman A; Girtanner RE; Adam E; Kaufman RH; Kaplan AL
Second primary cancer arising from the neovulva following reconstruction with myocutaneous flaps.
Gynecol Oncol 53: 128-30 (1994)
- Gallego MI; Zimonjic DB; Popescu NC; DiPaolo JA; Lazo PA
Integration site of human papillomavirus type-18 DNA in chromosome band 8q22.1 of C4-I cervical carcinoma: DNase I hypersensitivity and methylation of cellular flanking sequences.
Genes Chromosom Cancer 9: 28-32 (1994)
- Gao L; Chain B; Sinclair C; Crawford L; Zhou J; Morris J; Zhu X; Stauss H
Immune response to human papillomavirus type 16 E6 gene in a live vaccinia vector.
J Gen Virol 75 (Pt 1): 157-64 (1994)
- Gregoire L; Lawrence WD; Kukuruga D; Eisenbrey AB; Lancaster WD
Association between HLA-DQB1 alleles and risk for cervical cancer in African-American women.
Int J Cancer 57: 504-7 (1994)

- Griep AE; Lambert PF
Role of papillomavirus oncogenes in human cervical cancer: transgenic animal studies.
Proc Soc Exp Biol Med 206: 24-34 (1994)
- Gu Z; Pim D; Labrecque S; Banks L; Matlashewski G
DNA damage induced p53 mediated transcription is inhibited by human papillomavirus type 18 E6.
Oncogene 9: 629-33 (1994)
- Ham J; Steger G; Yaniv M
Cooperativity in vivo between the E2 transactivator and the TATA box binding protein depends on core promoter structure.
EMBO J 13: 147-57 (1994)
- Hei TK; Piao CQ; Willey JC; Thomas S; Hall EJ
Malignant transformation of human bronchial epithelial cells by radon-simulated alpha-particles.
Carcinogenesis 15: 431-7 (1994)
- Hermonat PL
Adeno-associated virus inhibits human papillomavirus type 16: a viral interaction implicated in cervical cancer.
Cancer Res 54: 2278-81 (1994)
- Hildesheim A; Schiffman MH; Gravitt PE; Glass AG; Greer CE; Zhang T; Scott DR; Rush BB; Lawler P; Sherman ME; et al
Persistence of type-specific human papillomavirus infection among cytologically normal women.
J Infect Dis 169: 235-40 (1994)
- Hodivala KJ; Pei XF; Liu QY; Jones PH; Rytina ER; Gilbert C; Singer A; Watt FM
Integrin expression and function in HPV 16-immortalised human keratinocytes in the presence or absence of v-Ha-ras. Comparison with cervical intraepithelial neoplasia.
Oncogene 9: 943-8 (1994)
- Holt SE; Schuller G; Wilson VG
DNA binding specificity of the bovine papillomavirus E1 protein is determined by sequences contained within an 18-base-pair inverted repeat element at the origin of replication.
J Virol 68: 1094-102 (1994)
- Kahn T; Turazza E; Ojeda R; Bercovich A; Stremlau A; Lichter P; Poustka A; Grinstein S; zur Hausen H
Integration of human papillomavirus type 6a DNA in a tonsillar carcinoma: chromosomal localization and nucleotide sequence of the genomic target region.
Cancer Res 54: 1305-12 (1994)
- Kashanchi F; Thompson J; Sadaie MR; Doniger J; Duvall J; Brady JN; Rosenthal LJ
Transcriptional activation of minimal HIV-1 promoter by ORF-1 protein expressed from the SalI-L fragment of human herpesvirus 6.
Virology 201: 95-106 (1994)
- REFERENCE Kast WM; Brandt RM; Sidney J; Drijfhout JW; Kubo RT; Grey HM; Melief CJ; Sette A
Role of HLA-A motifs in identification of potential CTL epitopes in human papillomavirus type 16 E6 and E7 proteins.
J Immunol 152: 3904-12 (1994)
- Kitasato H; Delius H; zur Hausen H; Sorger K; Rosl F; de Villiers EM
Sequence rearrangements in the upstream regulatory region of human papillomavirus type 6: are these involved in malignant transition?
J Gen Virol 75 (Pt 5): 1157-62 (1994)
- Klingelhutz AJ; Barber SA; Smith PP; Dyer K; McDougall JK
Restoration of telomeres in human papillomavirus-immortalized human anogenital epithelial cells.
Mol Cell Biol 14: 961-9 (1994)
- Kristiansen E; Jenkins A; Holm R
Coexistence of episomal and integrated HPV16 DNA in squamous cell carcinoma of the cervix.
J Clin Pathol 47: 253-6 (1994)
- Kristiansen E; Jenkins A; Kristensen G; Ask E; Kaern J; Abeler V; Lindqvist BH; Trope C; Kristiansen BE
Human papillomavirus infection in Norwegian women with cervical cancer.
APMIS 102: 122-8 (1994)
- Kurvinen K; Tervahauta A; Syrjanen S; Chang F; Syrjanen K
The state of the p53 gene in human papillomavirus (HPV)-positive and HPV-negative genital precancer lesions and carcinomas as determined by single-strand conformation polymorphism analysis and sequencing.
Anticancer Res 14: 177-81 (1994)
- Kyo S; Inoue M; Hayasaka N; Inoue T; Yutsudo M; Tanizawa O; Hakura A
Regulation of early gene expression of human papillomavirus type 16 by inflammatory cytokines.
Virology 200: 130-9 (1994)
- Lam EW; Morris JD; Davies R; Crook T; Watson RJ; Vousden KH
HPV16 E7 oncoprotein deregulates B-myb expression: correlation with targeting of p107/E2F complexes.
EMBO J 13: 871-8 (1994)

References (1994)

- Lappalainen K; Urtti A; Jaaskelainen I; Syrjanen K; Syrjanen S
Cationic liposomes mediated delivery of antisense oligonucleotides targeted to HPV 16 E7 mRNA in CaSki cells.
Antiviral Res 23: 119-30 (1994)
- Larsen ET; Duggan MA; Inoue M
Absence of human papilloma virus DNA in oropharyngeal spindle-cell squamous carcinomas.
Am J Clin Pathol 101: 514-8 (1994)
- Le Moal MA; Yaniv M; Thierry F
The bovine papillomavirus type 1 (BPV1) replication protein E1 modulates transcriptional activation by interacting with BPV1 E2.
J Virol 68: 1085-93 (1994)
- Lee YY; Wilczynski SP; Chumakov A; Chih D; Koeffler HP
Carcinoma of the vulva: HPV and p53 mutations.
Oncogene 9: 1655-9 (1994)
- List HJ; Patzel V; Zeidler U; Schopen A; Ruhl G; Stollwerk J; Klock G
Methylation sensitivity of the enhancer from the human papillomavirus type 16.
J Biol Chem 269: 11902-11 (1994)
- Liu Z; Ghai J; Ostrow RS; McGlennen RC; Faras AJ
The E6 gene of human papillomavirus type 16 is sufficient for transformation of baby rat kidney cells in cotransfection with activated Ha-ras.
Virology 201: 388-96 (1994)
- Liu Z; Ghai J; Ostrow RS; McGlennen RC; Faras AJ
The E6 gene of human papillomavirus type 16 is sufficient for transformation of baby rat kidney cells in cotransfection with activated Ha-ras.
Virology 201: 388-96 (1994)
- Lukas J; Muller H; Bartkova J; Spitkovsky D; Kjerulff AA; Jansen-Durr P; Strauss M; Bartek J
DNA tumor virus oncoproteins and retinoblastoma gene mutations share the ability to relieve the cell's requirement for cyclin D1 function in G1.
J Cell Biol 125: 625-38 (1994)
- Mackenzie SL; Gillespie MT; Scurry JP; Planner RS; Martin TJ; Danks JA
Parathyroid hormone-related protein and human papillomavirus in gynecological tumors.
Int J Cancer 56: 324-30 (1994)
- Majewski S; Szmurlo A; Marczak M; Jablonska S; Bollag W
Synergistic effect of retinoids and interferon alpha on tumor-induced angiogenesis: anti-angiogenic effect on HPV-harboring tumor-cell lines.
Int J Cancer 57: 81-5 (1994)
- Malejczyk J; Malejczyk M; Majewski S; Breitburd F; Luger TA; Jablonska S; Orth G
Increased tumorigenicity of human keratinocytes harboring human papillomavirus type 16 is associated with resistance to endogenous tumor necrosis factor-alpha-mediated growth limitation.
Int J Cancer 56: 593-8 (1994)
- Maloney KE; Wiener JS; Walther PJ
Oncogenic human papillomaviruses are rarely associated with squamous cell carcinoma of the bladder: evaluation by differential polymerase chain reaction.
J Urol 151: 360-4 (1994)
- May M; Dong XP; Beyer-Finkler E; Stubenrauch F; Fuchs PG; Pfister H
The E6/E7 promoter of extrachromosomal HPV16 DNA in cervical cancers escapes from cellular repression by mutation of target sequences for YY1.
EMBO J 13: 1460-6 (1994)
- May M; Grassmann K; Pfister H; Fuchs PG
Transcriptional silencer of the human papillomavirus type 8 late promoter interacts alternatively with the viral trans activator E2 or with a cellular factor.
J Virol 68: 3612-9 (1994)
- McGregor JM; Farthing A; Crook T; Yu CC; Dublin EA; Levison DA; MacDonald DM
Posttransplant skin cancer: a possible role for p53 gene mutation but not for oncogenic human papillomaviruses.
J Am Acad Dermatol 30: 701-6 (1994)
- McLachlin CM; Tate JE; Zitz JC; Sheets EE; Crum CP
Human papillomavirus type 18 and intraepithelial lesions of the cervix.
Am J Pathol 144: 141-7 (1994)
- Melchers W; Ferrera A; Willemse D; Galama J; Walboomers J; De Barahona O; Figueroa M; Snijders P
Human papillomavirus and cervical cancer in Honduran women.
Am J Trop Med Hyg 50: 137-42 (1994)
- Morelli AE; Belardi G; DiPaola G; Paredes A; Fainboim L
Cellular subsets and epithelial ICAM-1 and HLA-DR expression in human papillomavirus infection of the vulva.
Acta Derm Venereol (Stockh) 74: 45-50 (1994)

- Muller H; Lukas J; Schneider A; Warthoe P; Bartek J; Eilers M; Strauss M
Cyclin D1 expression is regulated by the retinoblastoma protein.
Proc Natl Acad Sci U S A 91: 2945-9 (1994)
- Mvula M; Iwasaka T; Matsuo N; Nakao Y; Kaku T; Hachisuga T; Fukuda K; Tsukamoto N; Sugimori H
Detection of human papillomavirus types 16 and 18 in primary and in metastatic lesions of cervical carcinomas.
Gynecol Oncol 53: 156-60 (1994)
- Nagai N
[Gene diagnosis of uterine cervical cancer]
Rinsho Byori 42: 372-9 (1994)
- Nallaseth FS; DePamphilis ML
Papillomavirus contains cis-acting sequences that can suppress but not regulate origins of DNA replication.
J Virol 68: 3051-64 (1994)
- Ono H; Takahashi A; Ogoshi S; Furihata M; Ohtsuki Y
Relationship between H-ras p21 product and p53 protein or high-risk human papillomaviruses in esophageal cancer from Kochi, Japan [letter]
Am J Gastroenterol 89: 646-7 (1994)
- Park DJ; Wilczynski SP; Paquette RL; Miller CW; Koeffler HP
p53 mutations in HPV-negative cervical carcinoma.
Oncogene 9: 205-10 (1994)
- Patrick DR; Oliff A; Heimbrook DC
Identification of a novel retinoblastoma gene product binding site on human papillomavirus type 16 E7 protein.
J Biol Chem 269: 6842-50 (1994)
- Peacock JW; Benchimol S
Mutation of the endogenous p53 gene in cells transformed by HPV-16 E7 and EJ c-ras confers a growth advantage involving an autocrine mechanism.
EMBO J 13: 1084-92 (1994)
- Petricoin E 3rd; David M; Fang H; Grimley P; Larner AC; Vande Pol S
Human cancer cell lines express a negative transcriptional regulator of the interferon regulatory factor family of DNA binding proteins.
Mol Cell Biol 14: 1477-86 (1994)
- Petti L; DiMaio D
Specific interaction between the bovine papillomavirus E5 transforming protein and the beta receptor for platelet-derived growth factor in stably transformed and acutely transfected cells.
J Virol 68: 3582-92 (1994)
- Pushko P; Sasagawa T; Cuzick J; Crawford L
Sequence variation in the capsid protein genes of human papillomavirus type 16.
J Gen Virol 75 (Pt 4): 911-6 (1994)
- Rainey WH; Sawetawan C; Shay JW; Michael MD; Mathis JM; Kutteh W; Byrd W; Carr BR
Transformation of human granulosa cells with the E6 and E7 regions of human papillomavirus.
J Clin Endocrinol Metab 78: 705-10 (1994)
- Rajpert-De Meyts E; Hording U; Nielsen HW; Skakkebaek NE
Human papillomavirus and Epstein-Barr virus in the etiology of testicular germ cell tumours.
APMIS 102: 38-42 (1994)
- Ratiarson A
An anti-oncogenic action of the polyomavirus large-T (LT) antigen in HeLa and SiHa tumor cells.
Cancer Lett 78: 183-7 (1994)
- Rose BR; Thompson CH; Jiang XM; Tattersall MH; Elliott PM; Dalrymple C; Cossart YE
Detection of human papillomavirus type 16 E6/E7 transcripts in histologically cancer-free pelvic lymph nodes of patients with cervical carcinoma.
Gynecol Oncol 52: 212-7 (1994)
- Rosl F; Lengert M; Albrecht J; Kleine K; Zawatzky R; Schraven B; zur Hausen H
Differential regulation of the JE gene encoding the monocyte chemoattractant protein (MCP-1) in cervical carcinoma cells and derived hybrids.
J Virol 68: 2142-50 (1994)
- Rusconi S; Agarossi A; Ravasi L; Tonta A; Conti M; Galli M
Serum 2'-5'-oligoadenylate synthetase levels and clinical response to interferon-beta therapy in women with genital human papillomavirus infection.
J Infect Dis 169: 1112-5 (1994)
- Ryan MJ; Johnson G; Kirk J; Fuerstenberg SM; Zager RA; Torok-Storb B
HK-2: an immortalized proximal tubule epithelial cell line from normal adult human kidney.
Kidney Int 45: 48-57 (1994)
- Sadovnikova E; Zhu X; Collins SM; Zhou J; Vousden K; Crawford L; Beverley P; Stauss HJ
Limitations of predictive motifs revealed by cytotoxic T lymphocyte epitope mapping of the human papilloma virus E7 protein.
Int Immunol 6: 289-96 (1994)

References (1994)

- Schuller G; Holt SE; Hsu J; Wilson VG
The bovine papillomavirus type 1 genome contains multiple loci of static DNA bending, but bends are absent from the functional origin of replication.
Virus Res 31: 203-17 (1994)
- Searle PF; Thomas DP; Faulkner KB; Tinsley JM
Stomach cancer in transgenic mice expressing human papillomavirus type 16 early region genes from a keratin promoter.
J Gen Virol 75 (Pt 5): 1125-37 (1994)
- Selvakumar R; Borenstein LA; Lin YL; Ahmed R; Wettstein FO
T-cell response to cottontail rabbit papillomavirus structural proteins in infected rabbits.
J Virol 68: 4043-8 (1994)
- Selvakumar R; Borenstein LA; Lin YL; Ahmed R; Wettstein FO
T-cell response to cottontail rabbit papillomavirus structural proteins in infected rabbits.
J Virol 68: 4043-8 (1994)
- Shamanin V; Delius H; de Villiers EM
Development of a broad spectrum PCR assay for papillomaviruses and its application in screening lung cancer biopsies.
J Gen Virol 75 (Pt 5): 1149-56 (1994)
- Shin KH; Min BM; Cherrick HM; Park NH
Combined effects of human papillomavirus-18 and N-methyl-N'-nitro-N-nitrosoguanidine on the transformation of normal human oral keratinocytes.
Mol Carcinog 9: 76-86 (1994)
- Smith-McCune KK; Weidner N
Demonstration and characterization of the angiogenic properties of cervical dysplasia.
Cancer Res 54: 800-4 (1994)
- Stacey SN; Eklund C; Jordan D; Smith NK; Stern PL; Dillner J; Arrand JR
Scanning the structure and antigenicity of HPV-16 E6 and E7 oncoproteins using anti-peptide antibodies.
Oncogene 9: 635-45 (1994)
- Stark LA; Arends MJ; McLaren KM; Benton EC; Shahidullah H; Hunter JA; Bird CC
Prevalence of human papillomavirus DNA in cutaneous neoplasms from renal allograft recipients supports a possible viral role in tumour promotion.
Br J Cancer 69: 222-9 (1994)
- Steinmann KE; Pei XF; Stoppler H; Schlegel R; Schlegel R
Elevated expression and activity of mitotic regulatory proteins in human papillomavirus-immortalized keratinocytes.
Oncogene 9: 387-94 (1994)
- Stewart AC; Jareborg N; Simonsson M; Alderborn A; Burnett S
Segregation properties of bovine papillomaviral plasmid DNA.
J Mol Biol 236: 480-90 (1994)
- Suzuki H; Sato N; Kodama T; Okano T; Isaka S; Shirasawa H; Simizu B; Shimazaki J
Detection of human papillomavirus DNA and state of p53 gene in Japanese penile cancer.
Jpn J Clin Oncol 24: 1-6 (1994)
- Sverdrup F; Khan SA
Replication of human papillomavirus (HPV) DNAs supported by the HPV type 18 E1 and E2 proteins.
J Virol 68: 505-9 (1994)
- Tachezy R; Hamsikova E; Valvoda J; Van Ranst M; Betka J; Burk RD; Vonka V
Antibody response to a synthetic peptide derived from the human papillomavirus type 6/11 L2 protein in recurrent respiratory papillomatosis: correlation between Southern blot hybridization, polymerase chain reaction, and serology.
J Med Virol 42: 52-9 (1994)
- Tait DL; MacDonald PC; Casey ML
Parathyroid hormone-related protein expression in gynecic squamous carcinoma cells.
Cancer 73: 1515-21 (1994)
- Tan CH; Tachezy R; Van Ranst M; Chan SY; Bernard HU; Burk RD
The *Mastomys natalensis* papillomavirus: nucleotide sequence, genome organization, and phylogenetic relationship of a rodent papillomavirus involved in tumorigenesis of cutaneous epithelia.
Virology 198: 534-41 (1994)
- Tarpey I; Stacey S; Hickling J; Birley HD; Renton A; McIndoe A; Davies DH
Human cytotoxic T lymphocytes stimulated by endogenously processed human papillomavirus type 11 E7 recognize a peptide containing a HLA-A2 (A*0201) motif.
Immunology 81: 222-7 (1994)
- Tate JE; Mutter GL; Prasad CJ; Berkowitz R; Goodman H; Crum CP
Analysis of HPV-positive and -negative vulvar carcinomas for alterations in c-myc, Ha-, Ki-, and N-ras genes.
Gynecol Oncol 53: 78-83 (1994)

- Tindle RW; Herd K; Londono P; Fernando GJ; Chatfield SN; Malcolm K; Dougan G
Chimeric hepatitis B core antigen particles containing B- and Th-epitopes of human papillomavirus type 16 E7 protein induce specific antibody and T-helper responses in immunised mice.
Virology 200: 547-57 (1994)
- Viallet J; Liu C; Emond J; Tsao MS
Characterization of human bronchial epithelial cells immortalized by the E6 and E7 genes of human papillomavirus type 16.
Exp Cell Res 212: 36-41 (1994)
- Volpers C; Schirmacher P; Streeck RE; Sapp M
Assembly of the major and the minor capsid protein of human papillomavirus type 33 into virus-like particles and tubular structures in insect cells.
Virology 200: 504-12 (1994)
- von Knebel Doeberitz M; Rittmuller C; Aengeneyndt F; Jansen-Durr P; Spitkovsky D
Reversible repression of papillomavirus oncogene expression in cervical carcinoma cells: consequences for the phenotype and E6-p53 and E7-pRB interactions.
J Virol 68: 2811-21 (1994)
- Voog E; Ricksten A; Lowhagen GB; Ternesten A
Demonstration of Epstein-Barr virus DNA in acetowhite lesions of the vulva.
Int J STD AIDS 5: 25-8 (1994)
- Yokoyama M; Tsutsumi K; Pater A; Pater MM
Human papillomavirus 18-immortalized endocervical cells with in vitro cytokeratin expression characteristics of adenocarcinoma.
Obstet Gynecol 83: 197-204 (1994)
- Zeltner R; Borenstein LA; Wettstein FO; Iftner T
Changes in RNA expression pattern during the malignant progression of cottontail rabbit papillomavirus-induced tumors in rabbits.
J Virol 68: 3620-30 (1994)
- Zhang Q; Gutsch D; Kenney S
Functional and physical interaction between p53 and BZLF1: implications for Epstein-Barr virus latency.
Mol Cell Biol 14: 1929-38 (1994)
- Zhou J; Sun XY; Louis K; Frazer IH
Interaction of human papillomavirus (HPV) type 16 capsid proteins with HPV DNA requires an intact L2 N-terminal sequence.
J Virol 68: 619-25 (1994)
- Zimonjic DB; Popescu ND; DiPaolo JA
Chromosomal organization of viral integration sites in human papillomavirus-immortalized human keratinocyte cell lines.
Cancer Genet Cytogenet 72: 39-43 (1994)
- zur Hausen H
Disrupted dichotomous intracellular control of human papillomavirus infection in cancer of the cervix.
Lancet 343: 955-7 (1994)