

- Adams PD; Kaelin WG Jr
Transcriptional control by E2F.
Semin Cancer Biol 6: 99-108 (1995)
- Adams V; Kempf W; Hassam S; Briner J; Schmid M; Moos R; Pfaltz M
Detection of several types of human papilloma viruses in AIDS-associated Kaposi's sarcoma.
J Med Virol 46: 189-93 (1995)
- Agorastos T; Bontis J; Lambropoulos AF; Constantinidis TC; Natsioutziki M; Tagou C; Katsouyiannopoulos V
Epidemiology of human papillomavirus infection in Greek asymptomatic women.
Eur J Cancer Prev 4: 159-67 (1995)
- Akasofu M; Oda Y
Immunohistochemical detection of p53 in cervical epithelial lesions with or without infection of human papillomavirus types 16 and 18.
Virchows Arch 425: 593-602 (1995)
- Akutsu N; Shirasawa H; Nakano K; Tanzawa H; Asano T; Kobayashi S; Isono K; Simizu B
Rare association of human papillomavirus DNA with esophageal cancer in Japan.
J Infect Dis 171: 425-8 (1995)
- Almadori G; Cadoni G; Maurizi M; Ottaviani F; Paludetti G; Cattani P; Scambia G
[Oncogenes and cancer of the larynx. EGFR, p21 ras and HPV-DNA infections]
Acta Otorhinolaryngol Ital 15: 1-22 (1995)
- Alvarez-Salas LM; Lopez-Bayghen E
[Genetic regulation of human genital papillomaviruses]
Salud Publica Mex 37: 241-7 (1995)
- Alvarez-Salas LM; Velazquez A; Lopez-Bayghen E; Woodworth CD; Garrido E; Gariglio P; DiPaolo JA
Transcriptional repression in normal human keratinocytes by wild-type and mutant p53.
Cancer Lett 91: 85-92 (1995)
- al-Ghamdi AA; Sanders CM; Keefe M; Coggon D; Maitland NJ
Human papillomavirus DNA and TP53 mutations in lung cancers from butchers.
Br J Cancer 72: 293-7 (1995)
- Anand P; Foley P; Navsaria HA; Sinicropi D; Williams-Chestnut RE; Leigh IM
Nerve growth factor levels in cultured human skin cells: effect of gestation and viral transformation.
Neurosci Lett 184: 157-60 (1995)
- Andresson T; Sparkowski J; Goldstein DJ; Schlegel R
Vacuolar H(+)-ATPase mutants transform cells and define a binding site for the papillomavirus E5 oncoprotein.
J Biol Chem 270: 6830-7 (1995)
- Arany I; Nagamani K; Tyring SK
Interferon resistance is independent from copy numbers in benign HPV-induced lesions.
Anticancer Res 15: 1003-6 (1995)
- Arends MJ; Wyllie AH; Bird CC
Human papillomavirus type 18 is associated with less apoptosis in fibroblast tumours than human papillomavirus type 16.
Br J Cancer 72: 646-9 (1995)
- Armstrong DJ; Roman A
Human papillomavirus type 6 E7 protein is a substrate in vitro of protein kinase C.
Biochem J 312 (Pt 3): 667-70 (1995)
- Baay MF; Duk JM; Burger MP; Walboomers J; ter Schegget J; Groenier KH; de Bruijn HW; Stolz E; Herbrink P
Antibodies to human papillomavirus type 16 E7 related to clinicopathological data in patients with cervical carcinoma.
J Clin Pathol 48: 410-4 (1995)
- Baay MF; Duk JM; Burger MP; de Bruijn HW; Stolz E; Herbrink P
Follow-up of antibody responses to human papillomavirus type 16 E7 in patients treated for cervical carcinoma.
J Med Virol 45: 342-7 (1995)
- Balaram P; Nalinakumari KR; Abraham E; Balan A; Hareendran NK; Bernard HU; Chan SY
Human papillomaviruses in 91 oral cancers from Indian betel quid chewers—high prevalence and multiplicity of infections.
Int J Cancer 61: 450-4 (1995)
- Barksdale S; Baker CC
Differentiation-specific alternative splicing of bovine papillomavirus late mRNAs.
J Virol 69: 6553-6 (1995)
- Barksdale SK; Baker CC
The human immunodeficiency virus type 1 Rev protein and the Rev-responsive element counteract the effect of an inhibitory 5' splice site in a 3' untranslated region.
Mol Cell Biol 15: 2962-71 (1995)
- Barten M; Ostwald C; Milde-Langosch K; Muller P; Wukasch Y; Loning T
HPV DNA and p53 alterations in oropharyngeal carcinomas.
Virchows Arch 427: 153-7 (1995)

References (1995)

- Bauer S; Heeg K; Wagner H; Lipford GB
Identification of H-2Kb binding and immunogenic peptides from human papilloma virus tumour antigens E6 and E7.
Scand J Immunol 42: 317-23 (1995)
- Bauknecht T; Jundt F; Herr I; Oehler T; Delius H; Shi Y; Angel P; Zur Hausen H
A switch region determines the cell type-specific positive or negative action of YY1 on the activity of the human papillomavirus type 18 promoter.
J Virol 69: 1-12 (1995)
- Belaguli NS; Pater MM; Pater A
Splice sites of human papillomavirus type 16 E6 gene or heterologous gene required for transformation by E7 and accumulation of E7 RNA.
J Med Virol 47: 445-53 (1995)
- Belyavskiy M; Miller J; Belyavskaya E; Wilson V
BPV E1 protein alters the kinetics of cell cycle entry of serum starved mouse fibroblasts.
Cytometry 21: 257-64 (1995)
- Benamouzig R; Jullian E; Chang F; Robaskiewicz M; Flejou JF; Raoul JL; Coste T; Couturier D; Pompidou A; Rautureau J
Absence of human papillomavirus DNA detected by polymerase chain reaction in French patients with esophageal carcinoma.
Gastroenterology 109: 1876-81 (1995)
- Benson JD; Howley PM
Amino-terminal domains of the bovine papillomavirus type 1 E1 and E2 proteins participate in complex formation.
J Virol 69: 4364-72 (1995)
- Berkhout RJ; Tieben LM; Smits HL; Bavinck JN; Vermeer BJ; ter Schegget J
Nested PCR approach for detection and typing of epidermodysplasia verruciformis-associated human papillomavirus types in cutaneous cancers from renal transplant recipients.
J Clin Microbiol 33: 690-5 (1995)
- Berumen J; Unger ER; Casas L; Figueroa P
Amplification of human papillomavirus types 16 and 18 in invasive cervical cancer.
Hum Pathol 26: 676-81 (1995)
- Bethwaite PB; Koreth J; Herrington CS; McGee JO
Loss of heterozygosity occurs at the D11S29 locus on chromosome 11q23 in invasive cervical carcinoma.
Br J Cancer 71: 814-8 (1995)
- Bochkarev A; Barwell JA; Pfuetzner RA; Furey W Jr; Edwards AM; Frappier L
Crystal structure of the DNA-binding domain of the Epstein-Barr virus origin-binding protein EBNA 1.
Cell 83: 39-46 (1995)
- Bonne-Andrea C; Santucci S; Clertant P
Bovine papillomavirus E1 protein can, by itself, efficiently drive multiple rounds of DNA synthesis in vitro.
J Virol 69: 3201-5 (1995)
- Bonne-Andrea C; Santucci S; Clertant P; Tillier F
Bovine papillomavirus E1 protein binds specifically DNA polymerase alpha but not replication protein A.
J Virol 69: 2341-50 (1995)
- Brune W; Durst M
Epithelial differentiation fails to support replication of cloned human papillomavirus type 16 DNA in transfected keratinocytes.
J Invest Dermatol 104: 277-81 (1995)
- Butz K; Hoppe-Seyley F
[Viruses and cancer: molecular pathologic mechanisms of viral carcinogenesis]
Immun Infekt 23: 179-84 (1995)
- Butz K; Shahabeddin L; Geisen C; Spitkovsky D; Ullmann A; Hoppe-Seyley F
Functional p53 protein in human papillomavirus-positive cancer cells.
Oncogene 10: 927-36 (1995)
- Caballero OL; Villa LL; Simpson AJ
Low stringency-PCR (LS-PCR) allows entirely internally standardized DNA quantitation.
Nucleic Acids Res 23: 192-3 (1995)
- Cairney M; Campo MS
The synergism between bovine papillomavirus type 4 and quercetin is dependent on the timing of exposure.
Carcinogenesis 16: 1997-2001 (1995)
- Campo MS
Infection by bovine papillomavirus and prospects for vaccination.
Trends Microbiol 3: 92-7 (1995)
- Campos-Banales ME; Perez-Pinero B; Barrios del Pino Y; Salido Ruiz E; Rivero Suarez J; Ruiz Casals E; Lopez Aguado D
[DNA papillomavirus detection in head and neck benign proliferative lesions]
An Otorrinolaringol Ibero Am 22: 487-94 (1995)

- Canman CE; Gilmer TM; Coutts SB; Kastan MB
Growth factor modulation of p53-mediated growth arrest versus apoptosis.
Genes Dev 9: 600-11 (1995)
- Carter JJ; Wipf GC; Hagensee ME; McKnight B; Habel LA; Lee SK; Kuypers J; Kiviat N; Daling JR; Koutsky LA; et al
Use of human papillomavirus type 6 capsids to detect antibodies in people with genital warts.
J Infect Dis 172: 11-8 (1995)
- Chan PJ; Seraj IM; Kalugdan TH; King A
Blastocysts exhibit preferential uptake of DNA fragments from the E6-E7 conserved region of the human papillomavirus.
Gynecol Oncol 58: 194-7 (1995)
- Chan SY; Delius H; Halpern AL; Bernard HU
Analysis of genomic sequences of 95 papillomavirus types: uniting typing, phylogeny, and taxonomy.
J Virol 69: 3074-83 (1995)
- Chandrachud LM; Grindlay GJ; McGarvie GM; O'Neil BW; Wagner ER; Jarrett WF; Campo MS
Vaccination of cattle with the N-terminus of L2 is necessary and sufficient for preventing infection by bovine papillomavirus-4.
Virology 211: 204-8 (1995)
- Chang DY; Hsieh CY; Chen RJ; Lee SC; Huang SC
Comparison of detection of human papillomavirus 16 DNA in cervical carcinoma tissues by Southern blot hybridisation and nested polymerase chain reaction.
J Med Microbiol 43: 430-5 (1995)
- Chee YH; Namkoong SE; Kim DH; Kim SJ; Park JS
Immunologic diagnosis and monitoring of cervical cancers using in vitro translated HPV proteins.
Gynecol Oncol 57: 226-31 (1995)
- Chen CM; Choo KB; Cheng WT
Frequent deletions and sequence aberrations at the transgene junctions of transgenic mice carrying the papillomavirus regulatory and the SV40 TAg gene sequences.
Transgenic Res 4: 52-9 (1995)
- Chen JJ; Reid CE; Band V; Androphy EJ
Interaction of papillomavirus E6 oncoproteins with a putative calcium-binding protein.
Science 269: 529-31 (1995)
- Chen SL; Tsao YP; Yang CM; Lin YK; Huang CH; Kuo SW
Differential induction and regulation of c-jun, junB, junD and c-fos by human papillomavirus type 11 E5a oncoprotein.
J Gen Virol 76 (Pt 11): 2653-9 (1995)
- Chen Z; Kamath P; Zhang S; Weil MM; Shillitoe EJ
Effectiveness of three ribozymes for cleavage of an RNA transcript from human papillomavirus type 18.
Cancer Gene Ther 2: 263-71 (1995)
- Cheng JY; Sheu LF; Lin JC; Meng CL
Detection of human papillomavirus DNA in colorectal adenomas.
Arch Surg 130: 73-6 (1995)
- Cheng JY; Sheu LF; Meng CL; Lee WH; Lin JC
Detection of human papillomavirus DNA in colorectal carcinomas by polymerase chain reaction.
Gut 37: 87-90 (1995)
- Cheng S; Schmidt-Grimminger DC; Murant T; Broker TR; Chow LT
Differentiation-dependent up-regulation of the human papillomavirus E7 gene reactivates cellular DNA replication in suprabasal differentiated keratinocytes.
Genes Dev 9: 2335-49 (1995)
- Choo CK; Rorke EA; Eckert RL
Retinoid regulation of cell differentiation in a series of human papillomavirus type 16-immortalized human cervical epithelial cell lines.
Carcinogenesis 16: 375-81 (1995)
- Choo KB; Huang CJ; Chen CM; Han CP; Au LC
Jun-B oncogene aberrations in cervical cancer cell lines.
Cancer Lett 93: 249-53 (1995)
- Chou CY; Shen MR; Wu SN
Volume-sensitive chloride channels associated with human cervical carcinogenesis.
Cancer Res 55: 6077-83 (1995)
- Clemens KE; Brent R; Gyuris J; Munger K
Dimerization of the human papillomavirus E7 oncoprotein in vivo.
Virology 214: 289-93 (1995)
- Comerford SA; Maika SD; Laimins LA; Messing A; Elsasser HP; Hammer RE
E6 and E7 expression from the HPV 18 LCR: development of genital hyperplasia and neoplasia in transgenic mice.
Oncogene 10: 587-97 (1995)
- Conroy SC; Hart CE; Perez-Reyes N; Giachelli CM; Schwartz SM; McDougall JK
Characterization of human aortic smooth muscle cells expressing HPV16 E6 and E7 open reading frames.
Am J Pathol 147: 753-62 (1995)

References (1995)

- Cooper K
p53 mutations in human papillomavirus-associated oesophageal squamous cell carcinoma [letter]
Br J Cancer 72: 1337 (1995)
- Cooper K; Taylor L; Govind S
Human papillomavirus DNA in oesophageal carcinomas in South Africa.
J Pathol 175: 273-7 (1995)
- Coutlee F; Provencher D; Voyer H
Detection of human papillomavirus DNA in cervical lavage specimens by a nonisotopic consensus PCR assay.
J Clin Microbiol 33: 1973-8 (1995)
- Craig J; Hopkins M; DeLucia A
Uterine cervix adenocarcinoma with both human papillomavirus type 18 and tumor suppressor gene p53 mutation from a woman having an intact hymen.
Gynecol Oncol 59: 423-6 (1995)
- Creek KE; Geslani G; Batova A; Pirisi L
Progressive loss of sensitivity to growth control by retinoic acid and transforming growth factor-beta at late stages of human papillomavirus type 16-initiated transformation of human keratinocytes.
Adv Exp Med Biol 375: 117-35 (1995)
- Crooke RM; Graham MJ; Cooke ME; Crooke ST
In vitro pharmacokinetics of phosphorothioate antisense oligonucleotides.
J Pharmacol Exp Ther 275: 462-73 (1995)
- Cupp MR; Malek RS; Goellner JR; Smith TF; Espy MJ
The detection of human papillomavirus deoxyribonucleic acid in intraepithelial, in situ, verrucous and invasive carcinoma of the penis.
J Urol 154: 1024-9 (1995)
- Czegledy J; Iosif C; Hansson BG; Evander M; Gergely L; Wadell G
Can a test for E6/E7 transcripts of human papillomavirus type 16 serve as a diagnostic tool for the detection of micrometastasis in cervical cancer?
Int J Cancer 64: 211-5 (1995)
- D'Amato J; Houbiers JG; Drijfhout JW; Brandt RM; Schipper R; Bavinck JN; Melief CJ; Kast WM
A computer program for predicting possible cytotoxic T lymphocyte epitopes based on HLA class I peptide-binding motifs.
Hum Immunol 43: 13-8 (1995)
- Daniel B; Mukherjee G; Seshadri L; Vallikad E; Krishna S
Changes in the physical state and expression of human papillomavirus type 16 in the progression of cervical intraepithelial neoplasia lesions analysed by PCR.
J Gen Virol 76 (Pt 10): 2589-93 (1995)
- Davidson B; Kopolovic J
[Molecular biology of human papilloma virus infection of the female genital tract]
Harefuah 129: 417-20 (1995)
- De Marco E; Marcante ML
Cellular and molecular analyses of interferon beta cytopathic effect on HPV-16 in vitro transformed human keratinocytes (HPK-IA).
J Biol Regul Homeost Agents 9: 24-30 (1995)
- De Marco F; Giannoni F; Marcante ML
Interferon-beta strong cytopathic effect on human papillomavirus type 16-immortalized HPK-IA cell line, unexpectedly not shared by interferon-alpha.
J Gen Virol 76 (Pt 2): 445-50 (1995)
- Delvenne P; al-Saleh W; Gilles C; Thiry A; Boniver J
Inhibition of growth of normal and human papillomavirus-transformed keratinocytes in monolayer and organotypic cultures by interferon-gamma and tumor necrosis factor-alpha.
Am J Pathol 146: 589-98 (1995)
- Demeret C; Le Moal M; Yaniv M; Thierry F
Control of HPV 18 DNA replication by cellular and viral transcription factors.
Nucleic Acids Res 23: 4777-84 (1995)
- de Roda Husman AM; Snijders PJ; Stel HV; van den Brule AJ; Meijer CJ; Walboomers JM
Processing of long-stored archival cervical smears for human papillomavirus detection by the polymerase chain reaction.
Br J Cancer 72: 412-7 (1995)
- Desaintes C; Hallez S; Detremmerie O; Burny A
Wild-type p53 down-regulates transcription from oncogenic human papillomavirus promoters through the epithelial specific enhancer.
Oncogene 10: 2155-61 (1995)
- DiLorenzo TP; Steinberg BM
Differential regulation of human papillomavirus type 6 and 11 early promoters in cultured cells derived from laryngeal papillomas.
J Virol 69: 6865-72 (1995)

- Dillner J; Wiklund F; Lenner P; Eklund C; Frederiksson-Shanazarian V; Schiller JT; Hibma M; Hallmans G; Stendahl U
Antibodies against linear and conformational epitopes of human papillomavirus type 16 that independently associate with incident cervical cancer.
Int J Cancer 60: 377-82 (1995)
- Dillner L; Zellbi A; Avall-Lundqvist E; Heino P; Eklund C; Pettersson CA; Forslund O; Hansson BG; Grandien M; Bistoletti P; et al
Association of serum antibodies against defined epitopes of human papillomavirus L1, E2, and E7 antigens and of HPV DNA with incident cervical cancer.
Cancer Detect Prev 19: 381-93 (1995)
- Donofrio V; Lo Muzio L; Mignogna MD; Troncone G; Staibano S; Boscaino A; De Rosa G
Prognostic evaluation of HPV-associated precancerous and microinvasive carcinoma of the oral cavity: combined use of nucleolar organiser regions (AgNOR) and proliferating cell nuclear antigen (PCNA).
Eur J Cancer B Oral Oncol 31B: 174-80 (1995)
- Dowhanick JJ; McBride AA; Howley PM
Suppression of cellular proliferation by the papillomavirus E2 protein.
J Virol 69: 7791-9 (1995)
- Dreyfus M; Baldauf JJ; Ritter J; Obert G
Seric and local antibodies against a synthetic peptide of HPV16.
Eur J Obstet Gynecol Reprod Biol 59: 187-91 (1995)
- Drijfhout JW; Brandt RM; D'Amaro J; Kast WM; Melief CJ
Detailed motifs for peptide binding to HLA-A*0201 derived from large random sets of peptides using a cellular binding assay.
Hum Immunol 43: 1-12 (1995)
- Drummond-Barbosa DA; Vaillancourt RR; Kazlauskas A; DiMaio D
Ligand-independent activation of the platelet-derived growth factor beta receptor: requirements for bovine papillomavirus E5-induced mitogenic signaling.
Mol Cell Biol 15: 2570-81 (1995)
- Duggan MA; Rowlands C; Kneafsey PD; Nation JG; Stuart GC
Uterine atypical polypoid adenomyoma and ovarian endometrioid carcinoma: metastatic disease or dual primaries?
Int J Gynecol Pathol 14: 81-6 (1995)
- Eklund C; Dillner J
A two-site enzyme immunoassay for quantitation of human papillomavirus type 16 particles.
J Virol Methods 53: 11-23 (1995)
- Ellis JR; Keating PJ; Baird J; Hounsell EF; Renouf DV; Rowe M; Hopkins D; Duggan-Keen MF; Bartholomew JS; Young LS; et al
The association of an HPV16 oncogene variant with HLA-B7 has implications for vaccine design in cervical cancer.
Nat Med 1: 464-70 (1995)
- Ennaji MM; Schwartz JL; Mealing G; Belbaraka L; Parker C; Parentaux M; Jouishomme H; Arella M; Whitfield JF; Phipps J
Alterations in cell-cell communication in human papillomavirus type 16 (HPV16) transformed rat myoblasts.
Cell Mol Biol (Noisy-le-grand) 41: 481-98 (1995)
- Fan S; Smith ML; Rivet DJ 2nd; Duba D; Zhan Q; Kohn KW; Fornace AJ Jr; O'Connor PM
Disruption of p53 function sensitizes breast cancer MCF-7 cells to cisplatin and pentoxifylline.
Cancer Res 55: 1649-54 (1995)
- Farin FM; Bigler LG; Oda D; McDougall JK; Omiecinski CJ
Expression of cytochrome P450 and microsomal epoxide hydrolase in cervical and oral epithelial cells immortalized by human papillomavirus type 16 E6/E7 genes.
Carcinogenesis 16: 1391-401 (1995)
- Farin FM; Bigler LG; Oda D; McDougall JK; Omiecinski CJ
Expression of cytochrome P450 and microsomal epoxide hydrolase in cervical and oral epithelial cells immortalized by human papillomavirus type 16 E6/E7 genes.
Carcinogenesis 16: 1670 (1995)
- Feltkamp MC; Vierboom MP; Toes RE; Ossendorp F; ter Schegget J; Melief CJ; Kast WM
Competition inhibition of cytotoxic T-lymphocyte (CTL) lysis, a more sensitive method to identify candidate CTL epitopes than induction of antibody-detected MHC class I stabilization.
Immunol Lett 47: 1-8 (1995)
- Feltkamp MC; Vreugdenhil GR; Vierboom MP; Ras E; van der Burg SH; ter Schegget J; Melief CJ; Kast WM
Cytotoxic T lymphocytes raised against a subdominant epitope offered as a synthetic peptide eradicate human papillomavirus type 16-induced tumors.
Eur J Immunol 25: 2638-42 (1995)
- Fernando GJ; Stenzel DJ; Tindle RW; Merza MS; Morein B; Frazer IH
Peptide polymerisation facilitates incorporation into ISCOMs and increases antigen-specific IgG2a production.
Vaccine 13: 1460-7 (1995)
- Fernando GJ; Tindle RW; Frazer IH
T-helper epitopes of the E7 transforming protein of cervical cancer associated human papillomavirus type 18 (HPV18).
Virus Res 36: 1-13 (1995)

References (1995)

- Fidalgo PO; Cravo ML; Chaves PP; Leitao CN; Mira FC
High prevalence of human papillomavirus in squamous cell carcinoma and matched normal esophageal mucosa: assessment by polymerase chain reaction.
Cancer 76: 1522-8 (1995)
- Fong KM; Schonrock J; Frazer IM; Zimmerman PV; Smith PJ
Human papillomavirus not found in squamous and large cell lung carcinomas by polymerase chain reaction [letter]
Cancer 75: 2400-1 (1995)
- Fontijn R; Hop C; Brinkman HJ; Slater R; Westerveld A; van Mourik JA; Pannekoek H
Maintenance of vascular endothelial cell-specific properties after immortalization with an amphotrophic replication-deficient retrovirus containing human papilloma virus 16 E6/E7 DNA.
Exp Cell Res 216: 199-207 (1995)
- Franch HA; Shay JW; Alpern RJ; Preisig PA
Involvement of pRB family in TGF beta-dependent epithelial cell hypertrophy.
J Cell Biol 129: 245-54 (1995)
- Frazer IH; Leippe DM; Dunn LA; Liem A; Tindle RW; Fernando GJ; Phelps WC; Lambert PF
Immunological responses in human papillomavirus 16 E6/E7-transgenic mice to E7 protein correlate with the presence of skin disease.
Cancer Res 55: 2635-9 (1995)
- Fujii T; Matsushima Y; Yajima M; Sugimura T; Terada M
Serum antibody against unfused recombinant E7 protein of human papillomavirus type 16 in cervical cancer patients.
Jpn J Cancer Res 86: 28-34 (1995)
- Fujii T; Oguni S; Kikuchi M; Kanai N; Saito K
p53 mutation in carcinomas arising in ovarian cystic teratomas.
Pathol Int 45: 649-54 (1995)
- Fujii T; Tsukazaki K; Kiguchi K; Kubushiro K; Yajima M; Nozawa S
The major E6/E7 transcript of HPV-16 in exfoliated cells from cervical neoplasia patients.
Gynecol Oncol 58: 210-5 (1995)
- Fujita A; Sakagami K; Kanegae Y; Saito I; Kobayashi I
Gene targeting with a replication-defective adenovirus vector.
J Virol 69: 6180-90 (1995)
- Fujiwara H; Mitchell MF; Arseneau J; Hale RJ; Wright TC Jr
Clear cell adenosquamous carcinoma of the cervix. An aggressive tumor associated with human papillomavirus-18.
Cancer 76: 1591-600 (1995)
- Furihata M; Yamasaki I; Ohtsuki Y; Sonobe H; Morioka M; Yamamoto A; Terao N; Kuwahara M; Fujisaki N
p53 and human papillomavirus DNA in renal pelvic and ureteral carcinoma including dysplastic lesions.
Int J Cancer 64: 298-303 (1995)
- Gallego MI; Lazo PA
Deletion in human chromosome region 12q13-15 by integration of human papillomavirus DNA in a cervical carcinoma cell line.
J Biol Chem 270: 24321-6 (1995)
- Gao L; Walter J; Travers P; Stauss H; Chain BM
Tumor-associated E6 protein of human papillomavirus type 16 contains an unusual H-2Kb-restricted cytotoxic T cell epitope.
J Immunol 155: 5519-26 (1995)
- Garry RF
Sequence similarities between latent membrane protein LMP-1 of Epstein-Barr virus, integral membrane protein p12I of human T cell leukemia/lymphotropic virus type 1, E5 transformation protein of bovine papillomavirus, and the transmembrane proteins of slowly transforming retroviruses [letter]
AIDS Res Hum Retroviruses 11: 431-2 (1995)
- Garzetti GG; Ciavattini A; Butini L; Vecchi A; Montroni M
Cervical dysplasia in HIV-seropositive women: role of human papillomavirus infection and immune status.
Gynecol Obstet Invest 40: 52-6 (1995)
- Garzetti GG; Ciavattini A; Goteri G; De Nictolis M; Menso S; Muzzioli M; Fabris N
HPV DNA positivity and natural killer cell activity in the clinical outcome of mild cervical dysplasia: integration between virus and immune system.
Gynecol Obstet Invest 39: 130-5 (1995)
- Geisen C; Delius H; Lichter P; Kahn T
A transcribed human sequence related to the mouse HC1 and the human papillomavirus type 18 E5 genes is located at chromosome 7p13-14.
Hum Mol Genet 4: 1337-45 (1995)
- Giannoudis A; Ergazaki M; Segas J; Giotakis J; Adamopoulos G; Gorgoulis V; Spandidos DA
Detection of Epstein-Barr virus and human papillomavirus in nasopharyngeal carcinoma by the polymerase chain reaction technique.
Cancer Lett 89: 177-81 (1995)
- Gomez MA; Drut R; Lojo MM; Drut RM
Detection of human papillomavirus in juvenile laryngeal papillomatosis using polymerase chain reaction.
Medicina (B Aires) 55: 213-7 (1995)

- Gopalakrishnan V; Walker S; Khan SA
Stimulation of human papillomavirus type 1a DNA replication by a multimerized AT-rich palindromic sequence.
Virology 214: 301-6 (1995)
- Gopalkrishna V; Srivastava AN; Hedau S; Sharma JK; Das BC
Detection of human papillomavirus DNA sequences in cancer of the urinary bladder by in situ hybridisation and polymerase chain reaction.
Genitourin Med 71: 231-3 (1995)
- Greaves RF; Brown JM; Vieira J; Mocarski ES
Selectable insertion and deletion mutagenesis of the human cytomegalovirus genome using the *Escherichia coli* guanosine phosphoribosyl transferase (gpt) gene.
J Gen Virol 76 (Pt 9): 2151-60 (1995)
- Gregoire L; Cubilla AL; Reuter VE; Haas GP; Lancaster WD
Preferential association of human papillomavirus with high-grade histologic variants of penile-invasive squamous cell carcinoma.
J Natl Cancer Inst 87: 1705-9 (1995)
- Grey HM; Ruppert J; Vitiello A; Sidney J; Kast WM; Kubo RT; Sette A
Class I MHC-peptide interactions: structural requirements and functional implications.
Cancer Surv 22: 37-49 (1995)
- Grinstein E; Royer HD
Multiple octamer-binding proteins are targets for the cell cycle-regulated nuclear inhibitor I-92.
DNA Cell Biol 14: 493-500 (1995)
- Gu Z; Matlashewski G
Effect of human papillomavirus type 16 oncogenes on MAP kinase activity.
J Virol 69: 8051-6 (1995)
- Haller K; Stubenrauch F; Pfister H
Differentiation-dependent transcription of the epidermodysplasia verruciformis-associated human papillomavirus type 5 in benign lesions.
Virology 214: 245-55 (1995)
- Havre PA; Yuan J; Hedrick L; Cho KR; Glazer PM
p53 inactivation by HPV16 E6 results in increased mutagenesis in human cells.
Cancer Res 55: 4420-4 (1995)
- Hecht JL; Kadish AS; Jiang G; Burk RD
Genetic characterization of the human papillomavirus (HPV) 18 E2 gene in clinical specimens suggests the presence of a subtype with decreased oncogenic potential.
Int J Cancer 60: 369-76 (1995)
- Hegde RS
Structure of the BPV-1 E2 DNA-binding domain bound to its DNA target.
J Nucl Med 36: 25S-27S (1995)
- Heino P; Dillner J; Schwartz S
Human papillomavirus type 16 capsid proteins produced from recombinant Semliki Forest virus assemble into virus-like particles.
Virology 214: 349-59 (1995)
- Heino P; Skyldberg B; Lehtinen M; Rantala I; Hagmar B; Kreider JW; Kirnbauer R; Dillner J
Human papillomavirus type 16 capsids expose multiple type-restricted and type-common antigenic epitopes.
J Gen Virol 76 (Pt 5): 1141-53 (1995)
- Heinzel PA; Chan SY; Ho L; O'Connor M; Balaram P; Campo MS; Fujinaga K; Kiviat N; Kuypers J; Pfister H; et al
Variation of human papillomavirus type 6 (HPV-6) and HPV-11 genomes sampled throughout the world.
J Clin Microbiol 33: 1746-54 (1995)
- Herrington CS
Human papillomaviruses (HPV) in gynaecological cytology: from molecular biology to clinical testing.
Cytopathology 6: 176-89 (1995)
- Herrington CS
Human papillomaviruses and cervical neoplasia. II. Interaction of HPV with other factors.
J Clin Pathol 48: 1-6 (1995)
- Hibma MH; Raj K; Ely SJ; Stanley M; Crawford L
The interaction between human papillomavirus type 16 E1 and E2 proteins is blocked by an antibody to the N-terminal region of E2.
Eur J Biochem 229: 517-25 (1995)
- Hietanen S; Grenman S; Syrjanen K; Lappalainen K; Kauppinen J; Carey T; Syrjanen S
Human papillomavirus in vulvar and vaginal carcinoma cell lines.
Br J Cancer 72: 134-9 (1995)
- Hietanen SH; Kurvinen K; Syrjanen K; Grenman S; Carey T; McClatchey K; Syrjanen S
Mutation of tumor suppressor gene p53 is frequently found in vulvar carcinoma cells.
Am J Obstet Gynecol 173: 1477-82 (1995)

References (1995)

- Hofmann KJ; Cook JC; Joyce JG; Brown DR; Schultz LD; George HA; Rosolowsky M; Fife KH; Jansen KU
Sequence determination of human papillomavirus type 6a and assembly of virus-like particles in *Saccharomyces cerevisiae*.
Virology 209: 506-18 (1995)
- Holt SE; Wilson VG
Mutational analysis of the 18-base-pair inverted repeat element at the bovine papillomavirus origin of replication: identification of critical sequences for E1 binding and in vivo replication.
J Virol 69: 6525-32 (1995)
- Honig JF; Becker HJ; Brinck U; Korabiowska M
Detection of human papillomavirus DNA sequences in leukocytes: a new approach to identify hematological markers of HPV infection in patients with oral SCC.
Bull Group Int Rech Sci Stomatol Odontol 38: 25-31 (1995)
- Hopfl R; Hristensen ND; Angell MG; Kreider JW
Leukocyte proliferation in vitro against cottontail rabbit papillomavirus in rabbits with persisting papillomas/cancer or after regression.
Arch Dermatol Res 287: 652-8 (1995)
- Horer M; Weger S; Butz K; Hoppe-Seyler F; Geisen C; Kleinschmidt JA
Mutational analysis of adeno-associated virus Rep protein-mediated inhibition of heterologous and homologous promoters.
J Virol 69: 5485-96 (1995)
- Hosone M; Ishiwata T; Nomura N; Kawanami O
[A molecular pathological study on the localization of the human papillomavirus using nested PCR and in situ hybridization methods in paraffin embedded tissues]
Nippon Ika Daigaku Zasshi 62: 131-41 (1995)
- Hsieh CY; Wu CC; Chen TM; Chen CA; Chen CL; Wang JF; Chang CF; Hsieh FJ
Clinical significance of intratumoral blood flow in cervical carcinoma assessed by color Doppler ultrasound.
Cancer 75: 2518-22 (1995)
- Hu G; Liu W; Hanania EG; Fu S; Wang T; Deisseroth AB
Suppression of tumorigenesis by transcription units expressing the antisense E6 and E7 messenger RNA (mRNA) for the transforming proteins of the human papilloma virus and the sense mRNA for the retinoblastoma gene in cervical carcinoma cells.
Cancer Gene Ther 2: 19-32 (1995)
- Huibregtse JM; Scheffner M; Beaudenon S; Howley PM
A family of proteins structurally and functionally related to the E6-AP ubiquitin-protein ligase.
Proc Natl Acad Sci U S A 92: 2563-7 (1995)
- Hummel M; Lim HB; Laimins LA
Human papillomavirus type 31b late gene expression is regulated through protein kinase C-mediated changes in RNA processing.
J Virol 69: 3381-8 (1995)
- Hwang ES; Nottoli T; Dimairo D
The HPV16 E5 protein: expression, detection, and stable complex formation with transmembrane proteins in COS cells.
Virology 211: 227-33 (1995)
- Icenogle JP; Clancy KA; Lin SY
Sequence variation in the capsid protein genes of human papillomavirus type 16 and type 31.
Virology 214: 664-9 (1995)
- Iglesias M; Plowman GD; Woodworth CD
Interleukin-6 and interleukin-6 soluble receptor regulate proliferation of normal, human papillomavirus-immortalized, and carcinoma-derived cervical cells in vitro.
Am J Pathol 146: 944-52 (1995)
- Ikenberg H; Matthay K; Schmitt B; Bauknecht T; Kiechle-Schwarz M; Goppinger A; Pfeleiderer A
p53 mutation and MDM2 amplification are rare even in human papillomavirus-negative cervical carcinomas.
Cancer 76: 57-66 (1995)
- Imura M; Fujiwara T; Ogura H
Human papillomavirus DNA in cell lines derived from malignancies.
Acta Med Okayama 49: 123-7 (1995)
- Jackson ME; Campo MS
Both viral E2 protein and the cellular factor PEBP2 regulate transcription via E2 consensus sites within the bovine papillomavirus type 4 long control region.
J Virol 69: 6038-46 (1995)
- Jacobs MV; de Roda Husman AM; van den Brule AJ; Snijders PJ; Meijer CJ; Walboomers JM
Group-specific differentiation between high- and low-risk human papillomavirus genotypes by general primer-mediated PCR and two cocktails of oligonucleotide probes.
J Clin Microbiol 33: 901-5 (1995)
- Jansen KU; Rosolowsky M; Schultz LD; Markus HZ; Cook JC; Donnelly JJ; Martinez D; Ellis RW; Shaw AR
Vaccination with yeast-expressed cottontail rabbit papillomavirus (CRPV) virus-like particles protects rabbits from CRPV-induced papilloma formation.
Vaccine 13: 1509-14 (1995)

- Jeon S; Allen-Hoffmann BL; Lambert PF
Integration of human papillomavirus type 16 into the human genome correlates with a selective growth advantage of cells.
J Virol 69: 2989-97 (1995)
- Jeon S; Lambert PF
Integration of human papillomavirus type 16 DNA into the human genome leads to increased stability of E6 and E7 mRNAs: implications for cervical carcinogenesis.
Proc Natl Acad Sci U S A 92: 1654-8 (1995)
- Jesudasan RA; Rahman RA; Chandrashekharappa S; Evans GA; Srivatsan ES
Deletion and translocation of chromosome 11q13 sequences in cervical carcinoma cell lines.
Am J Hum Genet 56: 705-15 (1995)
- Karlsen F; Kristensen G; Holm R; Chitemerere M; Berner A; Hagmar BM
High incidence of human papillomavirus in 146 cervical carcinomas. A study using three different pairs of consensus primers, and detecting viral genomes with putative deletions.
Eur J Cancer 31A: 1511-6 (1995)
- Kawana T
[Human papilloma virus and cervical cancer]
Gan To Kagaku Ryoho 22: 711-7 (1995)
- Kelley MJ; Otterson GA; Kaye FJ; Popescu NC; Johnson BE; Dipaolo JA
CDKN2 in HPV-positive and HPV-negative cervical carcinoma cell lines.
Int J Cancer 63: 226-30 (1995)
- Kelsall SR; Kulski JK
Expression of the major capsid protein of human papillomavirus type 16 in *Escherichia coli*.
J Virol Methods 53: 75-90 (1995)
- Kim CJ; Isono T; Tomoyoshi T; Seto A
Variable-region sequences for T-cell receptor-gamma and -delta chains of rabbit killer cell lines against Shope carcinoma cells.
Cancer Lett 89: 37-44 (1995)
- Kim CJ; Tomoyoshi T; Seto A
Varied expression of major histocompatibility complex and oncogenes in Shope carcinoma cell lines derived from a single tumor.
J Cancer Res Clin Oncol 121: 16-22 (1995)
- Kim JW; Cho YH; Lyu MS; Lee CG; Jung JK; Kim SJ; Namkoong SE
Establishment and characterization of a cell line (CUMC-3) derived from a human squamous carcinoma of the uterine cervix.
Gynecol Oncol 57: 47-60 (1995)
- Kim KH; Kim YS
Role of human papillomavirus and p53 tumor suppressor gene in cervical carcinogenesis.
Yonsei Med J 36: 412-25 (1995)
- Kim KH; Kim YS
Analysis of p53 tumor suppressor gene mutations and human papillomavirus infection in human bladder cancers.
Yonsei Med J 36: 322-31 (1995)
- Kim KH; Park TK; Yoon DJ; Kim YS
The effects of wild type p53 tumor suppressor gene expression on the normal human cervical epithelial cells or human epidermal keratinocytes transformed with human papillomavirus type 16 DNA.
Yonsei Med J 36: 287-98 (1995)
- Kinoshita I; Dosaka-Akita H; Shindoh M; Fujino M; Akie K; Kato M; Fujinaga K; Kawakami Y
Human papillomavirus type 18 DNA and E6-E7 mRNA are detected in squamous cell carcinoma and adenocarcinoma of the lung.
Br J Cancer 71: 344-9 (1995)
- Kleine K; Konig G; Kreuzer J; Komitowski D; Zur Hausen H; Rosl F
The effect of the JE (MCP-1) gene, which encodes monocyte chemoattractant protein-1, on the growth of HeLa cells and derived somatic-cell hybrids in nude mice.
Mol Carcinog 14: 179-89 (1995)
- Kondoh G; Hayasaka N; Li Q; Nishimune Y; Hakura A
An in vivo model for receptor tyrosine kinase autocrine/paracrine activation: auto-stimulated KIT receptor acts as a tumor promoting factor in papillomavirus-induced tumorigenesis.
Oncogene 10: 341-7 (1995)
- Koralnik IJ; Mulloy JC; Andresson T; Fullen J; Franchini G
Mapping of the intermolecular association of human T cell leukaemia/lymphotropic virus type I p12I and the vacuolar H⁺-ATPase 16 kDa subunit protein.
J Gen Virol 76 (Pt 8): 1909-16 (1995)
- Kreider JW; Cladel NM; Patrick SD; Welsh PA; DiAngelo SL; Bower JM; Christensen ND
High efficiency induction of papillomas in vivo using recombinant cottontail rabbit papillomavirus DNA.
J Virol Methods 55: 233-44 (1995)

References (1995)

- Kyo S; Tam A; Laimins LA
Transcriptional activity of human papillomavirus type 31b enhancer is regulated through synergistic interaction of AP1 with two novel cellular factors.
Virology 211: 184-97 (1995)
- Labrecque S; Matlashewski GJ
Viability of wild type p53-containing and p53-deficient tumor cells following anticancer treatment: the use of human papillomavirus E6 to target p53.
Oncogene 11: 387-92 (1995)
- Lam KY; Chan AC; Chan KW; Leung ML; Srivastava G
Expression of p53 and its relationship with human papillomavirus in penile carcinomas.
Eur J Surg Oncol 21: 613-6 (1995)
- Lau PW; Cook N; Andrews H; Bracka A; Myint SH
Detection of human papillomavirus types in balanitis xerotica obliterans and other penile conditions.
Genitourin Med 71: 228-30 (1995)
- Le Cann P; Chabaud M; Leboulleux D; Mouglin C; Mayelo V; Legrand MC; Calvet C; Afoutou JM; Coll-Seck AM; Coursaget P
Detection of antibodies to L1, L2, and E4 proteins of human papillomavirus types 6, 11, and 16 by ELISA using synthetic peptides.
J Med Virol 45: 410-4 (1995)
- Ledent C; Marcotte A; Dumont JE; Vassart G; Parmentier M
Differentiated carcinomas develop as a consequence of the thyroid specific expression of a thyroglobulin-human papillomavirus type 16 E7 transgene.
Oncogene 10: 1789-97 (1995)
- Lehtinen M; Hibma MH; Stellato G; Kuoppala T; Paavonen J
Human T helper cell epitopes overlap B cell and putative cytotoxic T cell epitopes in the E2 protein of human papillomavirus type 16.
Biochem Biophys Res Commun 209: 541-6 (1995)
- Levedakou EN; Kaufmann WK; Alcorta DA; Galloway DA; Paules RS
p21CIP1 is not required for the early G2 checkpoint response to ionizing radiation.
Cancer Res 55: 2500-2 (1995)
- Li HX; Zhu WY; Xia MY
Detection with the polymerase chain reaction of human papillomavirus DNA in condylomata acuminata treated with CO2 laser and microwave.
Int J Dermatol 34: 209-11 (1995)
- Li Q; Hu K; Pan X; Cao Z; Yang J; Hu S
Detection of human papillomavirus types 16, 18 DNA related sequences in bronchogenic carcinoma by polymerase chain reaction.
Chin Med J (Engl) 108: 610-4 (1995)
- Liang XH; Mungal S; Ayscue A; Meissner JD; Wodnicki P; Hockenbery D; Lockett S; Herman B
Bcl-2 protooncogene expression in cervical carcinoma cell lines containing inactive p53.
J Cell Biochem 57: 509-21 (1995)
- Lipford GB; Bauer S; Wagner H; Heeg K
Peptide engineering allows cytotoxic T-cell vaccination against human papilloma virus tumour antigen, E6.
Immunology 84: 298-303 (1995)
- Liu JS; Kuo SR; Broker TR; Chow LT
The functions of human papillomavirus type 11 E1, E2, and E2C proteins in cell-free DNA replication.
J Biol Chem 270: 27283-91 (1995)
- Liu Z; Ghai J; Ostrow RS; Faras AJ
The expression levels of the human papillomavirus type 16 E7 correlate with its transforming potential.
Virology 207: 260-70 (1995)
- Lungu O; Sun XW; Wright TC Jr; Ferenczy A; Richart RM; Silverstein S
A polymerase chain reaction-enzyme-linked immunosorbent assay method for detecting human papillomavirus in cervical carcinomas and high-grade cervical cancer precursors.
Obstet Gynecol 85: 337-42 (1995)
- Macdonald MR; Le KT; Freeman J; Hui MF; Cheung RK; Dosch HM
A majority of inverted sinonasal papillomas carries Epstein-Barr virus genomes.
Cancer 75: 2307-12 (1995)
- Maheswaran S; Englert C; Bennett P; Heinrich G; Haber DA
The WT1 gene product stabilizes p53 and inhibits p53-mediated apoptosis.
Genes Dev 9: 2143-56 (1995)
- Mansur CP; Marcus B; Dalal S; Androphy EJ
The domain of p53 required for binding HPV 16 E6 is separable from the degradation domain.
Oncogene 10: 457-65 (1995)
- Mao EJ
Prevalence of human papillomavirus 16 and nucleolar organizer region counts in oral exfoliated cells from normal and malignant epithelia.
Oral Surg Oral Med Oral Pathol Oral Radiol Endod 80: 320-9 (1995)

- Maran A; Amella CA; Di Lorenzo TP; Auburn KJ; Taichman LB; Steinberg BM
Human papillomavirus type 11 transcripts are present at low abundance in latently infected respiratory tissues.
Virology 212: 285-94 (1995)
- Mark HF; Hann E; Mikumo R; Lauchlan S; Beauregard L; Braun L
Cytogenetic characterization of three cell lines derived from primary cervical tumors.
Ann Clin Lab Sci 25: 185-99 (1995)
- Marston NJ; Jenkins JR; Vousden KH
Oligomerisation of full length p53 contributes to the interaction with mdm2 but not HPV E6.
Oncogene 10: 1709-15 (1995)
- Matsukura T; Sugase M
Identification of genital human papillomaviruses in cervical biopsy specimens: segregation of specific virus types in specific clinicopathologic lesions.
Int J Cancer 61: 13-22 (1995)
- May P; May E
[P53 and cancers]
Pathol Biol (Paris) 43: 165-73 (1995)
- Mazzarelli JM; Atkins GB; Geisberg JV; Ricciardi RP
The viral oncoproteins Ad5 E1A, HPV16 E7 and SV40 TAg bind a common region of the TBP-associated factor-110.
Oncogene 11: 1859-64 (1995)
- McGarvie GM; Grindlay GJ; Chandrachud LM; O'Neil BW; Jarrett WF; Campo MS
T cell responses to BPV-4 E7 during infection and mapping of T cell epitopes.
Virology 206: 504-10 (1995)
- McLachlin CM
Pathology of human papillomavirus in the female genital tract.
Curr Opin Obstet Gynecol 7: 24-9 (1995)
- McNicol P; Guijon F; Wayne S; Hidajat R; Paraskevas M
Expression of human papillomavirus type 16 E6-E7 open reading frame varies quantitatively in biopsy tissue from different grades of cervical intraepithelial neoplasia.
J Clin Microbiol 33: 1169-73 (1995)
- Meissner J
Complete nucleotide sequencing of an HPV-1a variant and determination of extant errors in the prototype HPV-1a sequence.
Virus Genes 9: 189-91 (1995)
- Meissner JD
On the nature of papillomavirus hell [letter]
J Infect Dis 172: 895-6 (1995)
- Melendy T; Sedman J; Stenlund A
Cellular factors required for papillomavirus DNA replication.
J Virol 69: 7857-67 (1995)
- Mendoza R; Gandhi L; Botchan MR
E1 recognition sequences in the bovine papillomavirus type 1 origin of DNA replication: interaction between half sites of the inverted repeats.
J Virol 69: 3789-98 (1995)
- Milde-Langosch K; Albrecht K; Joram S; Schlechte H; Giessing M; Loning T
Presence and persistence of HPV infection and p53 mutation in cancer of the cervix uteri and the vulva.
Int J Cancer 63: 639-45 (1995)
- Mitchell MF; Hittelman WK; Lotan R; Nishioka K; Tortolero-Luna G; Richards-Kortum R; Wharton JT; Hong WK
Chemoprevention trials and surrogate end point biomarkers in the cervix.
Cancer 76: 1956-77 (1995)
- Miwa K; Miyamoto S; Kato H; Imamura T; Nishida M; Yoshikawa Y; Nagata Y; Wake N
The role of p53 inactivation in human cervical cell carcinoma development.
Br J Cancer 71: 219-26 (1995)
- Molinari M; Milner J
p53 in complex with DNA is resistant to ubiquitin-dependent proteolysis in the presence of HPV-16 E6.
Oncogene 10: 1849-54 (1995)
- Monk BJ; Burger RA; Lin F; Parham G; Vasilev SA; Wilczynski SP
Prognostic significance of human papillomavirus DNA in vulvar carcinoma.
Obstet Gynecol 85: 709-15 (1995)
- Monsonog J
[Cellular and molecular pathogenesis of cancer of the cervix]
Contracept Fertil Sex 23: 731-40 (1995)
- Montenarh M
Marker genes for cytotoxic exposure: p53.
Stem Cells (Dayt) 13 Suppl 1: 136-41 (1995)
- Montgomery KD; Tedford KL; McDougall JK
Genetic instability of chromosome 3 in HPV-immortalized and tumorigenic human keratinocytes.
Genes Chromosomes Cancer 14: 97-105 (1995)

References (1995)

- Morozov A; Subjeck J; Raychaudhuri P
HPV16 E7 oncoprotein induces expression of a 110 kDa heat shock protein.
FEBS Lett 371: 214-8 (1995)
- Morrione A; DeAngelis T; Baserga R
Failure of the bovine papillomavirus to transform mouse embryo fibroblasts with a targeted disruption of the insulin-like growth factor I receptor genes.
J Virol 69: 5300-3 (1995)
- Moyret-Lalle C; Marçais C; Jacquemier J; Moles JP; Daver A; Soret JY; Jeanteur P; Ozturk M; Theillet C
ras, p53 and HPV status in benign and malignant prostate tumors.
Int J Cancer 64: 124-9 (1995)
- Muller M; Gissmann L; Cristiano RJ; Sun XY; Frazer IH; Jenson AB; Alonso A; Zentgraf H; Zhou J
Papillomavirus capsid binding and uptake by cells from different tissues and species.
J Virol 69: 948-54 (1995)
- Muller M; Viscidi RP; Ulken V; Bavinck JN; Hill PM; Fisher SG; Reid R; Munoz N; Schneider A; Shah KV; et al
Antibodies to the E4, E6, and E7 proteins of human papillomavirus (HPV) type 16 in patients with HPV-associated diseases and in the normal population.
J Invest Dermatol 104: 138-41 (1995)
- Nakagawa S; Watanabe S; Yoshikawa H; Taketani Y; Yoshiike K; Kanda T
Mutational analysis of human papillomavirus type 16 E6 protein: transforming function for human cells and degradation of p53 in vitro.
Virology 212: 535-42 (1995)
- Nakamura T; Ide H; Eguchi R; Hayashi K; Hanyu F; Nagasako K; Yukawa M; Asaka K; Fujimori T; Maeda S
Expression of p53 protein related to human papillomavirus and DNA ploidy in superficial esophageal carcinoma.
Surg Today 25: 591-7 (1995)
- Nilson LA; Gottlieb RL; Polack GW; DiMaio D
Mutational analysis of the interaction between the bovine papillomavirus E5 transforming protein and the endogenous beta receptor for platelet-derived growth factor in mouse C127 cells.
J Virol 69: 5869-74 (1995)
- Nishida M; Miyamoto S; Kato H; Miwa T; Imamura T; Miwa K; Yasumoto S; Barrett JC; Wake N
Transcriptional repression of smooth-muscle alpha-actin gene associated with human papillomavirus type 16 E7 expression.
Mol Carcinog 13: 157-65 (1995)
- Noffsinger AE; Hui YZ; Suzuk L; Yochman LK; Miller MA; Hurlbut P; Gal AA; Fenoglio-Preiser CM
The relationship of human papillomavirus to proliferation and ploidy in carcinoma of the anus.
Cancer 75: 958-67 (1995)
- Noffsinger AE; Suzuk L; Hui YZ; Gal AA; Fenoglio-Preiser CM
Differential sensitivities of E6 type-specific and L1 consensus primers in the detection of human papillomavirus in anal carcinoma.
Mod Pathol 8: 509-14 (1995)
- Nuorva K; Soini Y; Kamel D; Pollanen R; Bloigu R; Vahakangas K; Paakko P
p53 protein accumulation and the presence of human papillomavirus DNA in bronchiolo-alveolar carcinoma correlate with poor prognosis.
Int J Cancer 64: 424-9 (1995)
- Nurnberg W; Artuc M; Nawrath M; Lovric J; Stuting S; Moelling K; Czarnetzki BM; Schadendorf D
Human c-myc is expressed in cervical carcinomas and transactivates the HPV-16 promoter.
Cancer Res 55: 4432-7 (1995)
- Nurnberg W; Artuc M; Vorbrueggen G; Kalkbrenner F; Moelling K; Czarnetzki BM; Schadendorf D
Nuclear proto-oncogene products transactivate the human papillomavirus type 16 promoter.
Br J Cancer 71: 1018-24 (1995)
- O'Connor M; Bernard HU
Oct-1 activates the epithelial-specific enhancer of human papillomavirus type 16 via a synergistic interaction with NF1 at a conserved composite regulatory element.
Virology 207: 77-88 (1995)
- Odunsi K; Terry G; Ho L; Bell J; Cuzick J; Ganesan TS
Association between HLA DQB1 * 03 and cervical intra-epithelial neoplasia.
Mol Med 1: 161-71 (1995)
- Oelze I; Kartenbeck J; Crusius K; Alonso A
Human papillomavirus type 16 E5 protein affects cell-cell communication in an epithelial cell line.
J Virol 69: 4489-94 (1995)
- Oggioni MR; Manganelli R; Contorni M; Tommasino M; Pozzi G
Immunization of mice by oral colonization with live recombinant commensal streptococci.
Vaccine 13: 775-9 (1995)

- Ohe Y; Zhao D; Saijo N; Podack ER
Construction of a novel bovine papillomavirus vector without detectable transforming activity suitable for gene transfer.
Hum Gene Ther 6: 325-33 (1995)
- Orihuela E; Tying SK; Pow-Sang M; Dozier S; Cirelli R; Arany I; Rady P; Sanchez R
Development of human papillomavirus type 16 associated squamous cell carcinoma of the scrotum in a patient with Darier's disease treated with systemic isotretinoin.
J Urol 153: 1940-3 (1995)
- Ostrow RS; Coughlin SM; McGlennen RC; Johnson AN; Ratterree MS; Scheffler J; Yaegashi N; Galloway DA; Faras AJ
Serological and molecular evidence of rhesus papillomavirus type 1 infections in tissues from geographically distinct institutions.
J Gen Virol 76 (Pt 2): 293-9 (1995)
- Padayachee A; Sanders CM; Maitland NJ
A polymerase chain reaction (PCR) investigation of oral verrucae which contain HPV types 2 and 57 by in situ hybridization.
J Oral Pathol Med 24: 329-34 (1995)
- Pan H; Griep AE
Temporally distinct patterns of p53-dependent and p53-independent apoptosis during mouse lens development.
Genes Dev 9: 2157-69 (1995)
- Pao CC; Hor JJ; Wu CJ; Shi YF; Xie X; Lu SM
Human papillomavirus type 18 DNA in gestational trophoblastic tissues and choriocarcinomas.
Int J Cancer 63: 505-9 (1995)
- Pao CC; Lin CY; Yao DS; Tseng CJ
Differential expression of cytokine genes in cervical cancer tissues.
Biochem Biophys Res Commun 214: 1146-51 (1995)
- Park NH; Gujulova CN; Baek JH; Cherrick HM; Shin KH; Min BM
Combined
oral carcinogenicity of HPV-16 and benzo(a)pyrene: an in vitro multistep carcinogenesis model.
Oncogene 10: 2145-53 (1995)
- Park TW; Fujiwara H; Wright TC
Molecular biology of cervical cancer and its precursors.
Cancer 76: 1902-13 (1995)
- Payne D; Hoskins S; Schouten H; van Vleuten H; Tying S
Increased buffer pH enhances sensitivity and specificity of human papillomavirus detection using consensus primer based PCR.
J Virol Methods 52: 105-10 (1995)
- Peacock JW; Chung S; Bristow RG; Hill RP; Benchimol S
The p53-mediated G1 checkpoint is retained in tumorigenic rat embryo fibroblast clones transformed by the human papillomavirus type 16 E7 gene and EJ-ras.
Mol Cell Biol 15: 1446-54 (1995)
- Peng X; Lang CM; Kreider JW
Methylation of cottontail rabbit papillomavirus DNA and tissue-specific expression in transgenic rabbits.
Virus Res 35: 101-8 (1995)
- Perea SE; Lopez-Ocejo O; Garcia-Milian R; Arana MJ
Interferon-alpha elicits downregulation of human papillomavirus 18 mRNA in HeLa cells by selective repression of endogenous viral transcription.
J Interferon Cytokine Res 15: 495-501 (1995)
- Peto M; Tolle-Ersu I; Kreysch HG; Klock G
Epidermal growth factor induction of human papillomavirus type 16 E6/E7 mRNA in tumor cells involves two AP-1 binding sites in the viral enhancer.
J Gen Virol 76 (Pt 8): 1945-58 (1995)
- Piccini A; Storey A; Massimi P; Banks L
Mutations in the human papillomavirus type 16 E2 protein identify multiple regions of the protein involved in binding to E1.
J Gen Virol 76 (Pt 11): 2909-13 (1995)
- Pierrefite V; Cuzin F
Replication efficiency of bovine papillomavirus type 1 DNA depends on cis-acting sequences distinct from the replication origin.
J Virol 69: 7682-7 (1995)
- Pilotti S; D'Amato L; Della Torre G; Donghi R; Longoni A; Giarola M; Sampietro G; De Palo G; Pierotti MA; Rilke F
Papillomavirus, p53 alteration, and primary carcinoma of the vulva.
Diagn Mol Pathol 4: 239-48 (1995)
- Pizzighella S; Pisoni G; Bevilacqua F; Vaona A; Palu G
Simultaneous polymerase chain reaction detection and restriction typing for the diagnosis of human genital papillomavirus infection.
J Virol Methods 55: 245-56 (1995)
- Podust VN; Podust LM; Muller F; Hubscher U
DNA polymerase delta holoenzyme: action on single-stranded DNA and on double-stranded DNA in the presence of replicative DNA helicases.
Biochemistry 34: 5003-10 (1995)

References (1995)

- Poljak M; Orłowska J; Cerar A
Human papillomavirus infection in esophageal squamous cell papillomas: a study of 29 lesions.
Anticancer Res 15: 965-9 (1995)
- Radulescu RT; Bellitti MR; Ruvo M; Cassani G; Fassina G
Binding of the LXCXE insulin motif to a hexapeptide derived from retinoblastoma protein.
Biochem Biophys Res Commun 206: 97-102 (1995)
- Rady PL; Arany I; Hughes TK; Tyring SK
Type-specific primer-mediated direct sequencing of consensus primer-generated PCR amplicons of human papilloma viruses: a new approach for the simultaneous detection of multiple viral type infections.
J Virol Methods 53: 245-54 (1995)
- Ramael M; Segers K; Pannemans N; Wesling F; Van Marck E
Detection of human papillomavirus in cervical scrapings by in situ hybridization and the polymerase chain reaction in relation to cytology.
Histochem J 27: 54-9 (1995)
- Ramesar JE; Rybicki EP; Williamson AL
Sequence variation in the L1 gene of human papillomavirus type 16 from Africa.
Arch Virol 140: 1863-70 (1995)
- Rank NM; Lambert PF
Bovine papillomavirus type 1 E2 transcriptional regulators directly bind two cellular transcription factors, TFIID and TFIIB.
J Virol 69: 6323-34 (1995)
- Ranki A; Lassus J; Niemi KM
Relation of p53 tumor suppressor protein expression to human papillomavirus (HPV) DNA and to cellular atypia in male genital warts and in premalignant lesions.
Acta Derm Venereol 75: 180-6 (1995)
- Ravnan JB; Cohen SN
Transformed mouse cell lines that consist predominantly of cells maintaining bovine papilloma virus at high copy number.
Virology 213: 526-34 (1995)
- Ressing ME; Sette A; Brandt RM; Ruppert J; Wentworth PA; Hartman M; Oseroff C; Grey HM; Melief CJ; Kast WM
Human CTL epitopes encoded by human papillomavirus type 16 E6 and E7 identified through in vivo and in vitro immunogenicity studies of HLA-A*0201-binding peptides.
J Immunol 154: 5934-43 (1995)
- Riese DJ 2nd; DiMaio D
An intact PDGF signaling pathway is required for efficient growth transformation of mouse C127 cells by the bovine papillomavirus E5 protein.
Oncogene 10: 1431-9 (1995)
- Rolfe M; Beer-Romero P; Glass S; Eckstein J; Berdo I; Theodoras A; Pagano M; Draetta G
Reconstitution of p53-ubiquitinylation reactions from purified components: the role of human ubiquitin-conjugating enzyme UBC4 and E6-associated protein (E6AP).
Proc Natl Acad Sci U S A 92: 3264-8 (1995)
- Roman A; Brown D
Sequence variation in the extreme 5' end of the human papillomavirus type 6a long control region.
J Infect Dis 171: 697-700 (1995)
- Rorke EA; Jacobberger JW
Transforming growth factor-beta 1 (TGF beta 1) enhances apoptosis in human papillomavirus type 16-immortalized human ectocervical epithelial cells.
Exp Cell Res 216: 65-72 (1995)
- Rose BR; Thompson CH; Simpson JM; Jarrett CS; Elliott PM; Tattersall MH; Dalrymple C; Cossart YE
Human papillomavirus deoxyribonucleic acid as a prognostic indicator in early-stage cervical cancer: a possible role for type 18.
Am J Obstet Gynecol 173: 1461-8 (1995)
- Rose BR; Thompson CH; Tattersall MH; Elliott PM; Dalrymple C; Cossart YE
Identification of E6/E7 transcription patterns in HPV 16-positive cervical cancers using the reverse transcription/polymerase chain reaction.
Gynecol Oncol 56: 239-44 (1995)
- Routes JM; Ryan S
Oncogenicity of human papillomavirus- or adenovirus-transformed cells correlates with resistance to lysis by natural killer cells.
J Virol 69: 7639-47 (1995)
- Rubben A; Traidl C; Baron JM; Grussendorf-Conen EI
Evaluation of non-radioactive temperature gradient SSCP analysis and of temperature gradient gel electrophoresis for the detection of HPV 6-variants in condylomata acuminata and Buschke-Loewenstein tumours.
Eur J Epidemiol 11: 501-6 (1995)
- Runnebaum IB; Maier S; Tong XW; Rosenthal HE; Mobus VJ; Kieback DG; Kreienberg R
Human papillomavirus integration is not associated with advanced epithelial ovarian cancer in German patients [letter]
Cancer Epidemiol Biomarkers Prev 4: 573-5 (1995)

- Russell J; Botchan MR
cis-Acting components of human papillomavirus (HPV) DNA replication: linker substitution analysis of the HPV type 11 origin.
J Virol 69: 651-60 (1995)
- Russell KJ; Wiens LW; Demers GW; Galloway DA; Plon SE; Groudine M
Abrogation of the G2 checkpoint results in differential radiosensitization of G1 checkpoint-deficient and G1 checkpoint-competent cells.
Cancer Res 55: 1639-42 (1995)
- Saegusa M; Takano Y; Hashimura M; Shoji Y; Okayasu I
The possible role of bcl-2 expression in the progression of tumors of the uterine cervix.
Cancer 76: 2297-303 (1995)
- Saito J; Sumiyoshi M; Nakatani H; Ikeda M; Hoshiai H; Noda K
Dysplasia and HPV infection initially detected by DNA analysis in cytologically normal cervical smears.
Int J Gynaecol Obstet 51: 43-8 (1995)
- Salam M; Rockett J; Morris A
The prevalence of different human papillomavirus types and p53 mutations in laryngeal carcinomas: is there a reciprocal relationship?
Eur J Surg Oncol 21: 290-6 (1995)
- Salam MA; Rockett J; Morris A
General primer-mediated polymerase chain reaction for simultaneous detection and typing of human papillomavirus DNA in laryngeal squamous cell carcinomas.
Clin Otolaryngol 20: 84-8 (1995)
- Samoylova EV; Shaikhaiev GO; Petrov SV; Kissel'jova NP; Kissel'jov FL
HPV infection in cervical-cancer cases in Russia.
Int J Cancer 61: 337-41 (1995)
- Sanders CM; Stern PL; Maitland NJ
Characterization of human papillomavirus type 16 E2 protein and subdomains expressed in insect cells.
Virology 211: 418-33 (1995)
- Sarafi TR; McBride AA
Domains of the BPV-1 E1 replication protein required for origin-specific DNA binding and interaction with the E2 transactivator.
Virology 211: 385-96 (1995)
- Sasagawa T; Pushko P; Steers G; Gschmeissner SE; Hajibagheri MA; Finch J; Crawford L; Tommasino M
Synthesis and assembly of virus-like particles of human papillomaviruses type 6 and type 16 in fission yeast *Schizosaccharomyces pombe*.
Virology 206: 126-35 (1995)
- Sastre-Garau X; Couturier J; Favre M; Orth G
A recurrent human papillomavirus integration site at chromosome region 12q14-q15 in SW756 and SK-v cell lines derived from genital tumors.
C R Acad Sci III 318: 475-8 (1995)
- Scheffner M; Nuber U; Huibregtse JM
Protein ubiquitination involving an E1-E2-E3 enzyme ubiquitin thioester cascade.
Nature 373: 81-3 (1995)
- Sciubba JJ
Oral leukoplakia.
Crit Rev Oral Biol Med 6: 147-60 (1995)
- Sears WL; Goto-Mandeville R; Mirapuri M; Braun L
Effects of 12-O-tetradecanoylphorbol-13-acetate on human papillomavirus type 16-positive keratinocytes at different stages of transformation.
Mol Carcinog 13: 146-56 (1995)
- Sedman J; Stenlund A
Co-operative interaction between the initiator E1 and the transcriptional activator E2 is required for replicator specific DNA replication of bovine papillomavirus in vivo and in vitro.
EMBO J 14: 6218-28 (1995)
- Shao J; Zhou M; Ye Y
[Detection of human papillomavirus in condyloma acuminata tissues]
Chung Hua Ping Li Hsueh Tsa Chih 24: 46-8 (1995)
- Shay JW; Tomlinson G; Piatyszek MA; Gollahon LS
Spontaneous in vitro immortalization of breast epithelial cells from a patient with Li-Fraumeni syndrome.
Mol Cell Biol 15: 425-32 (1995)
- Shindoh M; Chiba I; Yasuda M; Saito T; Funaoka K; Kohgo T; Amemiya A; Sawada Y; Fujinaga K
Detection of human papillomavirus DNA sequences in oral squamous cell carcinomas and their relation to p53 and proliferating cell nuclear antigen expression.
Cancer 76: 1513-21 (1995)

References (1995)

- Shirasawa H; Kinoshita T; Shino Y; Mori K; Shimizu K; Simizu B
Cloning and sequencing of the murine farnesyltransferase alpha-encoding cDNA from a cell line which expresses the human papillomavirus type-16 E6 gene.
Gene 164: 373-4 (1995)
- Sibbet GJ; Cuthill S; Campo MS
The enhancer in the long control region of human papillomavirus type 16 is up-regulated by PEF-1 and down-regulated by Oct-1.
J Virol 69: 4006-11 (1995)
- Simons AM; Mugica van Herckenrode C; Rodriguez JA; Maitland N; Anderson M; Phillips DH; Coleman DV
Demonstration of smoking-related DNA damage in cervical epithelium and correlation with human papillomavirus type 16, using exfoliated cervical cells.
Br J Cancer 71: 246-9 (1995)
- Sizemore N; Mukhtar H; Couch LH; Howard PC; Rorke EA
Differential response of normal and HPV immortalized ectocervical epithelial cells to B[a]P.
Carcinogenesis 16: 2413-8 (1995)
- Skomedal H; Kristensen G; Helland A; Nesland JM; Kooi S; Borresen AL; Holm R
TP53 gene mutations and protein accumulation in primary vaginal carcinomas.
Br J Cancer 72: 129-33 (1995)
- Smith LH; Foster C; Hitchcock ME; Leiserowitz GS; Hall K; Isseroff R; Christensen ND; Kreider JW
Titration of HPV-11 infectivity and antibody neutralization can be measured in vitro.
J Invest Dermatol 105: 438-44 (1995)
- Smith ML; Chen IT; Zhan Q; O'Connor PM; Fornace AJ Jr
Involvement of the p53 tumor suppressor in repair of u.v.-type DNA damage.
Oncogene 10: 1053-9 (1995)
- Smits HL; Tjong-A-Hung SP; ter Schegget J; Nooter K; Kok T
Absence of human papillomavirus DNA from esophageal carcinoma as determined by multiple broad spectrum polymerase chain reactions.
J Med Virol 46: 213-5 (1995)
- Smits HL; van Gemen B; Schukkink R; van der Velden J; Tjong-A-Hung SP; Jebbink MF; ter Schegget J
Application of the NASBA nucleic acid amplification method for the detection of human papillomavirus type 16 E6-E7 transcripts.
J Virol Methods 54: 75-81 (1995)
- Somers GR; Tabrizi SN; Tiedemann K; Chow CW; Garland SM; Venter DJ
Squamous cell carcinoma of the tongue in a child with Fanconi anemia: a case report and review of the literature.
Pediatr Pathol Lab Med 15: 597-607 (1995)
- Sparkowski J; Anders J; Schlegel R
E5 oncoprotein retained in the endoplasmic reticulum/cis Golgi still induces PDGF receptor autophosphorylation but does not transform cells.
EMBO J 14: 3055-63 (1995)
- Stacey SN; Jordan D; Snijders PJ; Mackett M; Walboomers JM; Arrand JR
Translation of the human papillomavirus type 16 E7 oncoprotein from bicistronic mRNA is independent of splicing events within the E6 open reading frame.
J Virol 69: 7023-31 (1995)
- Staebler A; Pierce JH; Brazinski S; Heidaran MA; Li W; Schlegel R; Goldstein DJ
Mutational analysis of the beta-type platelet-derived growth factor receptor defines the site of interaction with the bovine papillomavirus type 1 E5 transforming protein.
J Virol 69: 6507-17 (1995)
- Stancovski I; Gonen H; Orian A; Schwartz AL; Ciechanover A
Degradation of the proto-oncogene product c-Fos by the ubiquitin proteolytic system in vivo and in vitro: identification and characterization of the conjugating enzymes.
Mol Cell Biol 15: 7106-16 (1995)
- Stastny JF; Ben-Ezra J; Stewart JA; Kornstein MJ; Kay S; Frable WJ
Condyloma and cervical intraepithelial neoplasia of the endometrium.
Gynecol Obstet Invest 39: 277-80 (1995)
- Steenbergen RD; Hermsen MA; Walboomers JM; Joenje H; Arwert F; Meijer CJ; Snijders PJ
Integrated human papillomavirus type 16 and loss of heterozygosity at 11q22 and 18q21 in an oral carcinoma and its derivative cell line.
Cancer Res 55: 5465-71 (1995)
- Steger G; Ham J; Lefebvre O; Yaniv M
The bovine papillomavirus 1 E2 protein contains two activation domains: one that interacts with TBP and another that functions after TBP binding.
EMBO J 14: 329-40 (1995)
- Steinbach E
[Pathogenesis of cholesteatoma (comment)]
HNO 43: 1-2 (1995)

- Storey A; Massimi P; Dawson K; Banks L
Conditional immortalization of primary cells by human papillomavirus type 18 E6 and EJ-ras defines an E6 activity in G0/G1 phase which can be substituted for mutations in p53. *Oncogene* 11: 653-61 (1995)
- Straight SW; Herman B; McCance DJ
The E5 oncoprotein of human papillomavirus type 16 inhibits the acidification of endosomes in human keratinocytes. *J Virol* 69: 3185-92 (1995)
- Street D; Delgado G
The role of p53 and HPV in cervical cancer [editorial] *Gynecol Oncol* 58: 287-8 (1995)
- Stremlau A; Helms J; Muller-Hermelink HK; Hoppe F; de Villiers EM
[Detection of DNA of human papillomaviruses (HPV) in an "aggressively" growing cholesteatoma. Is cholesteatoma a virus-induced tumor? (see comments)] *HNO* 43: 3-5 (1995)
- Sugimachi K; Sumiyoshi K; Nozoe T; Yasuda M; Watanabe M; Kitamura K; Tsutsui S; Mori M; Kuwano H
Carcinogenesis and histogenesis of esophageal carcinoma. *Cancer* 75: 1440-5 (1995)
- Sumantran VN; Ealovega MW; Nunez G; Clarke MF; Wicha MS
Overexpression of Bcl-XS sensitizes MCF-7 cells to chemotherapy-induced apoptosis. *Cancer Res* 55: 2507-10 (1995)
- Sun XW; Ellerbrock TV; Lungu O; Chiasson MA; Bush TJ; Wright TC Jr
Human papillomavirus infection in human immunodeficiency virus-seropositive women. *Obstet Gynecol* 85: 680-6 (1995)
- Sun XY; Frazer I; Muller M; Gissmann L; Zhou J
Sequences required for the nuclear targeting and accumulation of human papillomavirus type 6B L2 protein. *Virology* 213: 321-7 (1995)
- Sun Y; Si L
[The detection of p53 gene and HPV16 oncogene in cervical carcinoma] *Chung Hua Ping Li Hsueh Tsa Chih* 24: 29-32 (1995)
- Suzich JA; Ghim SJ; Palmer-Hill FJ; White WI; Tamura JK; Bell JA; Newsome JA; Jenson AB; Schlegel R
Systemic immunization with papillomavirus L1 protein completely prevents the development of viral mucosal papillomas. *Proc Natl Acad Sci U S A* 92: 11553-7 (1995)
- Suzuki T; Tomita Y; Nakano K; Shirasawa H; Simizu B
Deletion in the L1 open reading frame of human papillomavirus type 6a genomes associated with recurrent laryngeal papilloma. *J Med Virol* 47: 191-7 (1995)
- Sverdrup F; Khan SA
Two E2 binding sites alone are sufficient to function as the minimal origin of replication of human papillomavirus type 18 DNA. *J Virol* 69: 1319-23 (1995)
- Tan TM; Ting RC
In vitro and in vivo inhibition of human papillomavirus type 16 E6 and E7 genes. *Cancer Res* 55: 4599-605 (1995)
- Tan W; Felber BK; Zolotukhin AS; Pavlakis GN; Schwartz S
Efficient expression of the human papillomavirus type 16 L1 protein in epithelial cells by using Rev and the Rev-responsive element of human immunodeficiency virus or the cis-acting transactivation element of simian retrovirus type 1. *J Virol* 69: 5607-20 (1995)
- Tan W; Schwartz S
The Rev protein of human immunodeficiency virus type 1 counteracts the effect of an AU-rich negative element in the human papillomavirus type 1 late 3' untranslated region. *J Virol* 69: 2932-45 (1995)
- Ten Hagen KG; Ravnán JB; Cohen SN
Disparate replication properties of integrated and extrachromosomal forms of bovine papilloma virus in ID13 cells. *J Mol Biol* 254: 119-29 (1995)
- Tendler Y; Schwartz Y; Reshef R; Shasha SM; Rotter V; Shkolnik T
Immunohistochemical detection of p53 protein expression in HPV-induced condyloma acuminatum. *Acta Derm Venereol* 75: 177-9 (1995)
- Tenti P; Romagnoli S; Silini E; Pellegata NS; Zappatore R; Spinillo A; Zara C; Ranzani GN; Carnevali L
Analysis and clinical implications of K-ras gene mutations and infection with human papillomavirus types 16 and 18 in primary adenocarcinoma of the uterine cervix. *Int J Cancer* 64: 9-13 (1995)
- ter Harmse B; van Belkum A; Quint W; Pronk A; Kuijpers J; Ramaekers F; Tandon A; Smedts F
p53 and human papilloma virus type 16 in cervical intraepithelial neoplasia and carcinoma. *Int J Gynecol Pathol* 14: 125-33 (1995)

References (1995)

- Thomas M; Massimi P; Jenkins J; Banks L
HPV-18 E6 mediated inhibition of p53 DNA binding activity is independent of E6 induced degradation.
Oncogene 10: 261-8 (1995)
- Tindle RW; Croft S; Herd K; Malcolm K; Geczy AF; Stewart T; Fernando GJ
A vaccine conjugate of 'ISCAR' immunocarrier and peptide epitopes of the E7 cervical cancer-associated protein of human papillomavirus type 16 elicits specific Th1- and Th2-type responses in immunized mice in the absence of oil-based adjuvants.
Clin Exp Immunol 101: 265-71 (1995)
- Togawa K; Rustgi AK
A novel human papillomavirus sequence based on L1 general primers.
Virus Res 36: 293-7 (1995)
- Togawa K; Rustgi AK
Human papillomavirus-16 and -18 replication in esophagus squamous cancer cell lines does not require heterologous E1 and E2 proteins.
J Med Virol 45: 435-8 (1995)
- Tommasino M; Crawford L
Human papillomavirus E6 and E7: proteins which deregulate the cell cycle.
Bioessays 17: 509-18 (1995)
- Troungos C; Horti M; Kittas C
A rapid method for the screening and typing of high risk HPVs using molecular biology techniques.
Anticancer Res 15: 2045-8 (1995)
- Tsang NM; Nagasawa H; Li C; Little JB
Abrogation of p53 function by transfection of HPV16 E6 gene enhances the resistance of human diploid fibroblasts to ionizing radiation.
Oncogene 10: 2403-8 (1995)
- Tsao SW; Mok SC; Fey EG; Fletcher JA; Wan TS; Chew EC; Muto MG; Knapp RC; Berkowitz RS
Characterization of human ovarian surface epithelial cells immortalized by human papilloma viral oncogenes (HPV-E6E7 ORFs).
Exp Cell Res 218: 499-507 (1995)
- Tsao YP; Huang CH; Lin YK; Chen SL
Protein kinase C-and ras-dependent activation of c-jun gene by human papillomavirus type 11 E5a oncoprotein.
Cancer Lett 95: 201-5 (1995)
- Tsao YP; Kuo SW; Li SF; Tsai TC; Li LY; Chen SL
Human papillomavirus 11 E5a delays the growth restriction induced by temperature shift in temperature-sensitive simian virus 40 T antigen-immortalized keratinocytes.
Biochem Biophys Res Commun 217: 712-20 (1995)
- Tsuda H; Jiko K; Tsugane S; Yajima M; Yamada T; Tanemura K; Tsunematsu R; Ohmi K; Sonoda T; Hirohashi S
Prognostic value of p53 protein accumulation in cancer cell nuclei in adenocarcinoma of the uterine cervix.
Jpn J Cancer Res 86: 1049-53 (1995)
- Turner MA; Palefsky JM
Urokinase plasminogen activator expression by primary and HPV 16-transformed keratinocytes.
Clin Exp Metastasis 13: 260-8 (1995)
- Unger ER; Vernon SD; Thoms WW; Nisenbaum R; Spann CO; Horowitz IR; Icenogle JP; Reeves WC
Human papillomavirus and disease-free survival in FIGO stage Ib cervical cancer.
J Infect Dis 172: 1184-90 (1995)
- Vande Pol SB; Howley PM
Negative regulation of the bovine papillomavirus E5, E6, and E7 oncogenes by the viral E1 and E2 genes.
J Virol 69: 395-402 (1995)
- Vandebriel RJ; van der Kolk M; Geerse L; Steerenberg PA; Krul MR
A helper T-cell epitope of the E7 protein of human papillomavirus type 16 in BALB/c mice.
Virus Res 37: 13-22 (1995)
- van Belkum A
Low-stringency single specific primer PCR, DNA sequencing and single-strand conformation polymorphism of PCR products for identification of genetic variants of human papillomavirus type 16.
J Virol Methods 55: 435-43 (1995)
- van Belkum A; Juffermans L; Schrauwen L; van Doornum G; Burger M; Quint W
Genotyping human papillomavirus type 16 isolates from persistently infected promiscuous individuals and cervical neoplasia patients.
J Clin Microbiol 33: 2957-62 (1995)

- van Beurden M; ten Kate FJ; Smits HL; Berkhout RJ; de Craen AJ; van der Vange N; Lammes FB; ter Schegget J
Multifocal vulvar intraepithelial neoplasia grade III and multicentric lower genital tract neoplasia is associated with transcriptionally active human papillomavirus.
Cancer 75: 2879-84 (1995) Villa LL; Caballero OL; Levi JE; Pena SD; Simpson AJ
An approach to human papillomavirus identification using low stringency single specific primer PCR.
Mol Cell Probes 9: 45-8 (1995)
- Vivanco MD; Johnson R; Galante PE; Hanahan D; Yamamoto KR
A transition in transcriptional activation by the glucocorticoid and retinoic acid receptors at the tumor stage of dermal fibrosarcoma development.
EMBO J 14: 2217-28 (1995)
- Volpers C; Sapp M; Snijders PJ; Walboomers JM; Streeck RE
Conformational and linear epitopes on virus-like particles of human papillomavirus type 33 identified by monoclonal antibodies to the minor capsid protein L2.
J Gen Virol 76 (Pt 11): 2661-7 (1995)
- Voog E; Ricksten A; Lowhagen GB
Prevalence of Epstein-Barr virus and human papillomavirus in cervical samples from women attending an STD-clinic.
Int J STD AIDS 6: 208-10 (1995)
- Vousden KH
Regulation of the cell cycle by viral oncoproteins.
Semin Cancer Biol 6: 109-16 (1995)
- Wazer DE; Liu XL; Chu Q; Gao Q; Band V
Immortalization of distinct human mammary epithelial cell types by human papilloma virus 16 E6 or E7.
Proc Natl Acad Sci U S A 92: 3687-91 (1995)
- Wikstrom A; van Doornum GJ; Kirnbauer R; Quint WG; Dillner J
Prospective study on the development of antibodies against human papillomavirus type 6 among patients with condyloma acuminata or new asymptomatic infection.
J Med Virol 46: 368-74 (1995)
- Williams AT; Sexton CJ; Hanna NF; Leigh IM
Upregulation of integrin expression in benign vulvar warts.
J Pathol 175: 311-7 (1995)
- Wilson SE; Weng J; Blair S; He YG; Lloyd S
Expression of E6/E7 or SV40 large T antigen-coding oncogenes in human corneal endothelial cells indicates regulated high-proliferative capacity.
Invest Ophthalmol Vis Sci 36: 32-40 (1995)
- Woodworth CD; McMullin E; Iglesias M; Plowman GD
Interleukin 1 alpha and tumor necrosis factor alpha stimulate autocrine amphiregulin expression and proliferation of human papillomavirus-immortalized and carcinoma-derived cervical epithelial cells.
Proc Natl Acad Sci U S A 92: 2840-4 (1995)
- Wu TC; Guarnieri FG; Staveley-O'Carroll KF; Viscidi RP; Levitsky HI; Hedrick L; Cho KR; August JT; Pardoll DM
Engineering an intracellular pathway for major histocompatibility complex class II presentation of antigens.
Proc Natl Acad Sci U S A 92: 11671-5 (1995)
- Xi LF; Demers GW; Koutsky LA; Kiviat NB; Kuypers J; Watts DH; Holmes KK; Galloway DA
Analysis of human papillomavirus type 16 variants indicates establishment of persistent infection.
J Infect Dis 172: 747-55 (1995)
- Xu C; Meikrantz W; Schlegel R; Sager R
The human papilloma virus 16E6 gene sensitizes human mammary epithelial cells to apoptosis induced by DNA damage.
Proc Natl Acad Sci U S A 92: 7829-33 (1995)
- Yadav M; Nurhayati ZA; Padmanathan A; Abdul Aziz Y; Norhanom AW
Polymerase chain reaction detection and restriction enzyme typing of human papillomavirus in cervical carcinoma.
Med J Malaysia 50: 64-71 (1995)
- Yamada T; Wheeler CM; Halpern AL; Stewart AC; Hildesheim A; Jenison SA
Human papillomavirus type 16 variant lineages in United States populations characterized by nucleotide sequence analysis of the E6, L2, and L1 coding segments.
J Virol 69: 7743-53 (1995)
- Yamada T; Yamashita T; Nishikawa T; Fujimoto S; Fujinaga K
Biologic activity of human papillomavirus type 16 E6/E7 cDNA clones isolated from SiHa cervical carcinoma cell line.
Virus Genes 10: 15-25 (1995)
- Yang JT; Liu CZ; Iannaccone P
The HPV 16 genome induces carcinomas and T-cell lymphomas in transgenic mice.
Am J Pathol 147: 68-78 (1995)
- Yasumoto S
[Human papillomavirus type 16-gene functions relevant to molecular human carcinogenesis]
Nippon Rinsho 53: 2858-67 (1995)

References (1995)

- Yeudall WA; Paterson IC; Patel V; Prime SS
Presence of human papillomavirus sequences in tumour-derived human oral keratinocytes expressing mutant p53.
Eur J Cancer B Oral Oncol 31B: 136-43 (1995)
- Ylitalo N; Bergstrom T; Gyllensten U
Detection of genital human papillomavirus by single-tube nested PCR and type-specific oligonucleotide hybridization.
J Clin Microbiol 33: 1822-8 (1995)
- Yokoyama M; Nakao Y; Yang X; Sun Q; Tsutsumi K; Pater A; Pater MM
Alterations in physical state and expression of human papillomavirus type 18 DNA following crisis and establishment of immortalized ectocervical cells.
Virus Res 37: 139-51 (1995)
- Zerfass K; Schulze A; Spitkovsky D; Friedman V; Henglein B; Jansen-Durr P
Sequential activation of cyclin E and cyclin A gene expression by human papillomavirus type 16 E7 through sequences necessary for transformation.
J Virol 69: 6389-99 (1995)
- Zhang J; Rose BR; Thompson CH; Jarrett C; Russell P; Houghton RS; Cossart YE
Associations between oncogenic human papillomaviruses and local invasive patterns in cervical cancer.
Gynecol Oncol 57: 170-7 (1995)
- Zhang T; Chu Y; Yuan Y
[Relations between human papillomavirus and oral cancers using of nucleic acid hybridization]
Chung Hua Kou Chiang Hsueh Tsa Chih 30: 37-9 (1995)
- Zheng J; Saksela O; Matikainen S; Vaheri A
Keratinocyte growth factor is a bifunctional regulator of HPV16 DNA-immortalized cervical epithelial cells.
J Cell Biol 129: 843-51 (1995)
- Zheng J; Vaheri A
Human skin fibroblasts induce anchorage-independent growth of HPV-16-DNA-immortalized cervical epithelial cells.
Int J Cancer 61: 658-65 (1995)
- Zheng PS; Li SR; Iwasaka T; Song J; Cui MH; Sugimori H
Simultaneous detection by consensus multiplex PCR of high- and low-risk and other types of human papilloma virus in clinical samples.
Gynecol Oncol 58: 179-83 (1995)
- Zhu X; Tommasino M; Vousden K; Sadovnikava E; Rappuoli R; Crawford L; Kast M; Melief CJ; Beverley PC; Stauss HJ
Both immunization with protein and recombinant vaccinia virus can stimulate CTL specific for the E7 protein of human papilloma virus 16 in H-2d mice.
Scand J Immunol 42: 557-63 (1995)

- Akutsu N; Shirasawa H; Asano T; Isono K; Simizu B
p53-Dependent and -independent transactivation by the E6 protein of human papillomavirus type 16.
J Gen Virol 77 (Pt 3): 459-63 (1996)
- Antinore MJ; Birrer MJ; Patel D; Nader L; McCance DJ
The human papillomavirus type 16 E7 gene product interacts with and trans-activates the AP1 family of transcription factors.
EMBO J 15: 1950-60 (1996)
- Arbeit JM; Howley PM; Hanahan D
Chronic estrogen-induced cervical and vaginal squamous carcinogenesis in human papillomavirus type 16 transgenic mice.
Proc Natl Acad Sci U S A 93: 2930-5 (1996)
- Bauer-Hofmann R; Borghouts C; Auvinen E; Bourda E; Rosl F; Alonso A
Genomic cloning and characterization of the nonoccupied allele corresponding to the integration site of human papillomavirus type 16 DNA in the cervical cancer cell line SiHa.
Virology 217: 33-41 (1996)
- Behbakht K; DeGeest K; Turyk ME; Wilbanks GD
All-trans-retinoic acid inhibits the proliferation of cell lines derived from human cervical neoplasia.
Gynecol Oncol 61: 31-9 (1996)
- Belyavskiy M; Westerman M; DiMichele L; Wilson VG
Perturbation of the host cell cycle and DNA replication by the bovine papillomavirus replication protein E1.
Virology 219: 206-19 (1996)
- Bollen LJ; Tjong-A-Hung SP; van der Velden J; Mol BW; Lammes FB; ten Kate FW; ter Schegget J; Bleker OP
Human papillomavirus DNA after treatment of cervical dysplasia: low prevalence in normal cytologic smears.
Cancer 77: 2538-43 (1996)
- Bristow RG; Brail L; Jang A; Peacock J; Chung S; Benchimol S; Hill RP
P53-mediated radioresistance does not correlate with metastatic potential in tumorigenic rat embryo cell lines following oncogene transfection.
Int J Radiat Oncol Biol Phys 34: 341-55 (1996)
- Brokaw JL; Blanco M; McBride AA
Amino acids critical for the functions of the bovine papillomavirus type 1 E2 transactivator.
J Virol 70: 23-9 (1996)
- Chan MK; Lau KM; Tsui Y; Wong FW; Huang DP
Human papillomavirus infection in Hong Kong Chinese women with normal and abnormal cervix—detection by polymerase chain reaction method on cervical scrapes.
Gynecol Oncol 60: 217-23 (1996)
- Chen SL; Tsai TZ; Han CP; Tsao YP
Mutational analysis of human papillomavirus type 11 E5a oncoprotein.
J Virol 70: 3502-8 (1996)
- Chiba I; Shindoh M; Yasuda M; Yamazaki Y; Amemiya A; Sato Y; Fujinaga K; Notani K; Fukuda H
Mutations in the p53 gene and human papillomavirus infection as significant prognostic factors in squamous cell carcinomas of the oral cavity.
Oncogene 12: 1663-8 (1996)
- Chipev CC; Steinert PM; Woodworth CD
Characterization of an immortalized cell line from a patient with epidermolytic hyperkeratosis.
J Invest Dermatol 106: 385-90 (1996)
- Chou CY; Chen YH; Tzeng CC; Cheng YC; Chang CF; Chen TM
Establishment and characterization of a human-papillomavirus negative, p53-mutation negative human cervical cancer cell line.
Cancer Lett 102: 173-81 (1996)
- Crook T; Ludwig RL; Marston NJ; Willkomm D; Vousden KH
Sensitivity of p53 lysine mutants to ubiquitin-directed degradation targeted by human papillomavirus E6.
Virology 217: 285-92 (1996)
- Dalal S; Gao Q; Androphy EJ; Band V
Mutational analysis of human papillomavirus type 16 E6 demonstrates that p53 degradation is necessary for immortalization of mammary epithelial cells.
J Virol 70: 683-8 (1996)
- Dall P; Hekele A; Ikenberg H; Goppinger A; Bauknecht T; Pfeleiderer A; Moll J; Hofmann M; Ponta H; Herrlich P
Increasing incidence of CD44v7/8 epitope expression during uterine cervical carcinogenesis.
Int J Cancer 69: 79-85 (1996)
- Donnelly JJ; Martinez D; Jansen KU; Ellis RW; Montgomery DL; Liu MA
Protection against papillomavirus with a polynucleotide vaccine.
J Infect Dis 173: 314-20 (1996)
- Doorbar J; Medcalf E; Napthine S
Analysis of HPV1 E4 complexes and their association with keratins in vivo.
Virology 218: 114-26 (1996)

References (1996)

- Evans C; Bauer S; Grubert T; Brucker C; Baur S; Heeg K; Wagner H; Lipford GB
HLA-A2-restricted peripheral blood cytolytic T lymphocyte response to HPV type 16 proteins E6 and E7 from patients with neoplastic cervical lesions.
Cancer Immunol Immunother 42: 151-60 (1996)
- Fibi MR; Kunz J; Weimer T; Schulz R; Teifke P; Grzeschik KH; Johannsen R; Zettlmeissl G
Status and transcriptional activity of a bovine papillomavirus-I-based expression vector in a recombinant production cell lines.
Biotechnol Appl Biochem 23 (Pt 2): 187 (1996)
- Foster SA; Galloway DA
Human papillomavirus type 16 E7 alleviates a proliferation block in early passage human mammary epithelial cells.
Oncogene 12: 1773-9 (1996)
- Frattini MG; Lim HB; Laimins LA
In vitro synthesis of oncogenic human papillomaviruses requires episomal genomes for differentiation-dependent late expression.
Proc Natl Acad Sci U S A 93: 3062-7 (1996)
- Fujita M; Enomoto T; Wada H; Inoue M; Okudaira Y; Shroyer KR
Application of clonal analysis. Differential diagnosis for synchronous primary ovarian and endometrial cancers and metastatic cancer.
Am J Clin Pathol 105: 350-9 (1996)
- Ghim SJ; Young R; Jenson AB
Antigenicity of bovine papillomavirus type 1 (BPV-1) L1 virus-like particles compared with that of intact BPV-1 virions.
J Gen Virol 77 (Pt 2): 183-8 (1996)
- Gjoen K; Olsen AO; Magnus P; Grinde B; Sauer T; Orstavik I
Prevalence of human papillomavirus in cervical scrapes, as analyzed by PCR, in a population-based sample of women with and without cervical dysplasia.
APMIS 104: 68-74 (1996)
- Grassmann K; Rapp B; Maschek H; Petry KU; Iftner T
Identification of a differentiation-inducible promoter in the E7 open reading frame of human papillomavirus type 16 (HPV-16) in raft cultures of a new cell line containing high copy numbers of episomal HPV-16 DNA.
J Virol 70: 2339-49 (1996)
- Grinstein E; Weinert I; Droese B; Pagano M; Royer HD
Cell cycle regulation of nuclear factor p32 DNA-binding activity by novel phase-specific inhibitors.
J Biol Chem 271: 9215-22 (1996)
- Grossel MJ; Barsoum J; Prakash SS; Androphy EJ
The BPV-1 E2 DNA-contact helix cysteine is required for transcriptional activation but not replication in mammalian cells.
Virology 217: 301-10 (1996)
- Harry JB; Wettstein FO
Transforming properties of the cottontail rabbit papillomavirus oncoproteins Le6 and SE6 and of the E8 protein.
J Virol 70: 3355-62 (1996)
- Hawkins DS; Demers GW; Galloway DA
Inactivation of p53 enhances sensitivity to multiple chemotherapeutic agents.
Cancer Res 56: 892-8 (1996)
- Hembree JR; Agarwal C; Beard RL; Chandraratna RA; Eckert R
Retinoid X receptor-specific retinoids inhibit the ability of retinoic acid receptor-specific retinoids to increase the level of insulin-like growth factor binding protein-3 in human ectocervical epithelial cells.
Cancer Res 56: 1794-9 (1996)
- Herber R; Liem A; Pitot H; Lambert PF
Squamous epithelial hyperplasia and carcinoma in mice transgenic for the human papillomavirus type 16 E7 oncogene.
J Virol 70: 1873-81 (1996)
- Hofmann KJ; Neepser MP; Markus HZ; Brown DR; Muller M; Jansen KU
Sequence conservation within the major capsid protein of human papillomavirus (HPV) type 18 and formation of HPV-18 virus-like particles in *Saccharomyces cerevisiae*.
J Gen Virol 77 (Pt 3): 465-8 (1996)
- Hoos A; D'Incan C; Gissmann L; Altmann A; Momburg F; Nindl I; Osen W; Schonning BH; Jochmus I
Human papillomavirus type 16 (HPV 16) E7 and major histocompatibility complex (MHC) class I and II expression in human keratinocytes in culture.
Arch Virol 141: 449-58 (1996)
- Hwang ES; Naeger LK; DiMaio D
Activation of the endogenous p53 growth inhibitory pathway in HeLa cervical carcinoma cells by expression of the bovine papillomavirus E2 gene.
Oncogene 12: 795-803 (1996)
- Isacson C; Kesis TD; Hedrick L; Cho KR
Both cell proliferation and apoptosis increase with lesion grade in cervical neoplasia but do not correlate with human papillomavirus type.
Cancer Res 56: 669-74 (1996)

- Jenkins A; Kristiansen E; Ask E; Servold T; Christensen A; Kristiansen BE
Human papillomavirus infection in cervical biopsies from Norwegian gynecological in-patients.
APMIS 104: 30-4 (1996)
- Karlen S; Offord EA; Beard P
Functional promoters in the genome of human papillomavirus type 6b.
J Gen Virol 77 (Pt 1): 11-6 (1996)
- Khleif SN; DeGregori J; Yee CL; Otterson GA; Kaye FJ; Nevins JR; Howley PM
Inhibition of cyclin D-CDK4/CDK6 activity is associated with an E2F-mediated induction of cyclin kinase inhibitor activity.
Proc Natl Acad Sci U S A 93: 4350-4 (1996)
- Kim YT; Thomas NF; Kessis TD; Wilkinson EJ; Hedrick L; Cho KR
p53 mutations and clonality in vulvar carcinomas and squamous hyperplasias: evidence suggesting that squamous hyperplasias do not serve as direct precursors of human papillomavirus-negative vulvar carcinomas.
Hum Pathol 27: 389-95 (1996)
- Kitagawa K; Yoshikawa H; Onda T; Kawana T; Taketani Y; Yoshikura H; Iwamoto A
Genomic organization of human papillomavirus type 18 in cervical cancer specimens.
Jpn J Cancer Res 87: 263-8 (1996)
- Klingelutz AJ; Foster SA; McDougall JK
Telomerase activation by the E6 gene product of human papillomavirus type 16.
Nature 380: 79-82 (1996)
- Kowalski LA; Tsang SS; Davison AJ
Arsenic and chromium enhance transformation of bovine papillomavirus DNA-transfected C3H/10T1/2 cells.
Cancer Lett 103: 65-9 (1996)
- Lai YM; Yang FP; Pao CC
Human papillomavirus deoxyribonucleic acid and ribonucleic acid in seminal plasma and sperm cells.
Fertil Steril 65: 1026-30 (1996)
- Larson AA; Kern S; Sommers RL; Yokota J; Cavenee WK; Hampton GM
Analysis of replication error (RER+) phenotypes in cervical carcinoma.
Cancer Res 56: 1426-31 (1996)
- Li X; Coffino P
Identification of a region of p53 that confers lability.
J Biol Chem 271: 4447-51 (1996)
- Lizard G; Chignol MC; Chardonnet Y; Schmitt D
Differences of reactivity to interferon gamma in HeLa and CaSki cells: a combined immunocytochemical and flow-cytometric study.
J Cancer Res Clin Oncol 122: 223-30 (1996)
- Lopez-Beltran A; Escudero AL; Vicioso L; Munoz E; Carrasco JC
Human papillomavirus DNA as a factor determining the survival of bladder cancer patients.
Br J Cancer 73: 124-7 (1996)
- Maki H; Fujikawa-Adachi K; Yoshie O
Evidence for a promoter-like activity in the short non-coding region of human papillomaviruses.
J Gen Virol 77 (Pt 3): 453-8 (1996)
- Mao EJ; Schwartz SM; Daling JR; Oda D; Tickman L; Beckmann AM
Human papilloma viruses and p53 mutations in normal pre-malignant and malignant oral epithelia.
Int J Cancer 69: 152-8 (1996)
- Marionnet AV; Lizard G; Chignol MC; Chardonnet Y; Schmitt D
Differences in reactivity to cyclosporin A and interferon-gamma of normal and HPV-transformed keratinocytes.
Anticancer Res 16: 257-67 (1996)
- Massimi P; Pim D; Storey A; Banks L
HPV-16 E7 and adenovirus E1a complex formation with TATA box binding protein is enhanced by casein kinase II phosphorylation.
Oncogene 12: 2325-30 (1996)
- McIntyre MC; Ruesch MN; Laimins LA
Human papillomavirus E7 oncoproteins bind a single form of cyclin E in a complex with cdk2 and p107.
Virology 215: 73-82 (1996)
- Medina-Martinez O; Morales-Peza N; Yaniv M; Garcia-Carranca A; Thierry F
A single element mediates glucocorticoid hormone response of HPV18 with no functional interactions with AP1 or hbrm.
Virology 217: 392-6 (1996)
- Mitchell MF; Hamada K; Sastry KJ; Sarkar A; Tortolero-Luna G; Wharton JT; Roth JA
Transgene expression in the rhesus cervix mediated by an adenovirus expressing beta-galactosidase.
Am J Obstet Gynecol 174: 1094-101 (1996)
- Monini P; Rotola A; de Lellis L; Corallini A; Secchiero P; Albini A; Benelli R; Parravicini C; Barbanti-Brodano G; Cassai E
Latent BK virus infection and Kaposi's sarcoma pathogenesis.
Int J Cancer 66: 717-22 (1996)

References (1996)

- Muller F; Sapp M
Domains of the E1 protein of human papillomavirus type 33 involved in binding to the E2 protein.
Virology 219: 247-56 (1996)
- Mulloikandov MR; Kholodilov NG; Atkin NB; Burk RD; Johnson AB; Klinger HP
Genomic alterations in cervical carcinoma: losses of chromosome heterozygosity and human papilloma virus tumor status.
Cancer Res 56: 197-205 (1996)
- Nagano H; Yoshikawa H; Kawana T; Yokota H; Taketani Y; Igarashi H; Yoshikura H; Iwamoto A
Association of multiple human papillomavirus types with vulvar neoplasias.
J Obstet Gynaecol 22: 1-8 (1996)
- Naghashfar Z; DiPaolo JA; Woodworth CD; Passaniti A
Immortalization of human adult prostatic adenocarcinoma cells by human papilloma virus HPV16 and -18 DNA.
Cancer Lett 100: 47-54 (1996)
- Nilsson CH; Bakos E; Petry KU; Schneider A; Durst M
Promoter usage in the E7 ORF of HPV16 correlates with epithelial differentiation and is largely confined to low-grade genital neoplasia.
Int J Cancer 65: 6-12 (1996)
- Nuber U; Schwarz S; Kaiser P; Schneider R; Scheffner M
Cloning of human ubiquitin-conjugating enzymes UbcH6 and UbcH7 (E2-F1) and characterization of their interaction with E6-AP and RSP5.
J Biol Chem 271: 2795-800 (1996)
- Pan J; Nakanishi K; Yutsudo M; Inoue H; Li Q; Oka K; Yoshioka N; Hakura A
Isolation of a novel gene down-regulated by v-src.
FEBS Lett 383: 21-5 (1996)
- Pao CC; Yao DS; Lin CY; Lee SC; Su HT; Lin SC
Genomic aberrations of human papillomavirus recovered from cervical cancers.
Biochem Biophys Res Commun 222: 116-20 (1996)
- Park TW; Richart RM; Sun XW; Wright TC Jr
Association between human papillomavirus type and clonal status of cervical squamous intraepithelial lesions [see comments]
J Natl Cancer Inst 88: 355-8 (1996)
- Piirsoo M; Ustav E; Mandel T; Stenlund A; Ustav M
Cis and trans requirements for stable episomal maintenance of the BPV-1 replicator.
EMBO J 15: 1-11 (1996)
- Pusch O; Soucek T; Wawra E; Hengstschlager-Ottner E; Bernaschek G; Hengstschlager M
Specific transformation abolishes cyclin D1 fluctuation throughout the cell cycle.
FEBS Lett 385: 143-8 (1996)
- Qi YM; Peng SW; Hengst K; Evander M; Park DS; Zhou J; Frazer IH
Epithelial cells display separate receptors for papillomavirus VLPs and for soluble L1 capsid protein.
Virology 216: 35-45 (1996)
- Rasmussen M; Nielsen PE; Manoharan M; Buchardt O; Koch C
Specificity of antibodies raised against a specific phosphoromonothioate oligonucleotide sequence.
J Immunol Methods 189: 47-58 (1996)
- Ratiarson A
Analysis of the papillomavirus type 16 DNA amplification by PCR in SiHa and SLT cell lines.
Microbios 85: 127-31 (1996)
- Ressing ME; van Driel WJ; Celis E; Sette A; Brandt MP; Hartman M; Anholts JD; Schreuder GM; ter Harmsel WB; Fleuren GJ; Trimbos BJ; Kast WM; Melief CJ
Occasional memory cytotoxic T-cell responses of patients with human papillomavirus type 16-positive cervical lesions against a human leukocyte antigen-A *0201-restricted E7-encoded epitope.
Cancer Res 56: 582-8 (1996)
- Rowan, S; Ludwig RL; Haupt Y; Bates S; Lu X; Oren M; Vousden KH
Specific loss of apoptotic but not cell-cycle arrest function in a human tumor derived p53 mutant.
EMBO J 15: 827-38 (1996)
- Sagerman PM; Choi YJ; Hu Y; Niedt GW
Human papilloma virus, vulvar dystrophy, and vulvar carcinoma: differential expression of human papillomavirus and vulvar dystrophy in the presence and absence of squamous cell carcinoma of the vulva.
Gynecol Oncol 61: 328-32 (1996)
- Sakai H; Yasugi T; Benson JD; Dowhanick JJ; Howley PM
Targeted mutagenesis of the human papillomavirus type 16 E2 transactivation domain reveals separable transcriptional activation and DNA replication functions.
J Virol 70: 1602-11 (1996)
- Sandler AB; Baker CC; Spalholz BA
Sp1 is critical for basal and E2-transactivated transcription from the bovine papillomavirus type 1 P89 promoter.
J Gen Virol 77 (Pt 2): 189-98 (1996)

- Sarma D; Yang X; Jin G; Shindoh M; Pater MM; Pater A
Resistance to retinoic acid and altered cytokeratin expression of human papillomavirus type 16-immortalized endocervical cells after tumorigenesis.
Int J Cancer 65: 345-50 (1996)
- Shamanin V; zur Hausen H; Lavergne D; Proby CM; Leigh IM; Neumann C; Hamm H; Goos M; Hausteiner UF; Jung EG; Plewig G; Wolff H; de Villiers EM
Human papillomavirus infections in nonmelanoma skin cancers from renal transplant recipients and nonimmunosuppressed patients [see comments]
J Natl Cancer Inst 88: 802-11 (1996)
- Shen MR; Wu SN; Chou CY
Volume-sensitive chloride channels in the primary culture cells of human cervical carcinoma.
Biochim Biophys Acta 1315: 138-44 (1996)
- Shepherd PS; Rowe AJ; Cridland JC; Coletart T; Wilson P; Luxton JC
Proliferative T cell responses to human papillomavirus type 16 L1 peptides in patients with cervical dysplasia.
J Gen Virol 77 (Pt 4): 593-602 (1996)
- Sherman L; Schlegel R
Serum- and calcium-induced differentiation of human keratinocytes is inhibited by the E6 oncoprotein of human papillomavirus type 16.
J Virol 70: 3269-79 (1996)
- Shin KH; Tannyhill RJ; Liu X; Park NH
Oncogenic transformation of HPV-immortalized human oral keratinocytes is associated with the genetic instability of cells.
Oncogene 12: 1089-96 (1996)
- Shindoh M; Kimura S; Yamashita T; Fujinaga K; Amemiya A
[Histopathological application of in situ PCR (in situ hybridization after amplification by PCR)]
Tanpakushitsu Kakusan Koso 41: 660-3 (1996)
- Skiadopoulos MH; McBride AA
The bovine papillomavirus type 1 E2 transactivator and repressor proteins use different nuclear localization signals.
J Virol 70: 1117-24 (1996)
- Snijders PJ; Scholes AG; Hart CA; Jones AS; Vaughan ED; Woolgar JA; Meijer CJ; Walboomers JM; Field JK
Prevalence of mucosotropic human papillomaviruses in squamous-cell carcinoma of the head and neck.
Int J Cancer 66: 464-9 (1996)
- Soini Y; Pollanen R; Kempainen S; Paakko P; Lehto VP
The association of vascular proliferation with HPV status and epithelial PCNA positivity in cervical intraepithelial lesions.
APMIS 104: 183-90 (1996)
- Sparkowski J; Mense M; Anders J; Schlegel R
E5 oncoprotein transmembrane mutants dissociate fibroblast transforming activity from 16-kilodalton protein binding and platelet-derived growth factor receptor binding and phosphorylation.
J Virol 70: 2420-30 (1996)
- Stewart AC; Eriksson AM; Manos MM; Munoz N; Bosch FX; Peto J; Wheeler CM
Intratypic variation in 12 human papillomavirus types: a worldwide perspective.
J Virol 70: 3127-36 (1996)
- Stoppler H; Kirchoff T; Kartenbeck J; Gissmann L; Alonso A
Influence of human papillomavirus type 16 gene expression on in vitro differentiation of the human teratocarcinoma cell line 2102Ep.
Mol Carcinog 16: 109-14 (1996)
- Stoppler H; Stoppler MC; Adduci A; Koval D; Schlegel R
The serine protease inhibitors TLCK and TPCK react with the RB-binding core of HPV-18 E7 protein and abolish its RB-binding capability.
Virology 217: 542-53 (1996)
- Stubenrauch F; Leigh IM; Pfister H
E2 represses the late gene promoter of human papillomavirus type 8 at high concentrations by interfering with cellular factors.
J Virol 70: 119-26 (1996)
- Sun YN; Lu JZ; McCance DJ
Mapping of HPV-11 E1 binding site and determination of other important cis elements for replication of the origin.
Virology 216: 219-22 (1996)
- Suzuki H; Komiya A; Aida S; Ito H; Yatani R; Shimazaki J
Detection of human papillomavirus DNA and p53 gene mutations in human prostate cancer.
Prostate 28: 318-24 (1996)
- Taja-Chayeb L; Salas-Garcia M; Salcedo-Vargas M
[Molecular bases of papillomavirus and polyomavirus carcinogenesis]
Salud Publica Mex 38: 47-57 (1996)

References (1996)

- van der Burg SH; Visseren MJ; Brandt RM; Kast WM; Melief CJ
Immunogenicity of peptides bound to MHC class I molecules depends on the MHC-peptide complex stability.
J Immunol 156: 3308-14 (1996) Vieira KB; Goldstein DJ; Villa LL
Tumor necrosis factor alpha interferes with the cell cycle of normal and papillomavirus-immortalized human keratinocytes.
Cancer Res 56: 2452-7 (1996)
- Volter C; He Y; Delius H; Roy-Burman A; Greenspan JS; Greenspan D; de Villiers EM
Novel HPV types present in oral papillomatous lesions from patients with HIV infection.
Int J Cancer 66: 453-6 (1996)
- Wahl AF; Donaldson KL; Fairchild C; Lee FY; Foster SA; Demers GW; Galloway DA
Loss of normal p53 function confers sensitization to Taxol by increasing G2/M arrest and apoptosis.
Nat Med 2: 72-9 (1996)
- Willis SD; Seyfred MA
Pituitary-specific chromatin structure of the rat prolactin distal enhancer element.
Nucleic Acids Res 24: 1065-72 (1996)
- Wong HK; Ziff EB
The human papillomavirus type 16 E7 protein complements adenovirus type 5 E1A amino-terminus-dependent transactivation of adenovirus type 5 early genes and increases ATF and Oct-1 DNA binding activity.
J Virol 70: 332-40 (1996)
- Xiong Y; Kuppuswamy D; Li Y; Livanos EM; Hixon M; White A; Beach D; Tlsty TD
Alteration of cell cycle kinase complexes in human papillomavirus E6- and E7-expressing fibroblasts precedes neoplastic transformation.
J Virol 70: 999-1008 (1996)
- Yamashita T; Inoue Y; Fujinaga Y; Fujinaga K
[Detection and typing of HPV genome: consensus PCR method]
Tanpakushitsu Kakusan Koso 41: 664-8 (1996)
- Yang X; Jin G; Nakao Y; Rahimtula M; Pater MM; Pater A
Malignant transformation of HPV 16-immortalized human endocervical cells by cigarette smoke condensate and characterization of multistage carcinogenesis.
Int J Cancer 65: 338-44 (1996)
- Yang X; Nakao Y; Pater MM; Pater A
Identification of two novel cellular genes associated with multistage carcinogenesis of human endocervical cells by mRNA differential display.
Carcinogenesis 17: 563-7 (1996)
- Yukawa K; Butz K; Yasui T; Kikutani H; Hoppe-Seyler F
Regulation of human papillomavirus transcription by the differentiation-dependent epithelial factor Epc-1/skn-1a.
J Virol 70: 10-6 (1996)
- Zhang L; Kashanchi F; Zhan Q; Zhan S; Brady JN; Fornace AJ; Seth P; Helman LJ
Regulation of insulin-like growth factor II P3 promoter by p53: a potential mechanism for tumorigenesis.
Cancer Res 56: 1367-73 (1996)
- Zhao C; Tan W; Sokolowski M; Schwartz S
Identification of nuclear and cytoplasmic proteins that interact specifically with an AU-rich, cis-acting inhibitory sequence in the 3' untranslated region of human papillomavirus type 1 late mRNAs.
J Virol 70: 3659-67 (1996)
- Zou S; Brown EG
Stable expression of the reovirus mu2 protein in mouse L cells complements the growth of a reovirus ts mutant with a defect in its M1 gene.
Virology 217: 42-8 (1996)
- Zwerschke W; Joswig S; Jansen-Durr P
Identification of domains required for transcriptional activation and protein dimerization in the human papillomavirus type-16 E7 protein.
Oncogene 12: 213-20 (1996)