

- Abroi A; Kurg R; Ustav M  
Transcriptional and replicational activation functions in the bovine papillomavirus type 1 E2 protein are encoded by different structural determinants.  
*J Virol* 70: 6169-79 (1996)
- Adachi A; Kiyono T; Hayashi Y; Ohashi M; Ishibashi M  
Detection of human papillomavirus (HPV) type 47 DNA in malignant lesions from epidermodysplasia verruciformis by protocols for precise typing of related HPV DNAs.  
*J Clin Microbiol* 34: 369-75 (1996)
- Adams V; Moll C; Schmid M; Rodrigues C; Moos R; Briner J  
Detection and typing of human papillomavirus in biopsy and cytological specimens by polymerase chain reaction and restriction enzyme analysis: a method suitable for semiautomation.  
*J Med Virol* 48: 161-70 (1996)
- Agarwal C; Chandraratna RA; Teng M; Nagpal S; Rorke EA; Eckert RL  
Differential regulation of human ectocervical epithelial cell line proliferation and differentiation by retinoid X receptor- and retinoic acid receptor-specific retinoids.  
*Cell Growth Differ* 7: 521-30 (1996)
- Akutsu N; Shirasawa H; Asano T; Isono K; Simizu B  
p53-Dependent and -independent transactivation by the E6 protein of human papillomavirus type 16.  
*J Gen Virol* 77 ( Pt 3): 459-63 (1996)
- Alexander KA; Phelps WC  
A fluorescence anisotropy study of DNA binding by HPV-11 E2C protein: a hierarchy of E2-binding sites.  
*Biochemistry* 35: 9864-72 (1996)
- Alexander M; Salgaller ML; Celis E; Sette A; Barnes WA; Rosenberg SA; Steller MA  
Generation of tumor-specific cytolytic T lymphocytes from peripheral blood of cervical cancer patients by in vitro stimulation with a synthetic human papillomavirus type 16 E7 epitope.  
*Am J Obstet Gynecol* 175: 1586-93 (1996)
- Alvarez Alvarez I; Sanchez Lazo P; Ramos Gonzalez S; Rodrigo Tapia JP; Llorente Pendas JL; Suarez Nieto C  
Simultaneous screening of HPV-6b and 16 in pharyngolaryngeal cancer]  
*Acta Otorrinolaringol Esp* 47: 93-6 (1996)
- Antinore MJ; Birrer MJ; Patel D; Nader L; McCance DJ  
The human papillomavirus type 16 E7 gene product interacts with and trans-activates the AP1 family of transcription factors.  
*EMBO J* 15: 1950-60 (1996)
- Anwar K; Nakakuki K; Imai H; Shiraishi T; Inuzuka M  
Infection of human papillomavirus (HPV) and p53 over-expression in human female genital tract carcinoma.  
*JPMA J Pak Med Assoc* 46: 220-4 (1996)
- Apt D; Watts RM; Suske G; Bernard HU  
High Sp1/Sp3 ratios in epithelial cells during epithelial differentiation and cellular transformation correlate with the activation of the HPV-16 promoter.  
*Virology* 224: 281-91 (1996)
- Arany I; Brysk MM; Brysk H; Tyring SK  
Response to interferon treatment decreases with epidermal dedifferentiation in condylomas.  
*Antiviral Res* 32: 19-26 (1996)
- Arany I; Tyring SK  
Status of local cellular immunity in interferon-responsive and -nonresponsive human papillomavirus-associated lesions.  
*Sex Transm Dis* 23: 475-80 (1996)
- Arbeit JM; Howley PM; Hanahan D  
Chronic estrogen-induced cervical and vaginal squamous carcinogenesis in human papillomavirus type 16 transgenic mice.  
*Proc Natl Acad Sci U S A* 93: 2930-5 (1996)
- Arbeit JM; Olson DC; Hanahan D  
Upregulation of fibroblast growth factors and their receptors during multi-stage epidermal carcinogenesis in K14-HPV16 transgenic mice.  
*Oncogene* 13: 1847-57 (1996)
- Auborn KJ; Wang H; Vaccariello MA; Taichman LB  
Kinetics of HPV11 DNA replication after infection of keratinocytes with virions.  
*Virus Res* 43: 85-90 (1996)
- Baay MF; Quint WG; Koudstaal J; Hollema H; Duk JM; Burger MP; Stolz E; Herbrink P  
Comprehensive study of several general and type-specific primer pairs for detection of human papillomavirus DNA by PCR in paraffin-embedded cervical carcinomas.  
*J Clin Microbiol* 34: 745-7 (1996)
- Bagot M; Charue D; Cerni C; Revuz J; Meneguzzi G  
Induction of rat CD4+ proliferative and mouse CD8+ cytotoxic T-cell lines specific for human papillomavirus type 16 antigens.  
*Res Virol* 147: 301-11 (1996)
- Barbosa MS  
The oncogenic role of human papillomavirus proteins.  
*Crit Rev Oncog* 7: 1-18 (1996)

## References (1996)

- Bauer-Hofmann R; Borghouts C; Auvinen E; Bourda E; Rosl F; Alonso A  
Genomic cloning and characterization of the nonoccupied allele corresponding to the integration site of human papillomavirus type 16 DNA in the cervical cancer cell line SiHa.  
*Virology* 217: 33-41 (1996)
- Bauknecht T; See RH; Shi Y  
A novel C/EBP beta-YY1 complex controls the cell-type-specific activity of the human papillomavirus type 18 upstream regulatory region.  
*J Virol* 70: 7695-705 (1996)
- Behbakht K; DeGeest K; Turyk ME; Wilbanks GD  
All-trans-retinoic acid inhibits the proliferation of cell lines derived from human cervical neoplasia.  
*Gynecol Oncol* 61: 31-9 (1996)
- Belyavskiy M; Westerman M; DiMichele L; Wilson VG  
Perturbation of the host cell cycle and DNA replication by the bovine papillomavirus replication protein E1.  
*Virology* 219: 206-19 (1996)
- Benincasa D; Silberklang M; Mark GE 3rd; Ludmerer SW  
Rapid, high-level transient expression of papillomavirus-like particles in insect cells.  
*Biotechniques* 20: 890-5 (1996)
- Bollen LJ; Tjong-A-Hung SP; van der Velden J; Mol BW; Lammes FB; ten Kate FW; ter Schegget J; Bleker OP  
Human papillomavirus DNA after treatment of cervical dysplasia: low prevalence in normal cytologic smears.  
*Cancer* 77: 2538-43 (1996)
- Bornstein J; Shapiro S; Rahat M; Goldshmid N; Goldik Z; Abramovici H; Lahat N  
Polymerase chain reaction search for viral etiology of vulvar vestibulitis syndrome.  
*Am J Obstet Gynecol* 175: 139-44 (1996)
- Boylan MO; Athanassiou M; Houle B; Wang Y; Zarbl H  
Activation of tumor suppressor genes in nontumorigenic revertants of the HeLa cervical carcinoma cell line.  
*Cell Growth Differ* 7: 725-35 (1996)
- Breiding DE; Grossel MJ; Androphy EJ  
Genetic analysis of the bovine papillomavirus E2 transcriptional activation domain.  
*Virology* 221: 34-43 (1996)
- Bristow RG; Brail L; Jang A; Peacock J; Chung S; Benchimol S; Hill RP  
P53-mediated radioresistance does not correlate with metastatic potential in tumorigenic rat embryo cell lines following oncogene transfection.  
*Int J Radiat Oncol Biol Phys* 34: 341-55 (1996)
- Brokaw JL; Blanco M; McBride AA  
Amino acids critical for the functions of the bovine papillomavirus type 1 E2 transactivator.  
*J Virol* 70: 23-9 (1996)
- Brown DR; Pratt L; Bryan JT; Fife KH; Jansen K  
Virus-like particles and E1E4 protein expressed from the human papillomavirus type 11 bicistronic E1E4L1 transcript.  
*Virology* 222: 43-50 (1996)
- Bucher K; Szalai G; Marti E; Griot-Wenk ME; Lazary S; Pauli U  
Tumour suppressor gene p53 in the horse: identification, cloning, sequencing and a possible role in the pathogenesis of equine sarcoid.  
*Res Vet Sci* 61: 114-9 (1996)
- Butz K; Geisen C; Ullmann A; Spitkovsky D; Hoppe-Seyler F  
Cellular responses of HPV-positive cancer cells to genotoxic anti-cancer agents: repression of E6/E7-oncogene expression and induction of apoptosis.  
*Int J Cancer* 68: 506-13 (1996)
- Carmody MW; Jones M; Tarraza H; Vary CP  
Use of the polymerase chain reaction to specifically amplify integrated HPV-16 DNA by virtue of its linkage to interspersed repetitive DNA.  
*Mol Cell Probes* 10: 107-16 (1996)
- Cayrol C; Flemington EK  
The Epstein-Barr virus bZIP transcription factor Zta causes G0/G1 cell cycle arrest through induction of cyclin-dependent kinase inhibitors.  
*EMBO J* 15: 2748-59 (1996)
- Chan MK; Lau KM; Tsui Y; Wong FW; Huang DP  
Human papillomavirus infection in Hong Kong Chinese women with normal and abnormal cervix—detection by polymerase chain reaction method on cervical scrapes.  
*Gynecol Oncol* 60: 217-23 (1996)
- Chan PJ; Seraj IM; Kalugdan TH; King A  
Evidence for ease of transmission of human papillomavirus DNA from sperm to cells of the uterus and embryo.  
*J Assist Reprod Genet* 13: 516-9 (1996)
- Chang D; Tsai RT; Wang M; Ou WC  
Different genotypes of human polyomaviruses found in patients with autoimmune diseases in Taiwan.  
*J Med Virol* 48: 204-9 (1996)
- Chang T; Khalsa O; Wang H; Lee ME; Schlegel R  
Staurosporine resistance accompanies DNA tumor virus-induced immortalization and is independent of the expression and activities of ERK1, ERK2, cyclin A, cyclin-dependent kinase (cdk) 2, and cdk4.  
*Cell Growth Differ* 7: 361-72 (1996)

- Chen SL; Hsieh TB; Tsao YP; Han CP; Yang YF  
Coincidental expression of E5a and c-jun in human papillomavirus type 6/11-infected condylomata.  
*J Gen Virol* 77 ( Pt 6): 1145-9 (1996)
- Chen SL; Huang CH; Tsai TC; Lu KY; Tsao YP  
The regulation mechanism of c-jun and junB by human papillomavirus type 16 E5 oncoprotein.  
*Arch Virol* 141: 791-800 (1996)
- Chen SL; Lin YK; Li LY; Tsao YP; Lo HY; Wang WB; Tsai TC  
E5 proteins of human papillomavirus types 11 and 16 transactivate the c-fos promoter through the NF1 binding element.  
*J Virol* 70: 8558-63 (1996)
- Chen SL; Tsai TZ; Han CP; Tsao YP  
Mutational analysis of human papillomavirus type 11 E5a oncoprotein.  
*J Virol* 70: 3502-8 (1996)
- Chen TM; Chen YH; Wu CC; Chen CA; Chang CF; Hsieh CY  
Factors influencing tumor cell kinetics in cervical cancer.  
*J Cancer Res Clin Oncol* 122: 504-8 (1996)
- Chen YH; Huang LH; Chen TM  
Differential effects of progestins and estrogens on long control regions of human papillomavirus types 16 and 18.  
*Biochem Biophys Res Commun* 224: 651-9 (1996)
- Chen Z; Kamath P; Zhang S; St. John L; Adler-Storthz K; Shillitoe EJ  
Effects on tumor cells of ribozymes that cleave the RNA transcripts of human papillomavirus type 18.  
*Cancer Gene Ther* 3: 18-23 (1996)
- Chiang LC; Chiang W; Chang LL; Wu WJ; Huang CH  
Characterization of a new human transitional cell carcinoma cell line from the renal pelvis, RTCC-1/KMC.  
*Kao Hsiung I Hsueh Ko Hsueh Tsa Chih* 12: 448-52 (1996)
- Chiba I; Shindoh M; Yasuda M; Yamazaki Y; Amemiya A; Sato Y; Fujinaga K; Notani K; Fukuda H  
Mutations in the p53 gene and human papillomavirus infection as significant prognostic factors in squamous cell carcinomas of the oral cavity.  
*Oncogene* 12: 1663-8 (1996)
- Chinami M; Inoue M; Masunaga K; Fukuma T; Shingu M; Toyoda T  
Nucleic acid binding by zinc finger-like motif of human papillomavirus type 16 E7 oncoprotein.  
*J Virol Methods* 59: 173-6 (1996)
- Chipev CC; Steinert PM; Woodworth CD  
Characterization of an immortalized cell line from a patient with epidermolytic hyperkeratosis.  
*J Invest Dermatol* 106: 385-90 (1996)
- Choo KB; Chen CM; Han CP; Cheng WT; Au LC  
Molecular analysis of cellular loci disrupted by papillomavirus 16 integration in cervical cancer: frequent viral integration in topologically destabilized and transcriptionally active chromosomal regions.  
*J Med Virol* 49: 15-22 (1996)
- Chou CY; Chen YH; Tzeng CC; Cheng YC; Chang CF; Chen TM  
Establishment and characterization of a human-papillomavirus negative, p53-mutation negative human cervical cancer cell line.  
*Cancer Lett* 102: 173-81 (1996)
- Christensen ND; Cladel NM; Reed CA; Budgeon LR; Welsh PA; Patrick SD; Kreider JW  
Laboratory production of infectious stocks of rabbit oral papillomavirus.  
*J Gen Virol* 77 ( Pt 8): 1793-8 (1996)
- Christensen ND; Dillner J; Eklund C; Carter JJ; Wipf GC; Reed CA; Cladel NM; Galloway DA  
Surface conformational and linear epitopes on HPV-16 and HPV-18 L1 virus-like particles as defined by monoclonal antibodies.  
*Virology* 223: 174-84 (1996)
- Christensen ND; Reed CA; Cladel NM; Hall K; Leiserowitz GS  
Monoclonal antibodies to HPV-6 L1 virus-like particles identify conformational and linear neutralizing epitopes on HPV-11 in addition to type-specific epitopes on HPV-6.  
*Virology* 224: 477-86 (1996)
- Coppee F; Gerard AC; Denef JF; Ledent C; Vassart G; Dumont JE; Parmentier M  
Early occurrence of metastatic differentiated thyroid carcinomas in transgenic mice expressing the A2a adenosine receptor gene and the human papillomavirus type 16 E7 oncogene.  
*Oncogene* 13: 1471-82 (1996)
- Coussens LM; Hanahan D; Arbeit JM  
Genetic predisposition and parameters of malignant progression in K14-HPV16 transgenic mice.  
*Am J Pathol* 149: 1899-917 (1996)
- Crook T; Ludwig RL; Marston NJ; Willkomm D; Vousden KH  
Sensitivity of p53 lysine mutants to ubiquitin-directed degradation targeted by human papillomavirus E6.  
*Virology* 217: 285-92 (1996)

## References (1996)

- Da J; Chen L; Hu Y  
Human papillomavirus infection and p53 gene mutation in primary lung cancer] *Chung Hua Chung Liu Tsa Chih* 18: 27-9 (1996)
- Dalal S; Gao Q; Androphy EJ; Band V  
Mutational analysis of human papillomavirus type 16 E6 demonstrates that p53 degradation is necessary for immortalization of mammary epithelial cells. *J Virol* 70: 683-8 (1996)
- Dall P; Hekele A; Ikenberg H; Goppinger A; Bauknecht T; Pfeleiderer A; Moll J; Hofmann M; Ponta H; Herrlich P  
Increasing incidence of CD44v7/8 epitope expression during uterine cervical carcinogenesis. *Int J Cancer* 69: 79-85 (1996)
- Pavelic J; Gall-Troselj K; Bosnar MH; Kardum MM; Pavelic K  
PCR amplification of DNA from archival specimens. A methodological approach. *Neoplasma* 43: 75-81 (1996)
- Dellas A; Schultheiss E; Almendral AC; Torhost J; Gudat F  
Expression of CD44 and variant isoforms in cervical intraepithelial neoplasia. *Gynecol Oncol* 62: 218-25 (1996)
- Dellas A; Schultheiss E; Oberholzer M; Torhorst J; Gudat F  
Analysis of proliferative activity using Ki-67 on cervical precancerous lesions and the relationship to p53 expression. *Anticancer Res* 16: 3403-7 (1996)
- Dellas A; Schultheiss E; Oberholzer M; Torhorst J; Gudat F  
Analysis of proliferative activity using Ki-67 on cervical precancerous lesions and the relationship to p53 expression. *Anticancer Res* 16: 3403-7 (1996)
- Demers GW; Espling E; Harry JB; Etscheid BG; Galloway DA  
Abrogation of growth arrest signals by human papillomavirus type 16 E7 is mediated by sequences required for transformation. *J Virol* 70: 6862-9 (1996)
- de Gruijl TD; Bontkes HJ; Stukart MJ; Walboomers JM; Remmink AJ; Verheijen RH; Helmerhorst TJ; Meijer CJ; Scheper RJ  
T cell proliferative responses against human papillomavirus type 16 E7 oncoprotein are most prominent in cervical intraepithelial neoplasia patients with a persistent viral infection. *J Gen Virol* 77 ( Pt 9): 2183-91 (1996)
- Dias Neto E; Rahal P; de Moura RP; Caballero O; Villa LL; Simpson AJ  
DNA ligation quality control by competitive PCR. *Trends Genet* 12: 341-2 (1996)
- DiPaolo JA; Popescu NC; Woodworth CD; Zimonjic DB  
Papillomaviruses and potential copathogens. *Toxicol Lett* 88: 1-7 (1996)
- Dolcetti R; Maestro R  
Possible role of viral agents in the pathogenesis of squamous cell carcinomas of the head and neck] *Acta Otorhinolaryngol Ital* 16: 11-6 (1996)
- Dolcetti R; Maestro R  
Possible role of viral agents in the pathogenesis of squamous cell carcinomas of the head and neck] *Acta Otorhinolaryngol Ital* 16: 11-6 (1996)
- Doldi N; Origoni M; Bassan M; Ferrari D; Rossi M; Ferrari A  
Vascular endothelial growth factor. Expression in human vulvar neoplastic and nonneoplastic tissues. *J Reprod Med* 41: 844-8 (1996)
- Donnelly JJ; Martinez D; Jansen KU; Ellis RW; Montgomery DL; Liu MA  
Protection against papillomavirus with a polynucleotide vaccine. *J Infect Dis* 173: 314-20 (1996)
- Doorbar J; Medcalf E; Napthine S  
Analysis of HPV1 E4 complexes and their association with keratins in vivo. *Virology* 218: 114-26 (1996)
- Duggan-Keen MF; Keating PJ; Stevens FR; Sinnott P; Snijders PJ; Walboomers JM; Davidson S; Hunter RD; Dyer PA; Stern PL  
Immunogenetic factors in HPV-associated cervical cancer: influence on disease progression. *Eur J Immunogenet* 23: 275-84 (1996)
- Evans C; Bauer S; Grubert T; Brucker C; Baur S; Heeg K; Wagner H; Lipford GB  
HLA-A2-restricted peripheral blood cytolytic T lymphocyte response to HPV type 16 proteins E6 and E7 from patients with neoplastic cervical lesions. *Cancer Immunol Immunother* 42: 151-60 (1996)
- Faccini AM; Cairney M; Ashrafi GH; Finbow ME; Campo MS; Pitts JD  
The bovine papillomavirus type 4 E8 protein binds to ductin and causes loss of gap junctional intercellular communication in primary fibroblasts. *J Virol* 70: 9041-5 (1996)
- Ferguson MK; Botchan MR  
Genetic analysis of the activation domain of bovine papillomavirus protein E2: its role in transcription and replication. *J Virol* 70: 4193-9 (1996)

- Fibi MR; Kunz J; Weimer T; Schulz R; Teifke P; Grzeschik KH; Johannsen R; Zettlmeissl G  
Status and transcriptional activity of a bovine-papillomavirus-I-based expression vector in a recombinant production cell line.  
Biotechnol Appl Biochem 23 ( Pt 1): 83-93 (1996)
- Fibi MR; Kunz J; Weimer T; Schulz R; Teifke P; Grzeschik KH; Johannsen R; Zettlmeissl G  
Status and transcriptional activity of a bovine-papillomavirus-I-based expression vector in a recombinant production cell lines.  
Biotechnol Appl Biochem 23 ( Pt 2): 187 (1996)
- Field JK  
Genomic instability in squamous cell carcinoma of the head and neck.  
Anticancer Res 16: 2421-31 (1996)
- Fife KH; Fan L; Fritsch MH; Bryan J; Brown DR  
Association of human papillomavirus type 11 DNA with squamous cell carcinoma of the tongue.  
Arch Otolaryngol Head Neck Surg 122: 1404-8 (1996)
- Fornatora M; Jones AC; Kerpel S; Freedman P  
Human papillomavirus-associated oral epithelial dysplasia (koilocytic dysplasia): an entity of unknown biologic potential.  
Oral Surg Oral Med Oral Pathol Oral Radiol Endod 82: 47-56 (1996)
- Forslund O; Hansson BG  
Human papillomavirus type 70 genome cloned from overlapping PCR products: complete nucleotide sequence and genomic organization.  
J Clin Microbiol 34: 802-9 (1996)
- Foster SA; Galloway DA  
Human papillomavirus type 16 E7 alleviates a proliferation block in early passage human mammary epithelial cells.  
Oncogene 12: 1773-9 (1996)
- Frattini MG; Lim HB; Laimins LA  
In vitro synthesis of oncogenic human papillomaviruses requires episomal genomes for differentiation-dependent late expression.  
Proc Natl Acad Sci U S A 93: 3062-7 (1996)
- Fujita M; Enomoto T; Wada H; Inoue M; Okudaira Y; Shroyer KR  
Application of clonal analysis. Differential diagnosis for synchronous primary ovarian and endometrial cancers and metastatic cancer.  
Am J Clin Pathol 105: 350-9 (1996)
- Funaoka K; Shindoh M; Yamashita T; Fujinaga K; Amemiya A; Totsuka Y  
High-risk HPV-positive human cancer cell lines show different sensitivity to cisplatin-induced apoptosis correlated with the p21 Waf1/Cip1 level.  
Cancer Lett 108: 15-23 (1996)
- Furukawa T; Duguid WP; Rosenberg L; Viallet J; Galloway DA; Tsao MS  
Long-term culture and immortalization of epithelial cells from normal adult human pancreatic ducts transfected by the E6E7 gene of human papilloma virus 16.  
Am J Pathol 148: 1763-70 (1996)
- Furuya H; Yabushita H; Noguchi M; Nakanishi M  
Apoptosis and cell growth fraction in normal, dysplastic and neoplastic squamous epithelium of uterine cervix  
Nippon Rinsho 54: 1916-21 (1996)
- Gangopadhyay SB; Peacock JW; Benchimol S  
A transcript exhibiting homology to endogenous rat retroviral-like elements is regulated by p53.  
Oncogene 13: 2731-5 (1996)
- Gao Q; Hauser SH; Liu XL; Wazer DE; Madoc-Jones H; Band V  
Mutant p53-induced immortalization of primary human mammary epithelial cells.  
Cancer Res 56: 3129-33 (1996)
- Garrido-Guerrero E; Carrillo E; Guido M; Zamorano R; Garcia-Carranca A; Gariglio P  
Different arrangement of human papillomavirus E2 binding sites distinguishes cutaneous types from those associated with mucosal lesions.  
Arch Med Res 27: 389-94 (1996)
- Garzino-Demo A; Chen M; Lusso P; Berneman Z; DiPaolo JA  
Enhancement of TAT-induced transactivation of the HIV-1 LTR by two genomic fragments of HHV-6.  
J Med Virol 50: 20-4 (1996)
- Geisen C; Kahn T  
Promoter activity of sequences located upstream of the human papillomavirus types of 16 and 18 late regions.  
J Gen Virol 77 ( Pt 9): 2193-200 (1996)
- Ghai J; Ostrow RS; Tolar J; McGlennen RC; Lemke TD; Tobolt D; Liu Z; Faras AJ  
The E5 gene product of rhesus papillomavirus is an activator of endogenous Ras and phosphatidylinositol-3'-kinase in NIH 3T3 cells.  
Proc Natl Acad Sci U S A 93: 12879-84 (1996)

## References (1996)

- Ghim SJ; Young R; Jenson AB  
Antigenicity of bovine papillomavirus type 1 (BPV-1) L1 virus-like particles compared with that of intact BPV-1 virions.  
*J Gen Virol* 77 ( Pt 2 ): 183-8 (1996)
- Gilles C; Piette J; Ploton D; Doco-Fenzy M; Foidart JM  
Viral integration sites in human papilloma virus-33-immortalized cervical keratinocyte cell lines.  
*Cancer Genet Cytogenet* 90: 63-9 (1996)
- Gjooen K; Olsen AO; Magnus P; Grinde B; Sauer T; Orstavik I  
Prevalence of human papillomavirus in cervical scrapes, as analyzed by PCR, in a population-based sample of women with and without cervical dysplasia.  
*APMIS* 104: 68-74 (1996)
- Grassmann K; Kratzer F; Petry KU; Iftner T  
Functional characterization of naturally occurring mutants of human papillomavirus type 16 with large deletions in the non-coding region [letter]  
*Int J Cancer* 68: 265-9 (1996)
- Grassmann K; Rapp B; Maschek H; Petry KU; Iftner T  
Identification of a differentiation-inducible promoter in the E7 open reading frame of human papillomavirus type 16 (HPV-16) in raft cultures of a new cell line containing high copy numbers of episomal HPV-16 DNA.  
*J Virol* 70: 2339-49 (1996)
- Grassmann K; Wilczynski SP; Cook N; Rapp B; Iftner T  
HPV6 variants from malignant tumors with sequence alterations in the regulatory region do not reveal differences in the activities of the oncogene promoters but do contain amino acid exchanges in the E6 and E7 proteins.  
*Virology* 223: 185-97 (1996)
- Grayson W; Taylor L; Cooper K  
Detection of integrated high risk human papillomavirus in adenoid cystic carcinoma of the uterine cervix.  
*J Clin Pathol* 49: 805-9 (1996)
- Grinstein E; Weinert I; Droese B; Pagano M; Royer HD  
Cell cycle regulation of nuclear factor p92 DNA-binding activity by novel phase-specific inhibitors.  
*J Biol Chem* 271: 9215-22 (1996)
- Grossel MJ; Barsoum J; Prakash SS; Androphy EJ  
The BPV-1 E2 DNA-contact helix cysteine is required for transcriptional activation but not replication in mammalian cells.  
*Virology* 217: 301-10 (1996)
- Grossel MJ; Sverdrup F; Breiding DE; Androphy EJ  
Transcriptional activation function is not required for stimulation of DNA replication by bovine papillomavirus type 1 E2.  
*J Virol* 70: 7264-9 (1996)
- Grossman SR; Laimins LA  
EBNA1 and E2: a new paradigm for origin-binding proteins?  
*Trends Microbiol* 4: 87-9 (1996)
- Hamada K; Sakaue M; Alemany R; Zhang WW; Horio Y; Roth JA; Mitchell MF  
Adenovirus-mediated transfer of HPV 16 E6/E7 antisense RNA to human cervical cancer cells.  
*Gynecol Oncol* 63: 219-27 (1996)
- Hamada K; Zhang WW; Alemany R; Wolf J; Roth JA; Mitchell MF  
Growth inhibition of human cervical cancer cells with the recombinant adenovirus p53 in vitro.  
*Gynecol Oncol* 60: 373-9 (1996)
- Harry JB; Wettstein FO  
Transforming properties of the cottontail rabbit papillomavirus oncoproteins Le6 and SE6 and of the E8 protein.  
*J Virol* 70: 3355-62 (1996)
- Hawkins DS; Demers GW; Galloway DA  
Inactivation of p53 enhances sensitivity to multiple chemotherapeutic agents.  
*Cancer Res* 56: 892-8 (1996)
- Heinzel PA; Balaram P; Bernard HU  
Mutations and polymorphisms in the p53, p21 and p16 genes in oral carcinomas of Indian betel quid chewers.  
*Int J Cancer* 68: 420-3 (1996)
- Hembree JR; Agarwal C; Beard RL; Chandraratna RA; Eckert R  
Retinoid X receptor-specific retinoids inhibit the ability of retinoic acid receptor-specific retinoids to increase the level of insulin-like growth factor binding protein-3 in human ectocervical epithelial cells.  
*Cancer Res* 56: 1794-9 (1996)
- Herber R; Liem A; Pitot H; Lambert PF  
Squamous epithelial hyperplasia and carcinoma in mice transgenic for the human papillomavirus type 16 E7 oncogene.  
*J Virol* 70: 1873-81 (1996)
- Hermonat PL; Santin AD; Batchu RB  
The adeno-associated virus Rep78 major regulatory/transformation suppressor protein binds cellular Sp1 in vitro and evidence of a biological effect.  
*Cancer Res* 56: 5299-304 (1996)

- Herrington CS  
Control of HPV replication: implications for squamous neoplasia [editorial]  
J Pathol 178: 237-8 (1996)
- Heselmeyer K; Schrock E; du Manoir S; Blegen H; Shah K; Steinbeck R; Auer G; Ried T  
Gain of chromosome 3q defines the transition from severe dysplasia to invasive carcinoma of the uterine cervix.  
Proc Natl Acad Sci U S A 93: 479-84 (1996)
- Hinchliffe SA; Smith MD; Boon ME; Howard CV; van Velzen D; Rees JL  
Evidence for dissociation of histone mRNA expression from cellular proliferation in cutaneous human papillomavirus infection.  
J Pathol 178: 249-54 (1996)
- Hiraiwa A; Kiyono T; Suzuki S; Ohashi M; Ishibashi M  
E7 proteins of four groups of human papillomaviruses, irrespective of their tissue tropism or cancer association, possess the ability to transactivate transcriptional promoters E2F site dependently.  
Virus Genes 12: 27-35 (1996)
- Hofmann KJ; Neepier MP; Markus HZ; Brown DR; Muller M; Jansen KU  
Sequence conservation within the major capsid protein of human papillomavirus (HPV) type 18 and formation of HPV-18 virus-like particles in *Saccharomyces cerevisiae*.  
J Gen Virol 77 ( Pt 3): 465-8 (1996)
- Hoos A; D'Incan C; Gissmann L; Altmann A; Momburg F; Nindl I; Osen W; Schonning BH; Jochmus I  
Human papillomavirus type 16 (HPV 16) E7 and major histocompatibility complex (MHC) class I and II expression in human keratinocytes in culture.  
Arch Virol 141: 449-58 (1996)
- Huang H; Li CY; Little JB  
Abrogation of P53 function by transfection of HPV16 E6 gene does not enhance resistance of human tumour cells to ionizing radiation.  
Int J Radiat Biol 70: 151-60 (1996)
- Hwang ES; Naeger LK; DiMaio D  
Activation of the endogenous p53 growth inhibitory pathway in HeLa cervical carcinoma cells by expression of the bovine papillomavirus E2 gene.  
Oncogene 12: 795-803 (1996)
- Inoue H; Yamashita A; Hakura A  
Adhesion-dependency of serum-induced p42/p44 MAP kinase activation is released by retroviral oncogenes.  
Virology 225: 223-6 (1996)
- Isacson C; Kesis TD; Hedrick L; Cho KR  
Both cell proliferation and apoptosis increase with lesion grade in cervical neoplasia but do not correlate with human papillomavirus type.  
Cancer Res 56: 669-74 (1996)
- Iyengar S; Shah KV; Kotloff KL; Ghim SJ; Viscidi RP  
Self-assembly of in vitro-translated human papillomavirus type 16 L1 capsid protein into virus-like particles and antigenic reactivity of the protein.  
Clin Diagn Lab Immunol 3: 733-9 (1996)
- Jenkins A; Kristiansen E; Ask E; Servold T; Christensen A; Kristiansen BE  
Human papillomavirus infection in cervical biopsies from Norwegian gynecological in-patients.  
APMIS 104: 30-4 (1996)
- Jurk M; Weissinger F; Lottspeich F; Schwarz U; Winnacker EL  
Characterization of the single-strand-specific BPV-1 origin binding protein, SPSF I, as the HeLa Pur alpha factor.  
Nucleic Acids Res 24: 2799-806 (1996)
- Kandikonda S; Oda D; Niederman R; Sorkin BC  
Cadherin-mediated adhesion is required for normal growth regulation of human gingival epithelial cells.  
Cell Adhes Commun 4: 13-24 (1996)
- Karlen S; Offord EA; Beard P  
Functional promoters in the genome of human papillomavirus type 6b.  
J Gen Virol 77 ( Pt 1): 11-6 (1996)
- Kawase M; Orth G; Jablonska S; Blanchet-Bardon C; Rueda LA; Favre M  
Variability and phylogeny of the L1 capsid protein gene of human papillomavirus type 5: contribution of clusters of non-synonymous mutations and of a 30-nucleotide duplication.  
Virology 221: 189-98 (1996)
- Kaye JN; Starkey WG; Kell B; Biswas C; Raju KS; Best JM; Cason J  
Human papillomavirus type 16 in infants: use of DNA sequence analyses to determine the source of infection.  
J Gen Virol 77 ( Pt 6): 1139-43 (1996)
- Kesis TD; Connolly DC; Hedrick L; Cho KR  
Expression of HPV16 E6 or E7 increases integration of foreign DNA.  
Oncogene 13: 427-31 (1996)

## References (1996)

- Khare S; Kumar KU; Tang SC; Pater MM; Pater A  
Up-regulation of hormone response of human papillomavirus type 16 expression and increased DNA-protein binding by consensus mutations of viral glucocorticoid response elements.  
*J Med Virol* 50: 254-62 (1996)
- Khleif SN; DeGregori J; Yee CL; Otterson GA; Kaye FJ; Nevins JR; Howley PM  
Inhibition of cyclin D-CDK4/CDK6 activity is associated with an E2F-mediated induction of cyclin kinase inhibitor activity.  
*Proc Natl Acad Sci U S A* 93: 4350-4 (1996)
- Kikuchi K; Taniguchi A; Yasumoto S  
Induction of the HPV16 enhancer activity by Jun-B and c-Fos through cooperation of the promoter-proximal AP-1 site and the epithelial cell type-specific regulatory element in fibroblasts.  
*Virus Genes* 13: 45-52 (1996)
- Kim JW; Lee CG; Cho YH; Kim JH; Kim SJ; Kim HK; Park TC; Song SK; Namkoong SE  
CUMC-6, a new diploid human cell line derived from a squamous carcinoma of the uterine cervix.  
*Gynecol Oncol* 62: 230-40 (1996)
- Kim YT; Thomas NF; Kessiss TD; Wilkinson EJ; Hedrick L; Cho KR  
p53 mutations and clonality in vulvar carcinomas and squamous hyperplasias: evidence suggesting that squamous hyperplasias do not serve as direct precursors of human papillomavirus-negative vulvar carcinomas.  
*Hum Pathol* 27: 389-95 (1996)
- Kinoshita T; Shirasawa H; Shino Y; Shimizu K; Moriya H; Simizu B  
Human papillomavirus type 16 E6 protein up-regulates the expression of the high mobility group protein HMG-I(Y) gene in mouse 10T1/2 cells.  
*Virus Res* 42: 119-25 (1996)
- Kissel'jov F; Semionova L; Samoylova E; Mazurenko N; Komisarova E; Zourbitskaya V; Gritzko T; Kozachenko V; Netchushkin M; Petrov S; Smirnov A; Alonso A  
Instability of chromosome 6 microsatellite repeats in human cervical tumors carrying papillomavirus sequences.  
*Int J Cancer* 69: 484-7 (1996)
- Kitagawa K; Yoshikawa H; Onda T; Kawana T; Taketani Y; Yoshikura H; Iwamoto A  
Genomic organization of human papillomavirus type 18 in cervical cancer specimens.  
*Jpn J Cancer Res* 87: 263-8 (1996)
- Klingelutz AJ; Foster SA; McDougall JK  
Telomerase activation by the E6 gene product of human papillomavirus type 16.  
*Nature* 380: 79-82 (1996)
- Knowles G; O'Neil BW; Campo MS  
Phenotypical characterization of lymphocytes infiltrating regressing papillomas.  
*J Virol* 70: 8451-8 (1996)
- Kovelman R; Bilter GK; Glezer E; Tsou AY; Barbosa MS  
Enhanced transcriptional activation by E2 proteins from the oncogenic human papillomaviruses.  
*J Virol* 70: 7549-60 (1996)
- Kowalski LA; Tsang SS; Davison AJ  
Arsenic and chromium enhance transformation of bovine papillomavirus DNA-transfected C3H/10T1/2 cells.  
*Cancer Lett* 103: 65-9 (1996)
- Kremer R; Woodworth CD; Goltzman D  
Expression and action of parathyroid hormone-related peptide in human cervical epithelial cells.  
*Am J Physiol* 271: C164-71 (1996)
- Krul MR; Tijhaar EJ; Kleijne JA; Van Loon AM; Nievers MG; Schipper H; Geerse L; Van der Kolk M; Steerenberg PA; Mooi FR; Den Otter W  
Induction of an antibody response in mice against human papillomavirus (HPV) type 16 after immunization with HPV recombinant Salmonella strains.  
*Cancer Immunol Immunother* 43: 44-8 (1996)
- Kurvinen K; Syrjanen K; Syrjanen S  
p53 and bcl-2 proteins as prognostic markers in human papillomavirus-associated cervical lesions.  
*J Clin Oncol* 14: 2120-30 (1996)
- Labropoulou V; Diakomanolis E; Dailianas S; Kalpaktsoglou K; Rodolakis A; Beaudenon S; Kakkanas A; Mavromara P  
Genital papillomavirus in Greek women with high-grade cervical intraepithelial neoplasia and cervical carcinoma.  
*J Med Virol* 48: 80-7 (1996)
- Lai YM; Yang FP; Pao CC  
Human papillomavirus deoxyribonucleic acid and ribonucleic acid in seminal plasma and sperm cells.  
*Fertil Steril* 65: 1026-30 (1996)
- Lappalainen K; Pirila L; Jaaskelainen I; Syrjanen K; Syrjanen S  
Effects of liposomal antisense oligonucleotides on mRNA and protein levels of the HPV 16 E7 oncogene.  
*Anticancer Res* 16: 2485-92 (1996)



- Larson AA; Kern S; Sommers RL; Yokota J; Cavenee WK; Hampton GM  
Analysis of replication error (RER+) phenotypes in cervical carcinoma.  
*Cancer Res* 56: 1426-31 (1996)
- LeBoit PE  
Correspondence re: Nucci MR, Genest DR, Tate JE, Sparks CK, Crum CP: pseudobowenoid vulvar change: untreated condyloma acuminatum. *Mod Pathol* 9:375, 1996 letter  
*Mod Pathol* 9: 870-1 (1996)
- Ledent C  
Transgenic models of human thyroid pathologies: hyperfunctional adenoma, anaplastic cancer, differentiated cancer, hypothyroidism.  
*Bull Mem Acad R Med Belg* 151: 195-201 (1996)
- Ledent C; Coppee F; Dumont JE; Vassart G; Parmentier M  
Transgenic models for proliferative and hyperfunctional thyroid diseases.  
*Exp Clin Endocrinol Diabetes* 104 Suppl 3: 43-6 (1996)
- Lewensohn-Fuchs I; Ljungman P; Kjerrstrom A; Ringden O; Dalianis T  
Loss of seroreactivity against human papillomavirus (HPV) in bone marrow transplant recipients.  
*Bone Marrow Transplant* 18: 333-7 (1996)
- Li Q; Kondoh G; Inafuku S; Nishimune Y; Hakura A  
Abrogation of c-kit/Steel factor-dependent tumorigenesis by kinase defective mutants of the c-kit receptor: c-kit kinase defective mutants as candidate tools for cancer gene therapy.  
*Cancer Res* 56: 4343-6 (1996)
- Li X; Coffino P  
High-risk human papillomavirus E6 protein has two distinct binding sites within p53, of which only one determines degradation.  
*J Virol* 70: 4509-16 (1996)
- Li X; Coffino P  
Identification of a region of p53 that confers lability.  
*J Biol Chem* 271: 4447-51 (1996)
- Liang H; Petros AM; Meadows RP; Yoon HS; Egan DA; Walter K; Holzman TF; Robins T; Fesik SW  
Solution structure of the DNA-binding domain of a human papillomavirus E2 protein: evidence for flexible DNA-binding regions.  
*Biochemistry* 35: 2095-103 (1996)
- Lie ES; Heyden A; Johannesen MK; Boysen M; Brandtzaeg P  
Detection of human papillomavirus in routinely processed biopsy specimens from laryngeal papillomas: evaluation of reproducibility of polymerase chain reaction and DNA in situ hybridization procedures.  
*Acta Otolaryngol (Stockh)* 116: 627-32 (1996)
- Liggett WH Jr; Sewell DA; Rocco J; Ahrendt SA; Koch W; Sidransky D  
p16 and p16 beta are potent growth suppressors of head and neck squamous carcinoma cells in vitro.  
*Cancer Res* 56: 4119-23 (1996)
- Lizard G; Chignol MC; Chardonnet Y; Schmitt D  
Differences of reactivity to interferon gamma in HeLa and CaSki cells: a combined immunocytochemical and flow-cytometric study.  
*J Cancer Res Clin Oncol* 122: 223-30 (1996)
- Londesborough P; Ho L; Terry G; Cuzick J; Wheeler C; Singer A  
Human papillomavirus genotype as a predictor of persistence and development of high-grade lesions in women with minor cervical abnormalities.  
*Int J Cancer* 69: 364-8 (1996)
- Londono LP; Chatfield S; Tindle RW; Herd K; Gao XM; Frazer I; Dougan G  
Immunisation of mice using Salmonella typhimurium expressing human papillomavirus type 16 E7 epitopes inserted into hepatitis B virus core antigen.  
*Vaccine* 14: 545-52 (1996)
- Longuet M; Beaudenon S; Orth G  
Two novel genital human papillomavirus (HPV) types, HPV68 and HPV70, related to the potentially oncogenic HPV39.  
*J Clin Microbiol* 34: 738-44 (1996)
- Longuet M; Cassonnet P; Orth G  
A novel genital human papillomavirus (HPV), HPV type 74, found in immunosuppressed patients.  
*J Clin Microbiol* 34: 1859-62 (1996)
- Lopez-Beltran A; Escudero AL; Vicioso L; Munoz E; Carrasco JC  
Human papillomavirus DNA as a factor determining the survival of bladder cancer patients.  
*Br J Cancer* 73: 124-7 (1996)
- Lu S; Syrjanen SL; Havu VK; Syrjanen S  
Known HPV types have no association with keratoacanthomas.  
*Arch Dermatol Res* 288: 129-32 (1996)

## References (1996)

- Ludes-Meyers JH; Wilson VG  
Simple procedure for creation of in-frame deletion mutations throughout an open reading frame.  
*Biotechniques* 20: 433-8 (1996)
- Ludmerer SW; Benincasa D; Mark GE 3rd  
Two amino acid residues confer type specificity to a neutralizing, conformationally dependent epitope on human papillomavirus type 11.  
*J Virol* 70: 4791-4 (1996)
- Luxton JC; Rowe AJ; Cridland JC; Coletart T; Wilson P; Shepherd PS  
Proliferative T cell responses to the human papillomavirus type 16 E7 protein in women with cervical dysplasia and cervical carcinoma and in healthy individuals.  
*J Gen Virol* 77 ( Pt 7): 1585-93 (1996)
- Mahony JB  
Multiplex polymerase chain reaction for the diagnosis of sexually transmitted diseases.  
*Clin Lab Med* 16: 61-71 (1996)
- Maki CG; Huibregtse JM; Howley PM  
In vivo ubiquitination and proteasome-mediated degradation of p53(1).  
*Cancer Res* 56: 2649-54 (1996)
- Maki H; Fujikawa-Adachi K; Yoshie O  
Evidence for a promoter-like activity in the short non-coding region of human papillomaviruses.  
*J Gen Virol* 77 ( Pt 3): 453-8 (1996)
- Malejczyk J; Malejczyk M; Breitburd F; Majewski S; Schwarz A; Expert-Besancon N; Jablonska S; Orth G; Luger TA  
Progressive growth of human papillomavirus type 16-transformed keratinocytes is associated with an increased release of soluble tumour necrosis factor (TNF) receptor.  
*Br J Cancer* 74: 234-9 (1996)
- Mao EJ; Oda D; Haigh WG; Beckmann AM  
Loss of the adenomatous polyposis coli gene and human papillomavirus infection in oral carcinogenesis.  
*Eur J Cancer B Oral Oncol* 32B: 260-3 (1996)
- Mao EJ; Schwartz SM; Daling JR; Oda D; Tickman L; Beckmann AM  
Human papilloma viruses and p53 mutations in normal pre-malignant and malignant oral epithelia.  
*Int J Cancer* 69: 152-8 (1996)
- Marionnet AV; Lizard G; Chignol MC; Chardonnet Y; Schmitt D  
Differences in reactivity to cyclosporin A and interferon-gamma of normal and HPV-transformed keratinocytes.  
*Anticancer Res* 16: 257-67 (1996)
- Mark HF; Santoro K; Campbell W; Hann E; Lathrop J  
Integration of human papillomavirus sequences in cervical tumor cell lines.  
*Ann Clin Lab Sci* 26: 147-53 (1996)
- Massimi P; Pim D; Storey A; Banks L  
HPV-16 E7 and adenovirus E1a complex formation with TATA box binding protein is enhanced by casein kinase II phosphorylation.  
*Oncogene* 12: 2325-30 (1996)
- McIntyre MC; Ruesch MN; Laimins LA  
Human papillomavirus E7 oncoproteins bind a single form of cyclin E in a complex with cdk2 and p107.  
*Virology* 215: 73-82 (1996)
- Medina-Martinez O; Morales-Peza N; Yaniv M; Garcia-Carranca A; Thierry F  
A single element mediates glucocorticoid hormone response of HPV18 with no functional interactions with AP1 or hbrm.  
*Virology* 217: 392-6 (1996)
- Merrick DT; Winberg G; McDougall JK  
Re-expression of interleukin 1 in human papillomavirus 18 immortalized keratinocytes inhibits their tumorigenicity in nude mice.  
*Cell Growth Differ* 7: 1661-9 (1996)
- Miller CS; White DK  
Human papillomavirus expression in oral mucosa, premalignant conditions, and squamous cell carcinoma: a retrospective review of the literature.  
*Oral Surg Oral Med Oral Pathol Oral Radiol Endod* 82: 57-68 (1996)
- Mitchell MF; Hamada K; Sastry KJ; Sarkar A; Tortolero-Luna G; Wharton JT; Roth JA  
Transgene expression in the rhesus cervix mediated by an adenovirus expressing beta-galactosidase.  
*Am J Obstet Gynecol* 174: 1094-101 (1996)
- Mok YK; Bycroft M; de Prat-Gay G  
The dimeric DNA binding domain of the human papillomavirus E2 protein folds through a monomeric intermediate which cannot be native-like.  
*Nat Struct Biol* 3: 711-7 (1996)
- Mok YK; de Prat Gay G; Butler PJ; Bycroft M  
Equilibrium dissociation and unfolding of the dimeric human papillomavirus strain-16 E2 DNA-binding domain.  
*Protein Sci* 5: 310-9 (1996)

- Moller B; Lindeberg H  
The presence of HPV types 6/11 and 16 in a giant vulvar carcinoma, arising in pre-existing condylomas, demonstrated by PCR and DNA in-situ hybridization.  
*Eur J Gynaecol Oncol* 17: 497-500 (1996)
- Monini P; Rotola A; de Lellis L; Corallini A; Secchiero P; Albini A; Benelli R; Parravicini C; Barbanti-Brodano G; Cassai E  
Latent BK virus infection and Kaposi's sarcoma pathogenesis.  
*Int J Cancer* 66: 717-22 (1996)
- Moro A; Hernandez P  
Bacterial expression and immunological detection of human papillomavirus type 16 E7 protein.  
*Biochem Biophys Res Commun* 226: 895-9 (1996)
- Muller F; Sapp M  
Domains of the E1 protein of human papillomavirus type 33 involved in binding to the E2 protein.  
*Virology* 219: 247-56 (1996)
- Mullokanov MR; Kholodilov NG; Atkin NB; Burk RD; Johnson AB; Klinger HP  
Genomic alterations in cervical carcinoma: losses of chromosome heterozygosity and human papilloma virus tumor status.  
*Cancer Res* 56: 197-205 (1996)
- Nagai N  
Evaluation of prognostic factors in gynecological cancer examined by molecular biological study  
*Gan To Kagaku Ryoho* 23: 1004-9 (1996)
- Nagano H; Yoshikawa H; Kawana T; Yokota H; Taketani Y; Igarashi H; Yoshikura H; Iwamoto A  
Association of multiple human papillomavirus types with vulvar neoplasias.  
*J Obstet Gynaecol* 22: 1-8 (1996)
- Naghashfar Z; DiPaolo JA; Woodworth CD; Passaniti A  
Immortalization of human adult prostatic adenocarcinoma cells by human papilloma virus HPV16 and -18 DNA.  
*Cancer Lett* 100: 47-54 (1996)
- Nakagawa M; Stites DP; Farhat S; Judd A; Moscicki AB; Canchola AJ; Hilton JF; Palefsky JM  
T-cell proliferative response to human papillomavirus type 16 peptides: relationship to cervical intraepithelial neoplasia.  
*Clin Diagn Lab Immunol* 3: 205-10 (1996)
- Namba M; Mihara K; Fushimi K  
Immortalization of human cells and its mechanisms.  
*Crit Rev Oncog* 7: 19-31 (1996)
- Neeper MP; Hofmann KJ; Jansen KU  
Expression of the major capsid protein of human papillomavirus type 11 in *Saccharomyces cerevisiae*.  
*Gene* 180: 1-6 (1996)
- Nichols GE; Williams ME; Gaffey MJ; Stoler MH  
Cyclin D1 gene expression in human cervical neoplasia.  
*Mod Pathol* 9: 418-25 (1996)
- Nilsson CH; Bakos E; Petry KU; Schneider A; Durst M  
Promoter usage in the E7 ORF of HPV16 correlates with epithelial differentiation and is largely confined to low-grade genital neoplasia.  
*Int J Cancer* 65: 6-12 (1996)
- Noble JR; Rogan EM; Neumann AA; Maclean K; Bryan TM; Reddel RR  
Association of extended in vitro proliferative potential with loss of p16INK4 expression.  
*Oncogene* 13: 1259-68 (1996)
- Nuber U; Schwarz S; Kaiser P; Schneider R; Scheffner M  
Cloning of human ubiquitin-conjugating enzymes UbcH6 and UbcH7 (E2-F1) and characterization of their interaction with E6-AP and RSP5.  
*J Biol Chem* 271: 2795-800 (1996)
- O'Connor MJ; Tan SH; Tan CH; Bernard HU  
YY1 represses human papillomavirus type 16 transcription by quenching AP-1 activity.  
*J Virol* 70: 6529-39 (1996)
- Oda D; Bigler L; Lee P; Blanton R  
HPV immortalization of human oral epithelial cells: a model for carcinogenesis.  
*Exp Cell Res* 226: 164-9 (1996)
- Oda D; Bigler L; Mao EJ; Disteché CM  
Chromosomal abnormalities in HPV-16-immortalized oral epithelial cells.  
*Carcinogenesis* 17: 2003-8 (1996)
- Odunsi K; Terry G; Ho L; Bell J; Cuzick J; Ganesan TS  
Susceptibility to human papillomavirus-associated cervical intra-epithelial neoplasia is determined by specific HLA DR-DQ alleles.  
*Int J Cancer* 67: 595-602 (1996)
- Ozaki T; Hillmann A; Linder N; Blasius S; Winkelmann W  
Metastasis of chondrosarcoma.  
*J Cancer Res Clin Oncol* 122: 629-32 (1996)
- Paintsil J; Muller M; Picken M; Gissmann L; Zhou J  
Carboxyl terminus of bovine papillomavirus type-1 L1 protein is not required for capsid formation.  
*Virology* 223: 238-44 (1996)

## References (1996)

- Pan J; Nakanishi K; Yutsudo M; Inoue H; Li Q; Oka K; Yoshioka N; Hakura A  
Isolation of a novel gene down-regulated by v-src.  
FEBS Lett 383: 21-5 (1996)
- Pandita TK; Hall EJ; Hei TK; Piatyszek MA; Wright WE; Piao CQ; Pandita RK; Willey JC; Geard CR; Kastan MB; Shay JW  
Chromosome end-to-end associations and telomerase activity during cancer progression in human cells after treatment with alpha-particles simulating radon progeny.  
Oncogene 13: 1423-30 (1996)
- Pao CC; Yang FP; Lai YM  
Preferential retention of the E6 and E7 regions of the human papilloma virus type 18 genome by human sperm cells.  
Fertil Steril 66: 630-3 (1996)
- Pao CC; Yao DS; Lin CY; Lee SC; Su HT; Lin SC  
Genomic aberrations of human papillomavirus recovered from cervical cancers.  
Biochem Biophys Res Commun 222: 116-20 (1996)
- Park JS; Rhyu KS; Kim CJ; Kim HS; Han KT; Ahn HK; Kim SJ; Namkoong SE  
Presence of oncogenic HPV DNAs in cervical carcinoma tissues and pelvic lymph nodes associating with proliferating cell nuclear antigen expression.  
Gynecol Oncol 60: 418-23 (1996)
- Park TW; Richart RM; Sun XW; Wright TC Jr  
Association between human papillomavirus type and clonal status of cervical squamous intraepithelial lesions see comments  
J Natl Cancer Inst 88: 355-8 (1996)
- Parker GA; Crook T; Bain M; Sara EA; Farrell PJ; Allday MJ  
Epstein-Barr virus nuclear antigen (EBNA)3C is an immortalizing oncoprotein with similar properties to adenovirus E1A and papillomavirus E7.  
Oncogene 13: 2541-9 (1996)
- Payne D; Chan TS; Wagner R; Tying SK  
Cloning of mucosal and cutaneous HPV sequences in a metastatic squamous cell carcinoma from an epidermodysplasia verruciformis patient.  
Anticancer Res 16: 1165-6 (1996)
- Pei XF  
The human papillomavirus E6/E7 genes induce discordant changes in the expression of cell growth regulatory proteins.  
Carcinogenesis 17: 1395-401 (1996)
- Pelisson I; Soler C; Chardonnet Y; Euvrard S; Schmitt D  
A possible role for human papillomaviruses and c-myc, c-Ha-ras, and p53 gene alterations in malignant cutaneous lesions from renal transplant recipients.  
Cancer Detect Prev 20: 20-30 (1996)
- Peng X; Lang CM; Kreider JW  
Immortalization of inbred rabbit keratinocytes from a Shope papilloma and tumorigenic transformation of the cells by E7-ras.  
Cancer Lett 108: 101-9 (1996)
- Peters DM; Dowd N; Brandt C; Compton T  
Human papilloma virus E6/E7 genes can expand the lifespan of human corneal fibroblasts.  
In Vitro Cell Dev Biol Anim 32: 279-84 (1996)
- Pierrefite V; Binetruy B; Rassoulzadegan M; Cuzin F  
Conditional requirement for sequences distinct from the replication origin during episomal establishment of the BPV1 genome.  
Biochem Biophys Res Commun 229: 445-8 (1996)
- Piirsoo M; Ustav E; Mandel T; Stenlund A; Ustav M  
Cis and trans requirements for stable episomal maintenance of the BPV-1 replicator.  
EMBO J 15: 1-11 (1996)
- Pillai MR; Halabi S; McKalip A; Jayaprakash PG; Rajalekshmi TN; Nair MK; Herman B  
The presence of human papillomavirus-16/-18 E6, p53, and Bcl-2 protein in cervicovaginal smears from patients with invasive cervical cancer.  
Cancer Epidemiol Biomarkers Prev 5: 329-35 (1996)
- Poljak M; Cerar A; Orłowska J  
p53 protein expression in esophageal squamous cell papillomas: a study of 36 lesions.  
Scand J Gastroenterol 31: 10-3 (1996)
- Poljak M; Seme K  
Rapid detection and typing of human papillomaviruses by consensus polymerase chain reaction and enzyme-linked immunosorbent assay.  
J Virol Methods 56: 231-8 (1996)
- Pollanen R; Soini Y; Vuopala S; Laara E; Lehto VP  
Tenascin in human papillomavirus associated lesions of the uterine cervix.  
J Clin Pathol 49: 521-3 (1996)
- Popa LM; Schutz H; Winter S; Kittler L; Lober G  
Specific targeting of human papillomavirus type 16 E7 onco-gene with triple-helix forming purine oligodeoxyribonucleotides.  
Biochem Mol Biol Int 38: 285-95 (1996)

- Puranen M; Saarikoski S; Syrjanen K; Syrjanen S  
Polymerase chain reaction amplification of human papillomavirus DNA from archival, Papanicolaou-stained cervical smears.  
*Acta Cytol* 40: 391-5 (1996)
- Pusch O; Soucek T; Wawra E; Hengstschlager-Ottner E; Bernaschek G; Hengstschlager M  
Specific transformation abolishes cyclin D1 fluctuation throughout the cell cycle.  
*FEBS Lett* 385: 143-8 (1996)
- Puthenveetil JA; Frederickson SM; Reznikoff CA  
Apoptosis in human papillomavirus 16 E7-, but not E6-immortalized human uroepithelial cells.  
*Oncogene* 13: 1123-31 (1996)
- Qi YM; Peng SW; Hengst K; Evander M; Park DS; Zhou J; Frazer IH  
Epithelial cells display separate receptors for papillomavirus VLPs and for soluble L1 capsid protein.  
*Virology* 216: 35-45 (1996)
- Rader JS; Kamarasova T; Huettner PC; Li L; Li Y; Gerhard DS  
Allelotyping of all chromosomal arms in invasive cervical cancer.  
*Oncogene* 13: 2737-41 (1996)
- Rasmussen M; Nielsen PE; Manoharan M; Buchardt O; Koch C  
Specificity of antibodies raised against a specific phosphorothioate oligonucleotide sequence.  
*J Immunol Methods* 189: 47-58 (1996)
- Ratiarson A  
Analysis of the papillomavirus type 16 DNA amplification by PCR in SiHa and SLT cell lines.  
*Microbios* 85: 127-31 (1996)
- Renaud KJ; Cowsert LM  
Characterization of human papillomavirus-11 mRNAs expressed in the context of autonomously replicating viral genomes.  
*Virology* 220: 177-85 (1996)
- Ressing ME; van Driel WJ; Celis E; Sette A; Brandt MP; Hartman M; Anholts JD; Schreuder GM; ter Harmsel WB; Fleuren GJ; Trimbos BJ; Kast WM; Melief CJ  
Occasional memory cytotoxic T-cell responses of patients with human papillomavirus type 16-positive cervical lesions against a human leukocyte antigen-A \*0201-restricted E7-encoded epitope.  
*Cancer Res* 56: 582-8 (1996)
- Reznikoff CA; Yeager TR; Belair CD; Savelieva E; Puthenveetil JA; Stadler WM  
Elevated p16 at senescence and loss of p16 at immortalization in human papillomavirus 16 E6, but not E7, transformed human uroepithelial cells.  
*Cancer Res* 56: 2886-90 (1996)
- Rho J; de Villiers EM; Choe J  
Transforming activities of human papillomavirus type 59 E5, E6 and E7 open reading frames in mouse C127 cells.  
*Virus Res* 44: 57-65 (1996)
- Riethdorf L; Riethdorf S; Gutzlaff K; Prall F; Loning T  
Differential expression of the monocyte chemoattractant protein-1 gene in human papillomavirus-16-infected squamous intraepithelial lesions and squamous cell carcinomas of the cervix uteri.  
*Am J Pathol* 149: 1469-76 (1996)
- Rihet S; Lorenzato M; Clavel C  
Oncogenic human papillomaviruses and ploidy in cervical lesions.  
*J Clin Pathol* 49: 892-6 (1996)
- Roden RB; Greenstone HL; Kirnbauer R; Booy FP; Jessie J; Lowy DR; Schiller JT  
In vitro generation and type-specific neutralization of a human papillomavirus type 16 virion pseudotype.  
*J Virol* 70: 5875-83 (1996)
- Rodrigo Tapia JP; Alvarez Alvarez I; Martinez Sanchez JA; Lazo PS; Ramos S; Sampredo Nuno A; Suarez Nieto C  
The relation between human papilloma virus and cellular DNA content in epidermoid carcinoma of the head and neck  
*Acta Otorrinolaringol Esp* 47: 343-7 (1996)
- Rodriguez JA; Barandika O; Innes D; Mugica van Herckenrode C  
p53 functional status in HPV-positive human primary cervical carcinoma.  
*Int J Dev Biol Suppl* 1: 303S (1996)
- Roumen FJ; Boon ME; van Velzen D; Dieben TO; Coelingh Benink HJ  
The cervico-vaginal epithelium during 20 cycles' use of a combined contraceptive vaginal ring.  
*Hum Reprod* 11: 2443-8 (1996)
- Rowan S; Ludwig RL; Haupt Y; Bates S; Lu X; Oren M; Vousden KH  
Specific loss of apoptotic but not cell-cycle arrest function in a human tumor derived p53 mutant.  
*EMBO J* 15: 827-38 (1996)

## References (1996)

- Rubben A; Baron JM; Grussendorf-Conen EI  
Demonstration of human papillomavirus type 16-related DNA and absence of detectable p53 gene mutations in widespread cutaneous squamous cell carcinomas after oral psoralen with UV-A treatment letter  
*Arch Dermatol* 132: 1257-9 (1996)
- Runnebaum IB; Kohler T; Stickeler E; Kieback HR; Kreienberg R  
p53 mutation is associated with high S-phase fraction in primary fallopian tube adenocarcinoma.  
*Br J Cancer* 74: 1157-60 (1996)
- Sadler GP; Morgan JM; Jasani B; Douglas-Jones A; Wheeler MH  
Epidermal growth factor receptor status in hyperparathyroidism: immunocytochemical and In situ hybridization study.  
*World J Surg* 20: 736-42; discussion 742-3 (1996)
- Sagerman PM; Choi YJ; Hu Y; Niedt GW  
Human papilloma virus, vulvar dystrophy, and vulvar carcinoma: differential expression of human papillomavirus and vulvar dystrophy in the presence and absence of squamous cell carcinoma of the vulva.  
*Gynecol Oncol* 61: 328-32 (1996)
- Sakai H; Yasugi T; Benson JD; Dowhanick JJ; Howley PM  
Targeted mutagenesis of the human papillomavirus type 16 E2 transactivation domain reveals separable transcriptional activation and DNA replication functions.  
*J Virol* 70: 1602-11 (1996)
- Sandler AB; Baker CC; Spalholz BA  
Sp1 is critical for basal and E2-transactivated transcription from the bovine papillomavirus type 1 P89 promoter.  
*J Gen Virol* 77 ( Pt 2 ): 189-98 (1996)
- Sanjeevi CB; Hjelmstrom P; Hallmans G; Wiklund F; Lenner P; Angstrom T; Dillner J; Lernmark A  
Different HLA-DR-DQ haplotypes are associated with cervical intraepithelial neoplasia among human papillomavirus type-16 seropositive and seronegative Swedish women.  
*Int J Cancer* 68: 409-14 (1996)
- Sarma D; Yang X; Jin G; Shindoh M; Pater MM; Pater A  
Resistance to retinoic acid and altered cytokeratin expression of human papillomavirus type 16-immortalized endocervical cells after tumorigenesis.  
*Int J Cancer* 65: 345-50 (1996)
- Sasaoka R; Morimura T; Mihara M; Hagari Y; Aki T; Miyamoto T  
Detection of human papillomavirus type 16 DNA in two cases of verrucous carcinoma of the foot letter  
*Br J Dermatol* 134: 983-4 (1996)
- Sastre-Garau X; Loste MN; Vincent-Salomon A; Favre M; Mouret E; de la Rochefordiere A; Durand JC; Tartour E; Lepage V; Charron D  
Decreased frequency of HLA-DRB1 13 alleles in French-women with HPV-positive carcinoma of the cervix.  
*Int J Cancer* 69: 159-64 (1996)
- Schaller J; Rohwedder A; Fuchs M; Maron A; Kunze J  
HPV-5 typing with nested PCR and sequencing in epidermodysplasia verruciformis  
*Hautarzt* 47: 454-8 (1996)
- Schulz-Wendtland R; Kramer S; Keilholz L; Pfluger S; Lang N  
The dosage-performance effects on Ca-Ski and HPK cells in relation to the dose and fractionation  
*Strahlenther Onkol* 172: 439-45 (1996)
- Sedman J; Stenlund A  
The initiator protein E1 binds to the bovine papillomavirus origin of replication as a trimeric ring-like structure.  
*EMBO J* 15: 5085-92 (1996)
- Shally M; Alloul N; Jackman A; Muller M; Gissmann L; Sherman L  
The E6 variant proteins E6I-E6IV of human papillomavirus 16: expression in cell free systems and bacteria and study of their interaction with p53.  
*Virus Res* 42: 81-96 (1996)
- Shamanin V; zur Hausen H; Lavergne D; Proby CM; Leigh IM; Neumann C; Hamm H; Goos M; Hausteil UF; Jung EG; Plewig G; Wolff H; de Villiers EM  
Human papillomavirus infections in nonmelanoma skin cancers from renal transplant recipients and nonimmunosuppressed patients see comments  
*J Natl Cancer Inst* 88: 802-11 (1996)
- Shan B; Farmer AA; Lee WH  
The molecular basis of E2F-1/DP-1-induced S-phase entry and apoptosis.  
*Cell Growth Differ* 7: 689-97 (1996)
- Shen J; Tate JE; Crum CP; Goodman ML  
Prevalence of human papillomaviruses (HPV) in benign and malignant tumors of the upper respiratory tract.  
*Mod Pathol* 9: 15-20 (1996)
- Shen MR; Wu SN; Chou CY  
Volume-sensitive chloride channels in the primary culture cells of human cervical carcinoma.  
*Biochim Biophys Acta* 1315: 138-44 (1996)

- Shepherd PS; Rowe AJ; Cridland JC; Coletart T; Wilson P; Luxton JC  
Proliferative T cell responses to human papillomavirus type 16 L1 peptides in patients with cervical dysplasia.  
*J Gen Virol* 77 ( Pt 4): 593-602 (1996)
- Sherman L; Schlegel R  
Serum- and calcium-induced differentiation of human keratinocytes is inhibited by the E6 oncoprotein of human papillomavirus type 16.  
*J Virol* 70: 3269-79 (1996)
- Shimakage M; Miyata Y; Inoue H; Yutsudo M; Hakura A  
Increased sensitivity of EBNA2-transformed rat fibroblasts to ionizing radiation.  
*Int J Cancer* 68: 612-5 (1996)
- Shin KH; Tannyhill RJ; Liu X; Park NH  
Oncogenic transformation of HPV-immortalized human oral keratinocytes is associated with the genetic instability of cells.  
*Oncogene* 12: 1089-96 (1996)
- Shindoh M; Kimura S; Yamashita T; Fujinaga K; Amemiya A  
Histopathological application of in situ PCR (in situ hybridization after amplification by PCR)  
*Tanpakushitsu Kakusan Koso* 41: 660-3 (1996)
- Shoji Y; Saegusa M; Takano Y; Ohbu M; Okayasu I  
Correlation of apoptosis with tumour cell differentiation, progression, and HPV infection in cervical carcinoma.  
*J Clin Pathol* 49: 134-8 (1996)
- Skiadopoulos MH; McBride AA  
The bovine papillomavirus type 1 E2 transactivator and repressor proteins use different nuclear localization signals.  
*J Virol* 70: 1117-24 (1996)
- Smith ML; Kontny HU; Zhan Q; Sreenath A; O'Connor PM; Fornace AJ Jr  
Antisense GADD45 expression results in decreased DNA repair and sensitizes cells to u.v.-irradiation or cisplatin.  
*Oncogene* 13: 2255-63 (1996)
- Smits HL  
The use of consensus PCR and direct sequence analysis for the identification of HPV.  
*Methods Mol Biol* 50: 109-26 (1996)
- Snijders PJ; Scholes AG; Hart CA; Jones AS; Vaughan ED; Woolgar JA; Meijer CJ; Walboomers JM; Field JK  
Prevalence of mucosotropic human papillomaviruses in squamous-cell carcinoma of the head and neck.  
*Int J Cancer* 66: 464-9 (1996)
- Soini Y; Nuorva K; Kamel D; Pollanen R; Vahakangas K; Lehto VP; Paakko P  
Presence of human papillomavirus DNA and abnormal p53 protein accumulation in lung carcinoma see comments  
*Thorax* 51: 887-93 (1996)
- Soini Y; Pollanen R; Kempainen S; Paakko P; Lehto VP  
The association of vascular proliferation with HPV status and epithelial PCNA positivity in cervical intraepithelial lesions.  
*APMIS* 104: 183-90 (1996)
- Sparkowski J; Mense M; Anders J; Schlegel R  
E5 oncoprotein transmembrane mutants dissociate fibroblast transforming activity from 16-kilodalton protein binding and platelet-derived growth factor receptor binding and phosphorylation.  
*J Virol* 70: 2420-30 (1996)
- Spitkovsky D; Aengeneyndt F; Braspenning J; von Knebel Doeberitz M  
p53-independent growth regulation of cervical cancer cells by the papillomavirus E6 oncoprotein.  
*Oncogene* 13: 1027-35 (1996)
- Steenbergen RD; Walboomers JM; Meijer CJ; van der Raaij-Helmer EM; Parker JN; Chow LT; Broker TR; Snijders PJ  
Transition of human papillomavirus type 16 and 18 transfected human foreskin keratinocytes towards immortality: activation of telomerase and allele losses at 3p, 10p, 11q and/or 18q.  
*Oncogene* 13: 1249-57 (1996)
- Steger G; Ham J; Yaniv M  
E2 proteins: modulators of papillomavirus transcription and replication.  
*Methods Enzymol* 274: 173-85 (1996)
- Steller MA; Zou Z; Schiller JT; Baserga R  
Transformation by human papillomavirus 16 E6 and E7: role of the insulin-like growth factor 1 receptor.  
*Cancer Res* 56: 5087-91 (1996)
- Stewart AC; Eriksson AM; Manos MM; Munoz N; Bosch FX; Peto J; Wheeler CM  
Intratype variation in 12 human papillomavirus types: a worldwide perspective.  
*J Virol* 70: 3127-36 (1996)
- Stoppler H; Kirchhoff T; Kartenbeck J; Gissmann L; Alonso A  
Influence of human papillomavirus type 16 gene expression on in vitro differentiation of the human teratocarcinoma cell line 2102Ep.  
*Mol Carcinog* 16: 109-14 (1996)

## References (1996)

- Stoppler H; Koval D; Schlegel R  
The serine protease inhibitors TLCK and TPCK inhibit the in vitro immortalization of primary human keratinocytes by HPV-18 DNA.  
*Oncogene* 13: 1545-8 (1996)
- Stoppler H; Stoppler MC; Adduci A; Koval D; Schlegel R  
The serine protease inhibitors TLCK and TPCK react with the RB-binding core of HPV-18 E7 protein and abolish its RB-binding capability.  
*Virology* 217: 542-53 (1996)
- Stoppler MC; Ching K; Stoppler H; Clancy K; Schlegel R; Icenogle J  
Natural variants of the human papillomavirus type 16 E6 protein differ in their abilities to alter keratinocyte differentiation and to induce p53 degradation.  
*J Virol* 70: 6987-93 (1996)
- Stoppler MC; Straight SW; Tsao G; Schlegel R; McCance DJ  
The E5 gene of HPV-16 enhances keratinocyte immortalization by full-length DNA.  
*Virology* 223: 251-4 (1996)
- Stubenrauch F; Leigh IM; Pfister H  
E2 represses the late gene promoter of human papillomavirus type 8 at high concentrations by interfering with cellular factors.  
*J Virol* 70: 119-26 (1996)
- Su PF; Wu FY  
Differential suppression of the tumorigenicity of HeLa and SiHa cells by adeno-associated virus.  
*Br J Cancer* 73: 1533-7 (1996)
- Sun YN; Lu JZ; McCance DJ  
Mapping of HPV-11 E1 binding site and determination of other important cis elements for replication of the origin.  
*Virology* 216: 219-22 (1996)
- Sundaram P; Brandsma JL  
Rapid, efficient, large-scale purification of unfused, non-denatured E7 protein of cottontail rabbit papillomavirus.  
*J Virol Methods* 57: 61-70 (1996)
- Suzuk L; Noffsinger AE; Hui YZ; Fenoglio-Preiser CM  
Detection of human papillomavirus in esophageal squamous cell carcinoma.  
*Cancer* 78: 704-10 (1996)
- Suzuki H; Komiya A; Aida S; Ito H; Yatani R; Shimazaki J  
Detection of human papillomavirus DNA and p53 gene mutations in human prostate cancer.  
*Prostate* 28: 318-24 (1996)
- Suzuki T; Tsutsumi K; Nakajima T; Hara F; Kikuchi K; Yasumoto S  
Spontaneous mutation of p53 gene in human papillomavirus type 16 E7-immortalized human laryngeal epithelial cells.  
*Acta Otolaryngol Suppl (Stockh)* 522: 94-8 (1996)
- Taja-Chayeb L; Salas-Garcia M; Salcedo-Vargas M  
Molecular bases of papillomavirus and polyomavirus carcinogenesis  
*Salud Publica Mex* 38: 47-57 (1996)
- Tarunina M; Grimaldi M; Ruaro E; Pavlenko M; Schneider C; Jenkins JR  
Selective loss of endogenous p21waf1/cip1 induction underlies the G1 checkpoint defect of monomeric p53 proteins.  
*Oncogene* 13: 589-98 (1996)
- Tate JE; Yang YC; Shen J; McLachlin CM; Sheets EE; Crum CP  
A comparison of early (E7) and late (L1) primer-mediated amplification of papillomaviral DNA in cervical neoplasia.  
*Mol Cell Probes* 10: 347-51 (1996)
- Tenti P; Romagnoli S; Silini E; Zappatore R; Spinillo A; Giunta P; Cappellini A; Vesentini N; Zara C; Carnevali L  
Human papillomavirus types 16 and 18 infection in infiltrating adenocarcinoma of the cervix: PCR analysis of 138 cases and correlation with histologic type and grade.  
*Am J Clin Pathol* 106: 52-6 (1996)
- Tenti P; Zappatore R; Romagnoli S; Civardi E; Giunta P; Scelsi R; Stella G; Carnevali L  
p53 overexpression and human papillomavirus infection in transitional cell carcinoma of the urinary bladder: correlation with histological parameters.  
*J Pathol* 178: 65-70 (1996)
- Thain A; Jenkins O; Clarke AR; Gaston K  
CpG methylation directly inhibits binding of the human papillomavirus type 16 E2 protein to specific DNA sequences.  
*J Virol* 70: 7233-5 (1996)
- Thomas M; Massimi P; Banks L  
HPV-18 E6 inhibits p53 DNA binding activity regardless of the oligomeric state of p53 or the exact p53 recognition sequence.  
*Oncogene* 13: 471-80 (1996)
- Thomas M; Matlashewski G; Pim D; Banks L  
Induction of apoptosis by p53 is independent of its oligomeric state and can be abolished by HPV-18 E6 through ubiquitin mediated degradation.  
*Oncogene* 13: 265-73 (1996)
- Tindle RW  
Human papillomavirus vaccines for cervical cancer.  
*Curr Opin Immunol* 8: 643-50 (1996)



- Toledo-Cuevas EM; Garcia-Carranca A  
P53 protein and human papillomavirus oncogenes in carcinogenesis of the uterine cervix  
*Rev Invest Clin* 48: 59-68 (1996)
- Tomita Y; Shiga T; Simizu B  
Characterization of a promoter in the E7 open reading frame of human papillomavirus type 11.  
*Virology* 225: 267-73 (1996)
- Touze A; Dupuy C; Chabaud M; Le Cann P; Coursaget P  
Production of human papillomavirus type 45 virus-like particles in insect cells using a recombinant baculovirus.  
*FEMS Microbiol Lett* 141: 111-6 (1996)
- Trujillo JM; Mounts P  
Transforming activity of the E6 gene of HPV-11gt in NIH 3T3 and REF 52 cells: correlation with the level of E6 transcription.  
*Virology* 220: 1-9 (1996)
- Trujillo JM; Wu TC; Mounts P  
Characterization of human papillomavirus type 57b: transforming activity and comparative sequence analysis as probes for biological determinants associated with high-risk oncogenic viruses.  
*Virus Genes* 12: 165-78 (1996)
- Tsao YP; Li LY; Tsai TC; Chen SL  
Human papillomavirus type 11 and 16 E5 represses p21(Waf1/Sdi1/Cip1) gene expression in fibroblasts and keratinocytes.  
*J Virol* 70: 7535-9 (1996)
- Tsukui T; Hildesheim A; Schiffman MH; Lucci J 3rd; Contois D; Lawler P; Rush BB; Lorincz AT; Corrigan A; Burk RD; Qu W; Marshall MA; Mann D; Carrington M; Clerici M; Shearer GM; Carbone DP; Scott DR; Houghten RA; Berzofsky JA  
Interleukin 2 production in vitro by peripheral lymphocytes in response to human papillomavirus-derived peptides: correlation with cervical pathology.  
*Cancer Res* 56: 3967-74 (1996)
- Tsutsumi K; Iwatake H; Suzuki T  
An experimental model of multistep laryngeal carcinogenesis: combined effect of human papillomavirus type 16 genome and N-methyl-N'-nitro-N-nitrosoguanidine.  
*Acta Otolaryngol Suppl (Stockh)* 522: 89-93 (1996)
- Turek LP; Smith EM  
The genetic program of genital human papillomaviruses in infection and cancer.  
*Obstet Gynecol Clin North Am* 23: 735-58 (1996)
- Ullman CG; Haris PI; Galloway DA; Emery VC; Perkins SJ  
Predicted alpha-helix/beta-sheet secondary structures for the zinc-binding motifs of human papillomavirus E7 and E6 proteins by consensus prediction averaging and spectroscopic studies of E7.  
*Biochem J* 319 ( Pt 1): 229-39 (1996)
- Ulmer JB; Deck RR; Yawman A; Friedman A; Dewitt C; Martinez D; Montgomery DL; Donnelly JJ; Liu MA  
DNA vaccines for bacteria and viruses.  
*Adv Exp Med Biol* 397: 49-53 (1996)
- van der Burg SH; Visseren MJ; Brandt RM; Kast WM; Melief CJ  
Immunogenicity of peptides bound to MHC class I molecules depends on the MHC-peptide complex stability.  
*J Immunol* 156: 3308-14 (1996)
- Van Rensburg EJ; Engelbrecht S; Van Heerden WF; Raubenneheimer EJ; Schoub BD  
Human papillomavirus DNA in oral squamous cell carcinomas from an African population sample.  
*Anticancer Res* 16: 969-73 (1996)
- Vieira KB; Goldstein DJ; Villa LL  
Tumor necrosis factor alpha interferes with the cell cycle of normal and papillomavirus-immortalized human keratinocytes.  
*Cancer Res* 56: 2452-7 (1996)
- Volter C; He Y; Delius H; Roy-Burman A; Greenspan JS; Greenspan D; de Villiers EM  
Novel HPV types present in oral papillomatous lesions from patients with HIV infection.  
*Int J Cancer* 66: 453-6 (1996)
- Voog E  
Genital viral infections. Studies on human papillomavirus and Epstein-Barr virus.  
*Acta Derm Venereol Suppl (Stockh)* 198: 1-55 (1996)
- Vousden KH  
HPV E6: ensuring all's well at the end.  
*Trends Microbiol* 4: 337-8 (1996)
- Wahl AF; Donaldson KL; Fairchild C; Lee FY; Foster SA; Demers GW; Galloway DA  
Loss of normal p53 function confers sensitization to Taxol by increasing G2/M arrest and apoptosis.  
*Nat Med* 2: 72-9 (1996)
- Walker F; Darai E; Dauge-Geffroy MC; Joannes M; Soustre E; Lehy T; Potet F  
Genital infection with human papillomavirus (HPV): viral screening and typing, importance of in situ hybridization  
*Ann Pathol* 16: 364-73 (1996)

## References (1996)

- Wang H; Liu K; Yuan F; Berdichevsky L; Taichman LB; Auburn K  
C/EBPbeta is a negative regulator of human papillomavirus type 11 in keratinocytes.  
*J Virol* 70: 4839-44 (1996)
- Wang M; Fudge K; Rhim JS; Stearns ME  
Cytokine regulation of the matrix metalloproteinases and their inhibitors in human papillomavirus-18 transformed human prostatic tumor cell lines.  
*Oncol Res* 8: 303-15 (1996)
- Wang Q; Fan S; Eastman A; Worland PJ; Sausville EA; O'Connor PM  
UCN-01: a potent abrogator of G2 checkpoint function in cancer cells with disrupted p53.  
*J Natl Cancer Inst* 88: 956-65 (1996)
- Wang Y; Okan I; Pokrovskaja K; Wiman KG  
Abrogation of p53-induced G1 arrest by the HPV 16 E7 protein does not inhibit p53-induced apoptosis.  
*Oncogene* 12: 2731-5 (1996)
- Wilding J; Vousden KH; Soutter WP; McCrea PD; Del Buono R; Pignatelli M  
E-cadherin transfection down-regulates the epidermal growth factor receptor and reverses the invasive phenotype of human papilloma virus-transfected keratinocytes.  
*Cancer Res* 56: 5285-92 (1996)
- Wilke CM; Hall BK; Hoge A; Paradee W; Smith DI; Glover TW  
FRA3B extends over a broad region and contains a spontaneous HPV16 integration site: direct evidence for the coincidence of viral integration sites and fragile sites.  
*Hum Mol Genet* 5: 187-95 (1996)
- Willis SD; Seyfred MA  
Pituitary-specific chromatin structure of the rat prolactin distal enhancer element.  
*Nucleic Acids Res* 24: 1065-72 (1996)
- Winokur PL; McBride AA  
The transactivation and DNA binding domains of the BPV-1 E2 protein have different roles in cooperative origin binding with the E1 protein.  
*Virology* 221: 44-53 (1996)
- Wong HK; Ziff EB  
The human papillomavirus type 16 E7 protein complements adenovirus type 5 E1A amino-terminus-dependent transactivation of adenovirus type 5 early genes and increases ATF and Oct-1 DNA binding activity.  
*J Virol* 70: 332-40 (1996)
- Woodworth CD; Chung J; McMullin E; Plowman GD; Simpson S; Iglesias M  
Transforming growth factor beta 1 supports autonomous growth of human papillomavirus-immortalized cervical keratinocytes under conditions promoting squamous differentiation.  
*Cell Growth Differ* 7: 811-20 (1996)
- Xiao W; Brandsma JL  
High efficiency, long-term clinical expression of cottontail rabbit papillomavirus (CRPV) DNA in rabbit skin following particle-mediated DNA transfer.  
*Nucleic Acids Res* 24: 2620-2 (1996)
- Xiong Y; Kuppuswamy D; Li Y; Livanos EM; Hixon M; White A; Beach D; Tlsty TD  
Alteration of cell cycle kinase complexes in human papillomavirus E6- and E7-expressing fibroblasts precedes neoplastic transformation.  
*J Virol* 70: 999-1008 (1996)
- Yamashita T; Inoue Y; Fujinaga Y; Fujinaga K  
Detection and typing of HPV genome: consensus PCR method  
*Tanpakushitsu Kakusan Koso* 41: 664-8 (1996)
- Yan Y; Ouellette MM; Shay JW; Wright WE  
Age-dependent alterations of c-fos and growth regulation in human fibroblasts expressing the HPV16 E6 protein.  
*Mol Biol Cell* 7: 975-83 (1996)
- Yang X; Jin G; Nakao Y; Rahimtula M; Pater MM; Pater A  
Malignant transformation of HPV 16-immortalized human endocervical cells by cigarette smoke condensate and characterization of multistage carcinogenesis.  
*Int J Cancer* 65: 338-44 (1996)
- Yang X; Nakao Y; Pater MM; Pater A  
Identification of two novel cellular genes associated with multistage carcinogenesis of human endocervical cells by mRNA differential display.  
*Carcinogenesis* 17: 563-7 (1996)
- Yang YF; Tsao YP; Yin CS; Chen SL; Chu TY  
Overexpression of the proto-oncogene c-jun in association with low-risk type specific human papillomavirus infection in condyloma acuminata.  
*J Med Virol* 48: 302-7 (1996)
- Yasugi T; Howley PM  
Identification of the structural and functional human homolog of the yeast ubiquitin conjugating enzyme UBC9.  
*Nucleic Acids Res* 24: 2005-10 (1996)

- Yukawa K; Butz K; Yasui T; Kikutani H; Hoppe-Seyley F  
Regulation of human papillomavirus transcription by the differentiation-dependent epithelial factor Epc-1/skn-1a.  
*J Virol* 70: 10-6 (1996)
- Zehbe I; Rylander E; Edlund K; Wadell G; Wilander E  
Detection of human papillomavirus in cervical intra-epithelial neoplasia, using in situ hybridization and various polymerase chain reaction techniques.  
*Virchows Arch* 428: 151-7 (1996)
- Zehbe I; Sallstrom JF; Evander M; Edlund K; Rylander E; Wadell G; Wilander E  
Nonradioisotopic detection and typing of human papillomaviruses by use of polymerase chain reaction and single-strand conformation polymorphism.  
*Diagn Mol Pathol* 5: 206-13 (1996)
- Zehbe I; Strand A; Chua KL; Wilander E  
Cytological evaluation and molecular human papillomavirus test of cervical scrapings from women treated for condyloma.  
*Gynecol Obstet Invest* 42: 128-32 (1996)
- Zehbe I; Wilander E  
Two consensus primer systems and nested polymerase chain reaction for human papillomavirus detection in cervical biopsies: A study of sensitivity.  
*Hum Pathol* 27: 812-5 (1996)
- Zerfass-Thome K; Zwerschke W; Mannhardt B; Tindle R; Botz JW; Jansen-Durr P  
Inactivation of the cdk inhibitor p27KIP1 by the human papillomavirus type 16 E7 oncoprotein.  
*Oncogene* 13: 2323-30 (1996)
- Zhang L; Kashanchi F; Zhan Q; Zhan S; Brady JN; Fornace AJ; Seth P; Helman LJ  
Regulation of insulin-like growth factor II P3 promoter by p53: a potential mechanism for tumorigenesis.  
*Cancer Res* 56: 1367-73 (1996)
- Zhao C; Tan W; Sokolowski M; Schwartz S  
Identification of nuclear and cytoplasmic proteins that interact specifically with an AU-rich, cis-acting inhibitory sequence in the 3' untranslated region of human papillomavirus type 1 late mRNAs.  
*J Virol* 70: 3659-67 (1996)
- Zheng J; Siren V; Vaheri A  
Keratinocyte growth factor enhances urokinase-type plasminogen activator activity in HPV16 DNA-immortalized human uterine exocervical epithelial cells.  
*Eur J Cell Biol* 69: 128-34 (1996)
- Zheng ZM; He P; Baker CC  
Selection of the bovine papillomavirus type 1 nucleotide 3225 3' splice site is regulated through an exonic splicing enhancer and its juxtaposed exonic splicing suppressor.  
*J Virol* 70: 4691-9 (1996)
- Zou S; Brown EG  
Stable expression of the reovirus mu2 protein in mouse L cells complements the growth of a reovirus ts mutant with a defect in its M1 gene.  
*Virology* 217: 42-8 (1996)
- zur Hausen H  
Papillomavirus infections—a major cause of human cancers.  
*Biochim Biophys Acta* 1288: F55-78 (1996)
- Zwerschke W; Joswig S; Jansen-Durr P  
Identification of domains required for transcriptional activation and protein dimerization in the human papillomavirus type-16 E7 protein.  
*Oncogene* 12: 213-20 (1996)

## References (1997)

- al-Saleh W; Delvenne P; van den Brule FA; Menard S; Boniver J; Castronovo V  
Expression of the 67 KD laminin receptor in human cervical preneoplastic and neoplastic squamous epithelial lesions: an immunohistochemical study.  
J Pathol 181: 287-93 (1997) Beer-Romero P; Glass S; Rolfe M  
Antisense targeting of E6AP elevates p53 in HPV-infected cells but not in normal cells.  
Oncogene 14: 595-602 (1997)
- Belaguli NS; Pater MM; Pater A  
Identification and location of human papillomavirus type 16 antisense early promoter and characterisation of antisense RNA.  
J Med Virol 51: 344-54 (1997)
- Berg M; Stenlund A  
Functional interactions between papillomavirus E1 and E2 proteins.  
J Virol 71: 3853-63 (1997)
- Bergqvist A; Soderbarg K; Magnusson G  
Altered susceptibility to tumor necrosis factor alpha-induced apoptosis of mouse cells expressing polyomavirus middle and small T antigens.  
J Virol 71: 276-83 (1997)
- Bloch N; Sutton RH; Breen M; Spradbrow PB  
Identification of papillomaviruses in scrapings from bovine warts by use of the polymerase chain reaction.  
Vet Res Commun 21: 63-8 (1997)
- Bornstein J; Lahat N; Kinarty A; Revel M; Abramovici H; Shapiro S  
Interferon-beta and -gamma, but not tumor necrosis factor-alpha, demonstrate immunoregulatory effects on carcinoma cell lines infected with human papillomavirus.  
Cancer 79: 924-34 (1997)
- Boxman IL; Berkhout RJ; Mulder LH; Wolkers MC; Bouwes Bavinck JN; Vermeer BJ; ter Schegget J  
Detection of human papillomavirus DNA in plucked hairs from renal transplant recipients and healthy volunteers.  
J Invest Dermatol 108: 712-5 (1997)
- Brown JP; Pagano M  
Mechanism of p53 degradation.  
Biochim Biophys Acta 1332: O1-6 (1997)
- Burger MP; van Leeuwen AM; Hollema H; Quint WG; Pieters WJ  
Human papillomavirus type influences the extent of chromosomal lag during mitosis in cervical intraepithelial neoplasia grade III.  
Int J Gynecol Pathol 16: 10-4 (1997)
- Cabrera M; Chan PJ; Kalugdan TH; King A  
Transfection of the inner cell mass and lack of a unique DNA sequence affecting the uptake of exogenous DNA by sperm as shown by dideoxy sequencing analogues.  
J Assist Reprod Genet 14: 120-4 (1997)
- Chen Z; Storthz KA; Shillitoe EJ  
Mutations in the long control region of human papillomavirus DNA in oral cancer cells, and their functional consequences.  
Cancer Res 57: 1614-9 (1997)
- Chopra KF; Tyring SK  
The impact of the human immunodeficiency virus on the human papillomavirus epidemic.  
Arch Dermatol 133: 629-33 (1997)
- Cultraro CM; Bino T; Segal S  
Function of the c-Myc antagonist Mad1 during a molecular switch from proliferation to differentiation.  
Mol Cell Biol 17: 2353-9 (1997)
- Daniel B; Rangarajan A; Mukherjee G; Vallikad E; Krishna S  
The link between integration and expression of human papillomavirus type 16 genomes and cellular changes in the evolution of cervical intraepithelial neoplastic lesions.  
J Gen Virol 78 ( Pt 5): 1095-101 (1997)
- DeWeese TL; Walsh JC; Dillehay LE; Kessis TD; Hedrick L; Cho KR; Nelson WG  
Human papillomavirus E6 and E7 oncoproteins alter cell cycle progression but not radiosensitivity of carcinoma cells treated with low-dose-rate radiation.  
Int J Radiat Oncol Biol Phys 37: 145-54 (1997)
- Desaintes C; Demeret C; Goyat S; Yaniv M; Thierry F  
Expression of the papillomavirus E2 protein in HeLa cells leads to apoptosis.  
EMBO J 16: 504-14 (1997)
- Dietrich-Goetz W; Kennedy IM; Levins B; Stanley MA; Clements JB  
A cellular 65-kDa protein recognizes the negative regulatory element of human papillomavirus late mRNA.  
Proc Natl Acad Sci U S A 94: 163-8 (1997)
- Drummond-Barbosa D; DiMaio D  
Virocrine transformation.  
Biochim Biophys Acta 1332: M1-17 (1997)
- Duggan MA; McGregor SE; Stuart GC; Morris S; Chang-Poon V; Schepansky A; Honore L  
The HPV determinants of CIN I.  
Eur J Gynaecol Oncol 18: 117-23 (1997)

- Fallani MG; Pirami L; Penna C; Giache V; Zipoli E; Cardelli A; Marchionni M; Becciolini A  
Cervical intraepithelial neoplasia (CIN) and human papillomavirus (HPV) infection  
*Minerva Ginecol* 49: 13-8 (1997)
- Fan S; Chang JK; Smith ML; Duba D; Fornace AJ Jr; O'Connor PM  
Cells lacking CIP1/WAF1 genes exhibit preferential sensitivity to cisplatin and nitrogen mustard.  
*Oncogene* 14: 2127-36 (1997)
- Feldman SR; Lye KD; Smith ES; Lo AC; Borcik LD  
Polymerase chain reaction and its application for the diagnosis and management of skin diseases.  
*Adv Dermatol* 12: 115-36; discussion 137 (1997)
- Frank GA; Minkina GN; Kulagina LA; Simonova TV; Petrov AN  
Proliferating cell nuclear antigen in precancerous cervical disease  
*Arkh Patol* 59: 30-3 (1997)
- Frattini MG; Hurst SD; Lim HB; Swaminathan S; Laimins LA  
Abrogation of a mitotic checkpoint by E2 proteins from oncogenic human papillomaviruses correlates with increased turnover of the p53 tumor suppressor protein.  
*EMBO J* 16: 318-31 (1997)
- Gopalakrishnan R; Weghorst CM; Lehman TA; Calvert RJ; Bijur G; Sabourin CL; Mallery SR; Schuller DE; Stoner GD  
Mutated and wild-type p53 expression and HPV integration in proliferative verrucous leukoplakia and oral squamous cell carcinoma.  
*Oral Surg Oral Med Oral Pathol Oral Radiol Endod* 83: 471-7 (1997)
- Herd K; Fernando GJ; Dunn LA; Frazer IH; Lambert P; Tindle RW  
E7 oncoprotein of human papillomavirus type 16 expressed constitutively in the epidermis has no effect on E7-specific B- or Th-repertoires or on the immune response induced or sustained after immunization with E7 protein.  
*Virology* 231: 155-65 (1997)
- Herrington CS  
Cervical pathology.  
*Curr Opin Obstet Gynecol* 9: 57-62 (1997)
- Hickman ES; Bates S; Vousden KH  
Perturbation of the p53 response by human papillomavirus type 16 E7.  
*J Virol* 71: 3710-8 (1997)
- Hidajat R; McNicol P  
Primer-directed mutagenesis of an intact plasmid by using Pwo DNA polymerase in long distance inverse PCR.  
*Biotechniques* 22: 32-4 (1997)
- Iwasaki T; Maeda H; Kameyama Y; Moriyama M; Kanai S; Kurata T  
Presence of a novel hamster oral papillomavirus in dysplastic lesions of hamster lingual mucosa induced by application of dimethylbenzanthracene and excisional wounding: molecular cloning and complete nucleotide sequence.  
*J Gen Virol* 78 ( Pt 5): 1087-93 (1997)
- Jacobs MV; Snijders PJ; van den Brule AJ; Helmerhorst TJ; Meijer CJ; Walboomers JM  
A general primer GP5+/GP6(+)-mediated PCR-enzyme immunoassay method for rapid detection of 14 high-risk and 6 low-risk human papillomavirus genotypes in cervical scrapings.  
*J Clin Microbiol* 35: 791-5 (1997)
- Jenson AB; Jenson MC; Cowsert L; Ghim SJ; Sundberg JP  
Multiplicity of uses of monoclonal antibodies that define papillomavirus linear immunodominant epitopes.  
*Immunol Res* 16: 115-9 (1997)
- Kataoka A; Nishida T; Sugiyama T; Hori K; Honda S; Yakushiji M  
Squamous cell carcinoma of the endometrium with human papillomavirus type 31 and without tumor suppressor gene p53 mutation.  
*Gynecol Oncol* 65: 180-4 (1997)
- Khare S; Pater MM; Tang SC; Pater A  
Effect of glucocorticoid hormones on viral gene expression, growth, and dysplastic differentiation in HPV16-immortalized ectocervical cells.  
*Exp Cell Res* 232: 353-60 (1997)
- Kubbutat MH; Vousden KH  
Proteolytic cleavage of human p53 by calpain: a potential regulator of protein stability.  
*Mol Cell Biol* 17: 460-8 (1997)
- Kyo S; Klumpp DJ; Inoue M; Kanaya T; Laimins LA  
Expression of AP1 during cellular differentiation determines human papillomavirus E6/E7 expression in stratified epithelial cells.  
*J Gen Virol* 78 ( Pt 2): 401-11 (1997)
- Lazo PA  
Papillomavirus integration: prognostic marker in cervical cancer? letter  
*Am J Obstet Gynecol* 176: 1121-2 (1997)

## References (1997)

- Lehman CW; King DS; Botchan MR  
A papillomavirus E2 phosphorylation mutant exhibits normal transient replication and transcription but is defective in transformation and plasmid retention.  
*J Virol* 71: 3652-65 (1997)
- Leng X; Ludes-Meyers JH; Wilson VG  
Isolation of an amino-terminal region of bovine papillomavirus type 1 E1 protein that retains origin binding and E2 interaction capacity.  
*J Virol* 71: 848-52 (1997)
- Liu WJ; Gissmann L; Sun XY; Kanjanahaluethai A; Muller M; Doorbar J; Zhou J  
Sequence close to the N-terminus of L2 protein is displayed on the surface of bovine papillomavirus type 1 virions.  
*Virology* 227: 474-83 (1997)
- Liu X; Han S; Baluda MA; Park NH  
HPV-16 oncogenes E6 and E7 are mutagenic in normal human oral keratinocytes.  
*Oncogene* 14: 2347-53 (1997)
- Ludmerer SW; Benincasa D; Mark GE 3rd; Christensen ND  
A neutralizing epitope of human papillomavirus type 11 is principally described by a continuous set of residues which overlap a distinct linear, surface-exposed epitope.  
*J Virol* 71: 3834-9 (1997)
- Marionnet AV; Lizard G; Chardonnet Y; Schmitt D  
Comparative evaluation of the antiproliferative effect of cyclosporin A and gamma-interferon on normal and HPV-transformed keratinocytes by cell counting, MTT assay and tritiated thymidine incorporation.  
*Cell Biol Toxicol* 13: 115-23 (1997)
- Massimi P; Banks L  
Repression of p53 transcriptional activity by the HPV E7 proteins.  
*Virology* 227: 255-9 (1997)
- Matsumoto T; Nakashima N; Takase K; Hirochika H; Mizuno H  
A mutation study of the DNA binding domain of human papillomavirus type 11 E2 protein.  
*J Biochem (Tokyo)* 121: 138-44 (1997)
- Maurer TA; Christian KV; Kerschmann RL; Berzin B; Palefsky JM; Payne D; Tying SK; Berger TG  
Cutaneous squamous cell carcinoma in human immunodeficiency virus-infected patients. A study of epidemiologic risk factors, human papillomavirus, and p53 expression.  
*Arch Dermatol* 133: 577-83 (1997)
- McShan GD; Wilson VG  
Casein kinase II phosphorylates bovine papillomavirus type 1 E1 in vitro at a conserved motif.  
*J Gen Virol* 78 ( Pt 1): 171-7 (1997)
- Melero I; Singhal MC; McGowan P; Haugen HS; Blake J; Hellstrom KE; Yang G; Clegg CH; Chen L  
Immunological ignorance of an E7-encoded cytolytic T-lymphocyte epitope in transgenic mice expressing the E7 and E6 oncogenes of human papillomavirus type 16.  
*J Virol* 71: 3998-4004 (1997)
- Mitsuishi T; Sata T; Matsukura T; Iwasaki T; Kawashima M  
The presence of mucosal human papillomavirus in Bowen's disease of the hands.  
*Cancer* 79: 1911-7 (1997)
- Morozov A; Shiyanov P; Barr E; Leiden JM; Raychaudhuri P  
Accumulation of human papillomavirus type 16 E7 protein bypasses G1 arrest induced by serum deprivation and by the cell cycle inhibitor p21.  
*J Virol* 71: 3451-7 (1997)
- Muller F; Giroglou T; Sapp M  
Characterization of the DNA-binding activity of the E1 and E2 proteins and the E1/E2 complex of human papillomavirus type 33.  
*J Gen Virol* 78 ( Pt 4): 911-5 (1997)
- Musiani M; Zerbini M; Venturoli S; Gentilomi G; Gallinella G; Manaresi E; La Placa M; D'Antuono A; Roda A; Pasini P  
Sensitive chemiluminescence in situ hybridization for the detection of human papillomavirus genomes in biopsy specimens.  
*J Histochem Cytochem* 45: 729-35 (1997)
- Nakao Y; Yang X; Yokoyama M; Ferenczy A; Tang SC; Pater MM; Pater A  
Induction of p16 during immortalization by HPV 16 and 18 and not during malignant transformation.  
*Br J Cancer* 75: 1410-6 (1997)
- Ned R; Allen S; Vande Pol S  
Transformation by bovine papillomavirus type 1 E6 is independent of transcriptional activation by E6.  
*J Virol* 71: 4866-70 (1997)
- Ngan HY; Tsao SW; Liu SS; Stanley M  
Abnormal expression and mutation of p53 in cervical cancer—a study at protein, RNA and DNA levels.  
*Genitourin Med* 73: 54-8 (1997)

- Odunsi KO; Ganesan TS  
The roles of the human major histocompatibility complex and human papillomavirus infection in cervical intraepithelial neoplasia and cervical cancer.  
*Clin Oncol (R Coll Radiol)* 9: 4-13 (1997)
- Olshan AF; Weissler MC; Pei H; Conway K; Anderson S; Fried DB; Yarbrough WG  
Alterations of the p16 gene in head and neck cancer: frequency and association with p53, PRAD-1 and HPV.  
*Oncogene* 14: 811-8 (1997)
- Park JS; Hwang ES; Park SN; Ahn HK; Um SJ; Kim CJ; Kim SJ; Namkoong SE  
Physical status and expression of HPV genes in cervical cancers.  
*Gynecol Oncol* 65: 121-9 (1997)
- Pattison S; Skalnik DG; Roman A  
CCAAT displacement protein, a regulator of differentiation-specific gene expression, binds a negative regulatory element within the 5' end of the human papillomavirus type 6 long control region.  
*J Virol* 71: 2013-22 (1997)
- Pepinsky RB; Prakash SS; Corina K; Gressel MJ; Barsoum J; Androphy EJ  
Sequences flanking the core DNA-binding domain of bovine papillomavirus type 1 E2 contribute to DNA-binding function.  
*J Virol* 71: 828-31 (1997)
- Phillips AC; Vousden KH  
Analysis of the interaction between human papillomavirus type 16 E7 and the TATA-binding protein, TBP.  
*J Gen Virol* 78 ( Pt 4): 905-9 (1997)
- Pise-Masison CA; Dittmer J; Clemens KE; Brady JN  
Physical and functional interaction between the human T-cell lymphotropic virus type 1 Tax1 protein and the CCAAT binding protein NF-Y.  
*Mol Cell Biol* 17: 1236-43 (1997)
- Puranen MH; Yliskoski MH; Saarikoski SV; Syrjanen KJ; Syrjanen SM  
Exposure of an infant to cervical human papillomavirus infection of the mother is common.  
*Am J Obstet Gynecol* 176: 1039-45 (1997)
- Quinn TR; Young RH  
Epidermolytic hyperkeratosis in the lower female genital tract: an uncommon simulant of mucocutaneous papillomavirus infection—a report of two cases.  
*Int J Gynecol Pathol* 16: 163-8 (1997)
- Roberts S; Ashmole I; Rookes SM; Gallimore PH  
Mutational analysis of the human papillomavirus type 16 E1-E4 protein shows that the C terminus is dispensable for keratin cytoskeleton association but is involved in inducing disruption of the keratin filaments.  
*J Virol* 71: 3554-62 (1997)
- Rosl F; Das BC; Lengert M; Geletneký K; zur Hausen H  
Antioxidant-induced changes of the AP-1 transcription complex are paralleled by a selective suppression of human papillomavirus transcription.  
*J Virol* 71: 362-70 (1997)
- Rugge M; Bovo D; Busatto G; Parenti AR; Fawzy S; Guido M; Ancona E; Ninfo V; Ruol A; Shiao YH  
p53 alterations but no human papillomavirus infection in preinvasive and advanced squamous esophageal cancer in Italy.  
*Cancer Epidemiol Biomarkers Prev* 6: 171-6 (1997)
- Savelieva E; Belair CD; Newton MA; DeVries S; Gray JW; Waldman F; Reznikoff CA  
20q gain associates with immortalization: 20q13.2 amplification correlates with genome instability in human papillomavirus 16 E7 transformed human uroepithelial cells.  
*Oncogene* 14: 551-60 (1997)
- Scholes AG; Liloglou T; Snijders PJ; Hart CA; Jones AS; Woolgar JA; Vaughan ED; Walboomers JM; Field JK  
p53 mutations in relation to human papillomavirus type 16 infection in squamous cell carcinomas of the head and neck.  
*Int J Cancer* 71: 796-9 (1997)
- Shan L; Nakamura Y; Nakamura M; Zhang Z; Jing X; Hara T; Yokoi T; Kakudo K  
Synchronous and metachronous multicentric squamous cell carcinomas in the upper aerodigestive tract.  
*Pathol Int* 47: 68-72 (1997)
- Shino Y; Shirasawa H; Kinoshita T; Simizu B  
Human papillomavirus type 16 E6 protein transcriptionally modulates fibronectin gene expression by induction of protein complexes binding to the cyclic AMP response element.  
*J Virol* 71: 4310-8 (1997)
- Simpson S; Woodworth CD; DiPaolo JA  
Altered expression of Erg and Ets-2 transcription factors is associated with genetic changes at 21q22.2-22.3 in immortal and cervical carcinoma cell lines.  
*Oncogene* 14: 2149-57 (1997)
- Solinas-Toldo S; Durst M; Lichter P  
Specific chromosomal imbalances in human papillomavirus-transfected cells during progression toward immortality.  
*Proc Natl Acad Sci U S A* 94: 3854-9 (1997)

## References (1997)

- Steger G; Corbach S  
Dose-dependent regulation of the early promoter of human papillomavirus type 18 by the viral E2 protein.  
*J Virol* 71: 50-8 (1997)
- Stellato G; Paavonen J; Nieminen P; Hibma M; Vilja P; Lehtinen M  
Diagnostic phase antibody response to the human papillomavirus type 16 E2 protein is associated with successful treatment of genital HPV lesions with systemic interferon alpha-2b.  
*Clin Diagn Virol* 7: 167-72 (1997)
- Stoppler H; Hartmann DP; Sherman L; Schlegel R  
The human papillomavirus type 16 E6 and E7 oncoproteins dissociate cellular telomerase activity from the maintenance of telomere length.  
*J Biol Chem* 272: 13332-7 (1997)
- Swan DC; Tucker RA; Holloway BP; Icenogle JP  
A sensitive, type-specific, fluorogenic probe assay for detection of human papillomavirus DNA.  
*J Clin Microbiol* 35: 886-91 (1997)
- Thain A; Webster K; Emery D; Clarke AR; Gaston K  
DNA binding and bending by the human papillomavirus type 16 E2 protein. Recognition of an extended binding site.  
*J Biol Chem* 272: 8236-42 (1997)
- Tong X; Howley PM  
The bovine papillomavirus E6 oncoprotein interacts with paxillin and disrupts the actin cytoskeleton.  
*Proc Natl Acad Sci U S A* 94: 4412-7 (1997)
- Turner JR; Shen LH; Crum CP; Dean PJ; Odze RD  
Low prevalence of human papillomavirus infection in esophageal squamous cell carcinomas from North America: analysis by a highly sensitive and specific polymerase chain reaction-based approach.  
*Hum Pathol* 28: 174-8 (1997)
- Uchiyama M; Iwasaka T; Matsuo N; Hachisuga T; Mori M; Sugimori H  
Correlation between human papillomavirus positivity and p53 gene overexpression in adenocarcinoma of the uterine cervix.  
*Gynecol Oncol* 65: 23-9 (1997)
- Vernon SD; Unger ER; Miller DL; Lee DR; Reeves WC  
Association of human papillomavirus type 16 integration in the E2 gene with poor disease-free survival from cervical cancer.  
*Int J Cancer* 74: 50-6 (1997)
- Vernon SD; Unger ER; Miller DL; Lee DR; Reeves WC  
Association of human papillomavirus type 16 integration in the E2 gene with poor disease-free survival from cervical cancer.  
*Int J Cancer* 74: 50-6 (1997)
- Villa LL  
Human papillomaviruses and cervical cancer.  
*Adv Cancer Res* 71: 321-41 (1997)
- Voet M; Defoor E; Verhasselt P; Riles L; Robben J; Volckaert G  
The sequence of a nearly unclonable 22.8 kb segment on the left arm chromosome VII from *Saccharomyces cerevisiae* reveals ARO2, RPL9A, TIP1, MRF1 genes and six new open reading frames.  
*Yeast* 13: 177-82 (1997)
- Wan TS; Chan LC; Ngan HY; Tsao SW  
t(High) frequency of telomeric associations in human ovarian surface epithelial cells transformed by human papilloma viral oncogenes.  
*Cancer Genet Cytogenet* 95: 166-72 (1997)
- Yamada T; Manos MM; Peto J; Greer CE; Munoz N; Bosch FX; Wheeler CM  
Human papillomavirus type 16 sequence variation in cervical cancers: a worldwide perspective.  
*J Virol* 71: 2463-72 (1997)
- Yasugi T; Benson JD; Sakai H; Vidal M; Howley PM  
Mapping and characterization of the interaction domains of human papillomavirus type 16 E1 and E2 proteins.  
*J Virol* 71: 891-9 (1997)
- Yu Y; Li CY; Little JB  
Abrogation of p53 function by HPV16 E6 gene delays apoptosis and enhances mutagenesis but does not alter radiosensitivity in TK6 human lymphoblast cells.  
*Oncogene* 14: 1661-7 (1997)
- Zanardi TA; Stanley CM; Saville BM; Spacek SM; Lentz MR  
Modulation of bovine papillomavirus DNA replication by phosphorylation of the viral E1 protein.  
*Virology* 228: 1-10 (1997)
- Zehbe I; Hacker GW; Su H; Hauser-Kronberger C; Hainfeld JF; Tubbs R  
Sensitive in situ hybridization with catalyzed reporter deposition, streptavidin-Nanogold, and silver acetate autometallography: detection of single-copy human papillomavirus.  
*Am J Pathol* 150: 1553-61 (1997)
- Zehbe I; Wilander E  
Human papillomavirus infection and invasive cervical neoplasia: a study of prevalence and morphology.  
*J Pathol* 181: 270-5 (1997)



## References (1997)

Zehbe I; Wilander E  
Nonisotopic ELISA-based detection of human papilloma-  
virus-amplified DNA.  
Mod Pathol 10: 188-91 (1997)